

This article was downloaded by:

On: 27 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### Utility of Nitriles in Synthesis of Pyrido[2,3-d]pyrimidines, Thiazolo[3,2-a]pyridines, Pyrano[2,3-b]benzopyrrole, and Pyrido[2,3-d]benzopyrroles

M. R. Mahmoud<sup>a</sup>; E. A. A. El-Bordany<sup>a</sup>; N. F. Hassan<sup>a</sup>; F. S. M. Abu El-Azm<sup>a</sup>

<sup>a</sup> Chemistry Department, Faculty of Science, Ain Shams University, Cairo, Egypt

**To cite this Article** Mahmoud, M. R. , El-Bordany, E. A. A. , Hassan, N. F. and El-Azm, F. S. M. Abu(2007) 'Utility of Nitriles in Synthesis of Pyrido[2,3-d]pyrimidines, Thiazolo[3,2-a]pyridines, Pyrano[2,3-b]benzopyrrole, and Pyrido[2,3-d]benzopyrroles', Phosphorus, Sulfur, and Silicon and the Related Elements, 182: 11, 2507 — 2521

**To link to this Article:** DOI: 10.1080/10426500701506465

**URL:** <http://dx.doi.org/10.1080/10426500701506465>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Utility of Nitriles in Synthesis of Pyrido[2,3-d]pyrimidines, Thiazolo[3,2-a]pyridines, Pyrano[2,3-b]benzopyrrole, and Pyrido[2,3-d]benzopyrroles

M. R. Mahmoud  
E. A. A. El-Bordany  
N. F. Hassan  
F. S. M. Abu El-Azm

Chemistry Department, Faculty of Science, Ain Shams University,  
Cairo, Egypt

*3-Aryl-2-cyanoacrylonitriles (1a,b) reacted with 6-aminothiouracil, 2-ethoxy carbonylmethyl-4-oxo- $\Delta^2$ -thiazoline, 2-cyanomethyl-4-oxo- $\Delta^2$ -thiazoline and indan-2-one to give pyrido[2,3-d]pyrimidines, thiazolo[3,2-a]pyridines, pyrano [2,3-b]benzopyrrole and pyrido[2,3-b]benzopyrrole, respectively. The IR, MS and  $^1\text{H-NMR}$  of the synthesized compounds were discussed. The antimicrobial activity of some of the synthesized compounds was tested.*

**Keywords** Pyrido[2,3-d]pyrimidines; thiazolopyridines; pyrano-; pyridobenzopyrroles; antimicrobial activity

## INTRODUCTION

Pyrido[2,3-d]pyrimidines stand out for their antitumor,<sup>1</sup> anticonvulsive,<sup>2</sup> antiasthmatic, antiallergic,<sup>3</sup> antihypertensive,<sup>4</sup> and useful as diuretic compound.<sup>5–9</sup> Previously, we have reported several new synthesis of fused heterocyclic compounds utilizing laboratory available activated nitrile derivatives as starting materials.<sup>10–20</sup> In the present article, we report on the utility of the substituted acrylonitriles for synthesis of several pyrido[2,3-d]pyrimidine, thiazolo[3,2-a]pyridines with the aim of finding new chemotherapeutic agents.

## RESULTS AND DISCUSSION

The reaction of 3-aryl-2-cyanoacrylonitrile **1a,b** with 6-aminothiouracil **2** in ethanol in the presence of catalytic amount of piperidine


Received February 10, 2007; accepted March 4, 2007.

Address correspondence to F.S.M. Abu El-Azm, Chemistry Department, Faculty of Science, Ain Shams University, Abbassia 11566, Cairo, Egypt. E-mail: ftmsaber@yahoo.com

afforded the 1:1 adduct, the structure of the adduct was established as pyrido[2,3-d]pyrimidine derivative **3** rather than pyrano[2,3-d]pyrimidine or thiazino[3,2-a]pyrimidine on the basis of spectroscopic data which revealed a pattern completely in accord with structure **3**. Thus, the IR spectrum shows the stretching absorption bands characteristic for  $\text{NH}_2$ ,  $\text{C}=\text{N}$ ,  $\text{C}=\text{O}$  and  $\text{C}=\text{S}$  groups at 3378, 3220, 3113, 2206, 1687, 1640, and  $1128\text{ cm}^{-1}$ , respectively, which in accord with the proposed structure **3**. Furthermore,  $^1\text{H}$ -NMR spectrum of **3b** revealed two signals at  $\delta$  12.7 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ) and 12.2 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ) characteristic for -NH-CS-NH- group (**c.f. Exp.**). Alkylation of **3b** using ethylbromoacetate or ethyl iodide in ethanol containing anhydrous sodium acetate gave the S-alkylated products **4** and **5**, respectively. The reaction of **3b** with hydrazine hydrate in boiling ethanol yielded the Sulfur-free compound **6** which also obtained upon treatment of the S-alkylated product **5** with hydrazine hydrate. The hydrazino derivative **6** was used as the key intermediate for the synthesis of polyheterocyclic compounds. Treatment of compound **6** with freshly distilled acetic anhydride afforded the pyridotriazolopyrimidine derivative **7**. When a solution of **6** in dilute hydrochloric acid was refluxed with 2,4-pentadione and/or 1-phenylbut-1,3-dione in pyridine afforded pyrazol-1-ylpyridopyrimidine derivatives **8a** and **8b**, respectively (Scheme 1).

The reaction of **1a** with 2-ethoxy carbonylmethyl-4-oxo- $\Delta^2$ -thiazoline **9a** in absolute ethanol in the presence of piperidine afforded the adduct thiazolo[3,2-a]pyridine **10a**. Treatment of **10a** with triethylorthoformate yielded thiazolo[3,2-a]pyridine derivative **11**. The reaction of **11** with benzylamine results in the formation of the thiazolopyridine derivative **12**. The reaction of 2-cyanomethyl-4-oxo- $\Delta^2$ -thiazoline **9b** with **1a** in ethanol in the presence of catalytic amount of piperidine resulted in the formation of thiazolo[3,2-a]pyridine derivative **10b** (Scheme 2).

The carbanion derived from indan-2-one in absolute ethanol upon treatment with piperidine reacts with **1a** to afford 3-arylidene indan-2-one **13a** (ethanol insoluble fraction) and pyrano[2,3-b]benzopyrrole **14** (ethanol soluble fraction). It is worthy to mention that **13a** could be obtained from the reaction of indan-2-one with 1-naphthaldehyde in boiling ethanol and pyrano benzopyrrole **14** prepared upon treatment of **13a** with malononitrile in refluxing ethanol in the presence of piperidine. Similarly, the reaction of 2-bromobenzaldehyde with indanone under the same conditions yielded the 3-(2-bromobenzylidene)indan-2-one **13b**. Compound **13b** when fused with malononitrile in the presence of ammonium acetate at  $190\text{--}200^\circ\text{C}$  afforded pyrido[2,3-b]benzopyrrole **15**. Hydrazinolysis of **13a** in refluxing ethanol afforded the azine **16** not




SCHEME 1

pyrazolobenzopyrrole derivative **17**. Treatment of compound **13a,b** with freshly distilled acetic anhydride yielded the N-acyl derivative **18a,b**. The reaction of the N-acyl derivative **18b** with  $\beta$ -cyanoethylhydrazine in ethanol yielded pyrazolo[3,4-b]indole **19** (Scheme 3).


## Biological Investigation

### Antimicrobial Activity

The antimicrobial screening of some of the synthesized compounds was done using the agar diffusion assay. This screening was performed against the Gram-negative bacteria, *Escherichia coli* ATCC 10536 and Gram-positive bacteria, *Staphylococcus aureus* ATCC 06538 in addition to the pathogenic fungi *Candida albicans* ATCC 1023 and *Aspergillus flavus*. A moderate activity was observed with compounds **7**, **10a**, **14**,



Downloaded At: 17:53 27 January 2011



Downloaded At: 17:53 27 January 2011

**TABLE I** Antimicrobial Screening Results of the Tested Compounds at 1000  $\mu\text{g/ml}$ 

Compd. no.	<i>E. coli</i>	<i>S. aureus</i>	<i>C. albicans</i>
<b>3b</b>	00	00	00
<b>4</b>	00	00	00
<b>5</b>	00	00	00
<b>7</b>	13	12	12
<b>8a</b>	00	00	00
<b>8b</b>	00	00	00
<b>10a</b>	12	12	11
<b>10b</b>	00	00	00
<b>11</b>	00	00	00
<b>12</b>	00	00	00
<b>13a</b>	00	00	00
<b>14</b>	17	15	15
<b>15</b>	16	15	15
<b>18a</b>	10	11	11
<b>18b</b>	12	13	12
<b>19</b>	11	11	9
<b>Ampicillin</b>	00	22	00
<b>Streptomycin</b>	20	21	00
<b>Nystatin</b>	00	00	22

00, no activity (inhibition zone < 7 mm); weak activity (7–10 mm); moderate activity (11–15 mm); strong activity (>15 mm). Solvent: DMSO (6 mm).

**15**, **18a**, **18b**, and **19**, which proved to possess marked activity against *E. coli*, *S. aureus*, and *C. albicans*. The inhibitory concentration was determined for each of the active compounds along with Ampicillin, Streptomycin, and Nystatin as positive controlled. No activity was detected for all the synthesized compounds towards *Aspergillus flavus*. Results are shown in Table I.

## CONCLUSION

The results of this article indicates that activated nitriles are used as the key starting materials for synthesis of biologically active fused heterocyclic compounds such as Pyrido[2,3-d]pyrimidines, Thiazolo[3,2-a]pyridines, Pyrano[2,3-b]benzopyrrole, and Pyrido[2,3-d]benzopyrroles.

## EXPERIMENTAL

All melting points are uncorrected. The infrared spectra were recorded on FTIR Maltson (infinity series) spectrometers as KBr discs. The

$^1\text{H}$ -NMR spectra were measured on Varian Gemini 200 MHz instrument with chemical shift ( $\delta$ ) expressed in ppm downfield from TMS. Mass spectra were recorded on Shimadzu GC-MS, QP 1000 EX instrument operating at 70 eV. The monitoring of the progress of all reactions and homogeneity of the synthesized compounds was carried out by TLC. TLC was runned using TLC aluminum sheets silica gel  $\text{F}_{254}$  (Merck).

### 7-Amino-6-cyano-5-aryl-4-oxo-1H,3H-pyrido[2,3-d]pyrimidin-2-thione **3a,b**

A mixture of 3-aryl-2-cyanoacrylonitrile **1a,b** (10 mmol) and 6-amino-thiouracil (1.43 g, 10 mmol) in ethanol (50 ml) containing 1 ml piperidine was heated under reflux for 2 h (TLC). The pale yellow product that deposited during reflux, was collected by suction, washed with ethanol, dried, and then recrystallized from the proper solvent to give **3a** and/or **3b**.


### 7-Amino-6-cyano-5-[1'-naphthyl]-4-oxo-1H,3H-pyrido[2,3-d]pyrimidin-2(H)-thione **3a**

Recrystallized from ethanol as pale yellow crystals, m.p.: 326–328°C, yield 46%. IR (KBr): 3378, 3220, 3113  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2206  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1687  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1640  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ) and 1128  $\text{cm}^{-1}$  ( $\text{C}=\text{S}$ ).  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 12.08 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 11.7 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.1 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ) and 8.0–7.5 (m, 7 $\text{H}_{\text{arom.}}$ ).  $^{13}\text{C}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 173.8 (CS), 160.2 (CO), 114 (CN), 102 ( $\text{C}_{4a}$ ), 153.8 ( $\text{C}_5$ ), 80.1 ( $\text{C}_6$ ), 156 ( $\text{C}_7$ ), 160.7 ( $\text{N}=\text{C}-\text{N}$ ), 124.8–138.2 (naphthyl carbons). MS  $m/z$  (%): 345 ( $\text{M}^+$ , 98.2), 344 (M-1, 100), 343 (40.3), 285 (11.6), 215 (17.9). Anal. calcd. for  $\text{C}_{18}\text{H}_{11}\text{N}_5\text{OS}$  (345): C, 62.6%; H, 3.19%; N, 20.28%; S, 9.27%. Found: C, 62.48%; H, 3.32%; N, 19.77%; S, 9.30%.

### 7-Amino-6-cyano-5-(3',4'-dimethoxyphenyl)-4-oxo-1H,3H-pyrido[2,3-d] pyrimidin-2(H)-thione **3b**

Recrystallized from ethanol as yellow crystals, m.p.: 344–346°C, yield 48%. IR (KBr): 3377, 3299, 3216  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2206  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1696  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1640  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ), and 1128  $\text{cm}^{-1}$  ( $\text{C}=\text{S}$ ).  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 12.7 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 12.2 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.7 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.04–6.8 (m, 3 $\text{H}_{\text{arom.}}$ ), 3.84 (s, 3H, OMe) and 3.74 (s, 3H, OMe).

$^{13}\text{C}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 174.7 (CS), 161.3 (CO), 158.4 ( $\text{C}_5$ ), 82.4 ( $\text{C}_6$ ), 112.6 (CN), 160.1 ( $\text{C}_7$ ), 102 ( $\text{C}_{4a}$ ), 155.3 ( $\text{N}=\text{C}-\text{N}$ ), 132 ( $\text{C}_{8a}$ ), 158.9



(C-OMe), 160.2 (C-OMe), 56.1 (OMe), 55.7 (OMe), 121.8, 122.9, 126.3 ( $C_{\text{arom.}}$ ). MS  $m/z$  (%): 355 ( $M^+$ , 100). Anal. calcd. for  $C_{16}H_{13}N_5O_3S$  (355): C, 54.08%; H, 3.6%; N, 19.7%; S, 9.0%. Found: C, 52.86%; H, 4.0%; N, 20.09%; S, 8.66%.

#### 7-Amino-6-cyano-5-(3',4'-dimethoxyphenyl)-2-ethoxycarbonylmethylthiopyrido[2,3-d]pyrimidin-4(3H)-one 4

A mixture of **3b** (3.5 g, 10 mmol), ethylbromoacetate (1.67 ml, 10 mmol), and anhydrous sodium acetate (5 g, 160 mmol) in absolute ethanol (50 ml) was refluxed for 6 h (TLC). Most of the solvent was distilled and the reaction mixture was poured onto water to afford a white substance which was filtered off, washed several times with hot water, dried and recrystallized from dioxane to give **4**. m.p.: 321–323°C, yield 85%. IR (KBr): 3478, 3389, 3214, 3144  $\text{cm}^{-1}$  ( $\text{NH}_2$ , NH), 2212, 2182  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1737  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$  ester), 1669  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$  pyrimidone), and 1630  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ).  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$  (ppm): 12.5 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.6 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.03–6.8 (m, 3 $H_{\text{arom.}}$ ), 4.2 (q, 2H,  $\text{COOCH}_2\text{CH}_3$ ), 3.8 (s, 2H,  $\text{SCH}_2\text{COOEt}$ ), 3.7 (s, 6H, 2OMe) and 1.2 (t, 3H,  $\text{COOCH}_2\text{CH}_3$ ).  $^{13}\text{C-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$  (ppm): 158.7 ( $\text{C}_2$ ), 162.1 ( $\text{C}_4$ ), 99.3 ( $\text{C}_{4a}$ ), 155.2 ( $\text{C}_5$ ), 89.7 ( $\text{C}_6$ ), 161.7 ( $\text{C}_7$ ), 158.1 ( $\text{C}_{8a}$ ), 114.9 (CN), 132.4 ( $\text{C}'_1$ ), 126.3 ( $\text{C}'_2$ ), 156 ( $\text{C}'_3$ ), 150.9 ( $\text{C}'_4$ ), 120.3 ( $\text{C}'_5$ ), 124.3 ( $\text{C}'_6$ ), 55.8 (OMe), 55.7 (OMe), 32.4 ( $\text{SCH}_2$ ), 171.3 ( $\text{CO}_{\text{ester}}$ ), 57.7 ( $\text{OCH}_2$ ), 14.8 (Me). MS  $m/z$  (%): 441 ( $M^+$ , 73.8), 396 (33.6), 368 (87.9), 355 (100), 323 (17.4). Anal. calcd. for  $\text{C}_{20}\text{H}_{19}\text{N}_5\text{O}_5\text{S}$  (441): C, 54.4%; H, 4.3%; N, 15.8%; S, 7.2%. Found: C, 54.43%; H, 4.52%; N, 16.31%; S, 6.99%.

#### 7-Amino-6-cyano-5-(3',4'-dimethoxyphenyl)-2-ethylthiopyrido[2,3-d]pyrimidin-4(3H)-one 5

A mixture of **3b** (3.5 g, 10 mmol), ethyl iodide (1.5 ml, 10 mmol) and anhydrous sodium acetate (5 g, 160 mmol) in absolute ethanol (50 ml)



was heated under reflux for 3 h (TLC). Most of the solvent was distilled and the reaction mixture was stirred with hot water to give a white substance which was filtered off, dried and recrystallized from dioxane to give **5**. m.p.: 361–363°C, yield 90%. IR (KBr): 3449, 3301, 3217  $\text{cm}^{-1}$  ( $\text{NH}_2$ , NH), 2215  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1697  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1622  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ).  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 12.2 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.7 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.0–6.8 (m, 3 $\text{H}_{\text{arom.}}$ ), 3.84 (s, 3H, OMe), 3.74 (s, 3H, OMe), 3.2 (q, 2H,  $\text{SCH}_2\text{CH}_3$ ) and 1.3 (t, 3H,  $\text{SCH}_2\text{CH}_3$ ). MS  $m/z$  (%): 383 ( $\text{M}^+$ , 48), 355 (100), 340 (28.1). Anal. calcd. for  $\text{C}_{18}\text{H}_{17}\text{N}_5\text{O}_3\text{S}$  (383): C, 56.3%; H, 4.4%; N, 18.2%; S, 8.3%. Found: C, 56.1%; H, 4.8%; N, 17.68%; S, 8.0%.

### 7-Amino-6-cyano-2-hydrazino-5-(3',4'-dimethoxyphenyl)-pyrido[2,3-d]pyrimidin-4(3H)-one **6**

#### Method 1

A mixture of **3b** (3.5 g, 10 mmol) and hydrazine hydrate 80% (0.5 ml, 10 mmol) in absolute ethanol (50 ml) was heated under reflux for 3 h. The yellowish white product that deposited during reflux, was collected by suction and washed with ethanol to give **6**. m.p.: >360°C, yield 20%. IR (KBr): 3451, 3303, 3208  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2215  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1691  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ) $\delta$  (ppm): 10.9 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 8.4 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 8.1 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.3 (s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.2–6.9 (m, 3 $\text{H}_{\text{arom.}}$ ), 3.83 (s, 3H, OMe), and 3.79 (s, 3H, OMe). MS  $m/z$  (%): 353 ( $\text{M}^+$ , 100), 323 (40), 308 (31.2), 280 (23.1). Anal. calcd. for  $\text{C}_{16}\text{H}_{15}\text{N}_7\text{O}_3$  (353): C, 54.3%; H, 4.2%; N, 27.7%. Found: C, 54.71%; H, 3.93%; N, 28.21%.

#### Method 2

A mixture of **5** (3.83 g, 10 mmol) and hydrazine hydrate 80% (0.5 ml, 10 mmol) in absolute ethanol (50 ml) was heated under reflux for 3 h. The excess solvent was removed by suction. The deposited solid was filtered off and washed by ethanol to give **6** (yield 80%).

### 8-Diacetylamino-7-cyano-6-(3',4'-dimethoxyphenyl)-3-methyl-5-oxo-1H,1,2,4-triazolo[4,5-a]pyrido[2,3-d]pyrimidine **7**

A solution of **6** (1 g, 2.8 mmol) in freshly distilled acetic anhydride (10 ml) was heated under reflux for 6 h, left to cool, and diluted with cold water. The crude pale green solid that separated out was collected, washed with cold water, dried and then recrystallized from ethanol to give **7**. m.p.: 195–196°C, yield 82%. IR (KBr): 2229  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1740,

1704, 1691  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm): 8.9 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 6.9–6.8 (m,  $3\text{H}_{\text{arom.}}$ ), 3.95 (s, 3H, OMe), 3.88 (s, 3H, OMe), 2.5 (s, 3H,  $\text{COCH}_3$ ), 2.4 (s, 3H,  $\text{COCH}_3$ ) and 2.2 (s, 3H,  $\text{N}=\text{C}-\text{CH}_3$ ). MS  $m/z$  (%): 461 ( $\text{M}^+$ , 21.3), 419 (100), 404 (41.5), 377 (78.1), 362 (33.6). Anal. calcd. for  $\text{C}_{22}\text{H}_{19}\text{N}_7\text{O}_5$  (461): C, 57.2%; H, 4.1%; N, 21.2%. Found: C, 56.76%; H, 3.81%; N, 21.7%.

**7-Amino-6-cyano-5-(3',4'-dimethoxyphenyl)-2-(3'',5''-dimethylpyrazolo-1-yl)-3H-pyrido[2,3-d]pyrimidin-4(3H)-one **8a** and 7-Amino-6-cyano-5-(3', 4'-dimethoxyphenyl)-2-(3''-methyl-1-phenylpyrazolo-2''-yl)-3H-pyrido-[2,3-d]pyrimidin-4(3H)-one **8b****

A solution of **6** (1 g, 2.8 mmol) in dilute hydrochloric acid (10 ml, 50%) was heated under reflux with 2,4-pentadione (0.28 ml, 2.8 mmol) or 1-phenylbut-1,3-dione (0.45 g, 2.8 mmol) in pyridine (10 ml) for 0.5 h. The pale yellow product deposited on hot, was collected by suction, dried and then recrystallized from dioxane to give **8a** and **8b**, respectively.

**8a.** m.p.: 342–344°C, yield 76%. IR (KBr): 3459, 3301, 3211  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2213  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1682  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ) $\delta$  (ppm): 11.45 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.68 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.0–6.8 (m,  $3\text{H}_{\text{arom.}}$ ), 6.3 (s, 1H,  $=\text{CH}$ ), 3.86 (s, 3H, OMe), 3.76 (s, 3H, OMe), 2.6 (s, 3H,  $\text{C}'_5-\text{CH}_3$ ), and 2.25 (s, 3H,  $\text{C}'_3-\text{CH}_3$ ). MS  $m/z$  (%): 417 ( $\text{M}^+$ , 100), 402 (50.7), 95 (18.2). Anal. calcd. for  $\text{C}_{21}\text{H}_{19}\text{N}_7\text{O}_3$  (417): C, 60.4%; H, 4.5%; N, 23.5%. Found: C, 59.33%; H, 4.5%; N, 23.9%.

**8b.** m.p.: 311–313°C, yield 80%. IR (KBr): 3484, 3336, 3289  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2208  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1698  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ) $\delta$  (ppm): 12.03 (s, 1H, NH, exchangeable with  $\text{D}_2\text{O}$ ), 7.74 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 7.5–7.0 (m,  $8\text{H}_{\text{arom.}}$ ), 6.7 (s, 1H), 3.87 (s, 3H, OMe), 3.77 (s, 3H, OMe), and 2.52 (s, 3H,  $\text{C}'_3-\text{Me}$ ). MS  $m/z$  (%): 479 ( $\text{M}^+$ , 100), 464 (28.2), 157 (6.9). Anal. calcd. for  $\text{C}_{26}\text{H}_{21}\text{N}_7\text{O}_3$  (479): C, 65.1%; H, 4.38%; N, 20.4%. Found: C, 64.77%; H, 4.41%; N, 20.91%.

**5-Amino-7-aryl-2-arylidene-6-cyano-8-substituted-3-oxo-2H,3H,7H-thiazolo-[3,2-a]-pyridine **10a,b****

A mixture of **1a** (4.08 g, 20 mmol) and 2-ethoxycarbonylmethyl-4-oxo- $\Delta^2$ -thiazoline **9a** and/or 2-cyanomethyl-4-oxo- $\Delta^2$ -thiazoline **9b** (10 mmol) in absolute ethanol (50 ml) containing 1 ml piperidine was heated under reflux for 1 h. The yellow product that deposited during reflux,

was collected by suction, washed by ethanol, dried and recrystallized from the proper solvent to give **10a** and **10b**, respectively.

**5-Amino-6-cyano-8-ethoxycarbonyl-3-oxo-2-[1-naphthylidene]-7-[1-naphth-yl]-2H,3H,7H-thiazolo[3,2-a]pyridine 10a**

Recrystallized from dioxane to give a yellow crystals, m.p.: 265–266°C, yield 63%. IR (KBr): 3457, 3347  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2196  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1691  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$  unsaturated ester), 1643  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$  unsaturated cyclic imide).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm): 8.6–7.6 (m, 14 $\text{H}_{\text{arom.}}$ ), 7.3 (br.s, 2H,  $\text{NH}_2$  exchangeable with  $\text{D}_2\text{O}$ ), 6.6 (s, 1H, olefinic proton), 4.8 (s, 1H,  $\text{C}_7\text{-H}$ ), 3.89 (q, 2H,  $\text{COOCH}_2\text{CH}_3$ ), and 0.98 (t, 3H,  $\text{COOCH}_2\text{CH}_3$ ). MS  $m/z$  (%): 529 ( $\text{M}^+$ , 10.3), 402 (24), 403 (100), 152 (12.8). Anal. calcd. for  $\text{C}_{32}\text{H}_{23}\text{N}_3\text{O}_3\text{S}$  (529): C, 72.5%; H, 4.3%; N, 7.9%; S, 6.0%. Found: C, 73.06%; H, 4.09%; N, 8.32%; S, 5.49%.

**5-Amino-6,8-dicyano-3-oxo-2-(1-naphthylidene)-7-(1-naphthyl)-2H,3H,7H-thiazolo[3,2-d]pyridine 10b**

Recrystallized from dioxane as yellow crystals, m.p.: 290–291°C, yield 70%. IR (KBr): 3387, 3331, 3287  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2196  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1717  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1653  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ).  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ) $\delta$  (ppm): 8.81–7.4 (m, 14 $\text{H}_{\text{arom.}}$ ), 7.1 (s, 1H, olefinic proton), 6.5 (br.s, 2H,  $\text{NH}_2$  exchangeable with  $\text{D}_2\text{O}$ ) and 4.6 (s, 1H,  $\text{C}_7\text{-H}$ ). MS  $m/z$  (%): 355 ( $\text{M}^+ - \text{C}_{10}\text{H}_7$ , 19.4), 301 (100), 184 (62.2). Anal. calcd. for  $\text{C}_{30}\text{H}_{18}\text{N}_4\text{OS}$  (482): C, 74.6%; H, 3.7%; N, 11.6%; S, 6.6%. Found: C, 74.26%; H, 3.63%; N, 11.24%; S, 6.72%.

**6-Cyano-5-ethoxymethyleneamino-8-ethoxycarbonyl-3-oxo-2-(1-naphthylidene)-7-(1-naphthyl)-2H,3H,7H-thiazolo[3,2-a]pyridine 11**

A mixture of **10a** (1 g, 1.9 mmol) and triethylorthoformate (10 ml, 67 mmol) in freshly distilled acetic anhydride (10 ml) was heated under reflux for 8 h. The pale green solid deposited during reflux, was collected by filtration, washed with light-petroleum ether (b.p. 80–100) and, then, recrystallized from benzene to give **11**. m.p.: 236–238°C, yield 57%. IR (KBr): devoid  $\nu_{\text{NH}_2}$ , 2205  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1723  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$  ester), 1690  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$   $\alpha\beta$ -unsaturated ketone), 1626  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm): 8.5–7.3 (m, 15 $\text{H}_{\text{arom.}}$  +  $\text{N}=\text{CH}$ ), 7.2 (s, 1H,  $=\text{CH}$ ), 5.7 (s, 1H,  $\text{C}_7\text{-H}$ ), 4.4 (q, 2H,  $\text{CH}_3\text{CH}_2\text{-OCO}$ ), 3.9 (q, 2H,  $\text{CH}_3\text{CH}_2\text{-O}$ ), 1.5 (t, 3H,  $\text{CH}_3\text{CH}_2\text{-O-CO}$ ) and 0.7 (t, 3H,  $\text{CH}_3\text{CH}_2\text{O}$ ). MS  $m/z$  (%): 585 ( $\text{M}^+$ ,

12.1), 458 (100), 312 (10.1), 184 (16). Anal. calcd. for  $C_{35}H_{27}N_3O_4S$  (585): C, 71.79%; H, 4.6%; N, 7.17%; S, 5.47%. Found: C, 71.56%; H, 4.23%; N, 7.62%; S, 5.81%.

### Formation of the Thiazolo Pyridine Derivative 12

A mixture of **12** (5.85 g, 10 mmol) and benzylamine (1 ml, 10 mmol) in dry pyridine (30 ml) was heated under reflux for 6 h. The reaction mixture was cooled and acidified with cold dilute acetic acid. The solid deposited was filtered off, washed several times with water and dried, then recrystallized from dioxane to give **12**. m.p.: 257–259°C, yield 80%. IR (KBr): 3450, 3341  $\text{cm}^{-1}$  (NH), 2194  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1694  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1643  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm): 9.1 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), 8.5–7.4 (m, 24 $\text{H}_{\text{arom.}}$ ), 7.0 (s, 1H, olefinic proton), 6.93 (s, 1H,  $\text{N}=\text{CH}$ ), 4.9 (s, 1H,  $\text{C}_7\text{-H}$ ), and 4.34 (d,d, 4H, benzylic protons). MS  $m/z$  (%): 590 [ $\text{M}^+-(\text{PhCH}_2, \text{CN})$ , 2.6], 402 (15.5), 184 (15.5), 152 (20.38), 127 (100). Anal. calcd. for  $\text{C}_{45}\text{H}_{33}\text{N}_5\text{O}_2\text{S}$  (707): C, 76.37%; H, 4.6%; N, 9.9%; S, 4. %5. Found: C, 76.71%; H, 4.3%6; N, 10.21%; S, 4.32%.

### Reaction of 1a with 2-indanone: Formation of 13a and 14

A mixture of **1a** (2.04 g, 10 mmol) and indan-2-one (1.33 g, 10 mmol) in absolute ethanol (50 ml) containing 1 ml piperidine was heated under reflux for 2 h (TLC). The reaction mixture was cooled and the solid deposited was collected by suction, washed by ethanol and recrystallized from benzene to give **13a**. Most of the solvent of the mother liquor was distilled off and then acidified with cold dilute hydrochloric acid. The green solid that separated out was collected by filtration, washed by cold water, dried and recrystallized from ethanol to give **14**.

**13a**. m.p.: 190–192°C, yield 50%. IR (KBr): 3179, 3151  $\text{cm}^{-1}$  (NH), 1710  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ), 1616  $\text{cm}^{-1}$  ( $\text{C}=\text{C}$ ).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm): 8.3 (s, 1H, NH), 8.0–7.2 (m, 11 $\text{H}_{\text{arom.}}$ ), and 6.7 (s, 1H,  $\text{CH}=\text{}$ ). MS  $m/z$  (%): 271 ( $\text{M}^+$ , 79.3), 270 (100), 144 (11.2), 120 (9). Anal. calcd. for  $\text{C}_{19}\text{H}_{13}\text{NO}$  (271): C, 84.1%; H, 4.79%; N, 5.1%. Found: C, 83.75%; H, 5.0%; N, 4.77%.

### Authentic Sample

A solution of indan-2-one (1.33 g, 10 mmol) in absolute ethanol (30 ml) was stirred with 1-naphthaldehyde (1.56 ml, 10 mmol) for  $\frac{1}{2}$  h and then refluxed for 1 h. The excess solvent was removed and left a crude

product which recrystallized from benzene to give **13a** (identity by m.p., mixed m.p. and TLC).

14. m.p.: 327–329°C, yield 20%. IR (KBr): 3429, 3317, 3211  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2210  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), 1645  $\text{cm}^{-1}$  ( $\text{C}=\text{N}$ ).  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ) $\delta$  (ppm): 10.4 (s, 1H,  $\text{NH}$ , exchangeable with  $\text{D}_2\text{O}$ ), 8.18–7.0 (m, 11 $\text{H}_{\text{arom.}}$ ), 6.7 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ), and 4.86 (s, 1H). MS  $m/z$  (%): 337 ( $\text{M}^+$ , 37.6), 334 (100), 307 (17.8). Anal. calcd. for  $\text{C}_{22}\text{H}_{15}\text{N}_3\text{O}$  (337): C, 78.3%; H, 4.45%; N, 12.46%. Found: C, 78.0%; H, 4.61%; N, 12.09%.

### Reaction of **13a** with Malononitrile


A mixture of **13a** (2.71 g, 10 mmol) and malononitrile (0.99 g, 15 mmol) in absolute ethanol (50 ml) and 1 ml piperidine was refluxed for 3 h. The excess solvent was collected by distillation and the residual acidified with cold dilute acetic acid to give **14** (yield 28.7%) (identity by m.p., mixed m.p., IR, and TLC).

### Formation of **13b**

A mixture of indan-2-one (1.33 g, 10 mmol) and 2-bromobenzaldehyde (1.85 ml, 10 mmol) in absolute ethanol (50 ml) was refluxed for 4 h. The resulting yellow product (one spot in TLC) was collected by filtration and then recrystallized from benzene to give **13b**. m.p.: 166–167°C, yield 80%. IR (KBr): 3143  $\text{cm}^{-1}$  ( $\text{NH}$ ), 1713  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ). MS  $m/z$  (%): 300 ( $\text{M}^+$ , 100), 301 (9.6), 302 (30). Anal. calcd. For  $\text{C}_{15}\text{H}_{10}\text{NOBr}$  (300): C, 60.0%; H, 3.3%; N, 4.6%; Br, 26.6%. Found: C, 59.4%; H, 3.4%; N, 4.4%; Br, 27.0%.

### 2-Amino-3-cyano-4-(2-bromophenyl)-pyrido-[2,3-b]benzopyrrole **15**

A mixture of **13b** (1 g, 3.3 mmol), malononitrile (0.22 g, 3.3 mmol) and ammonium acetate (3 g, 40 mmol) was firstly fused for 2 min, then heated under reflux in absolute ethanol (25 ml) for 6 h. Most of the solvent was distilled off and the reaction mixture was poured onto water. The pale green solid that separated out was collected by filtration, dried and then, recrystallized from benzene to give **15**. m.p.: 165–167°C, yield 30%. IR (KBr): 3475, 3331, 3217  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 2196  $\text{cm}^{-1}$  ( $\text{C}\equiv\text{N}$ ), and no band for  $\text{C}=\text{O}$  group.  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ) $\delta$  (ppm): 10.3 (s, 1H,  $\text{NH}$ , exchangeable with  $\text{D}_2\text{O}$ ), 8.1–7.3 (m, 8 $\text{H}_{\text{arom.}}$ ), and 6.88 (br.s, 2H,  $\text{NH}_2$ , exchangeable with  $\text{D}_2\text{O}$ ).  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ) $\delta$  (ppm):



MS  $m/z$  (%): 363 ( $M^+$ , 22.5), 364 (9.2),  $M+2$  (9.2), 283 (100). Anal. calcd. for  $C_{18}H_{11}N_4Br$  (363): C, 59.5%; H, 3.0%; N, 15.4%; Br, 22.0%. Found: C, 59.28%; H, 3.01%; N, 15.63%; Br, 22.46%.

### Reaction of 13a with Hydrazine Hydrate; Formation of the Azine 16

A mixture of **13a** (10 mmol) and hydrazine hydrate (1.5 mmol) in ethanol (25 ml) was heated under reflux for 2 h. The solid deposited after evaporation of the solvent, was filtered off, dried, and recrystallized from light petroleum ether (b.p. 80–100°C) to give **16**. m.p.: 63–65°C, yield 43%. MS  $m/z$  (%): 308 ( $M^+$ , 32.9), 280 (100), 181 (59), 154 (71.1), 127 (87.3). Identity (m.p, mixed m.p, TLC) with authentic sample resulted from the reaction of 1-naphthaldehyde with hydrazine hydrate in ethanol.

### Acetylation of 13a,b; Formation of 18a,b

A solution of **13a** or **13b** (1 g) in freshly distilled acetic anhydride (10 ml) was heated under reflux on water bath for 2 h. The solid deposited on hot was collected by suction, washed several times with light petroleum ether and recrystallized from the proper solvent to give **18a** and/or **18b**.

**18a.** Recrystallized from light petroleum ether (b.p. 80–100°C), m.p.: 112–114°C, yield 65%. IR (KBr): 1733, 1704  $cm^{-1}$  (C=O).  $^1H$ -NMR ( $CDCl_3$ ) $\delta$  (ppm): 8.4–7.2 (m, 11H<sub>arom.</sub>), 6.9 (d, 1H, C=CH), and 2.8 (s, 3H, COMe). MS  $m/z$  (%): 313 ( $M^+$ , 32.2), 271 (95.3), 270 (95.3), 270 (100), 242 (19.5), 144 (12.6). Anal. calcd. for  $C_{21}H_{15}NO_2$  (313): C, 80.5%; H, 4.79%; N, 4.47%. Found : C, 80.23%; H, 4.72%; N, 4.39%.

**18b.** Recrystallized from benzene, m.p.: 138–139°C, yield 90%. IR (KBr): 1734, 1708  $cm^{-1}$  (C=O).  $^1H$ -NMR ( $CDCl_3$ ) $\delta$  (ppm) : 8.8–7.6 (m, 8H<sub>arom.</sub>), 7.2 (s, 1H, C=CH) and 2.45 (s, 3H, COMe). MS  $m/z$  (%): 342 ( $M^+$ , 3.1), 262 (17.4), 220 (100). Anal. calcd. for  $C_{17}H_{12}NO_2Br$  (342): C,

59.6%; H, 3.5%; N, 4.1%; Br, 23.39%. Found : C, 59.1%; H, 3.84%; N, 4.44%; Br, 23.69%.

### 8-Acetyl-1-(2'-cyanoethyl)-3-(2'-bromophenyl)-2,3-dihydropyrazolo [3,4-b]indole 19

A solution of compound **18b** (3.42 g, 10 mmol) in pyridine (50 ml) was stirred with  $\beta$ -cyanoethylhydrazine (0.7 ml, 10 mmol) for 1 h and then refluxed for 3 h on water bath. The reaction mixture was concentrated and acidified with cold dilute acetic acid. The deposited solid was filtered off, dried and then recrystallized from ethanol to give **19** as yellow crystals. m.p.: 177–178°C, yield 40%. IR (KBr): 3242, 3183  $\text{cm}^{-1}$  (NH), 2253  $\text{cm}^{-1}$  (C $\equiv$ N), 1695  $\text{cm}^{-1}$  (C=O), 1647  $\text{cm}^{-1}$  (C=N).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$  (ppm) : 8.3–7.3 (m, 8H<sub>arom.</sub>), 5.3 (d, 1H), 4.2 (d, 1H, NH, exchangeable with D<sub>2</sub>O), 3.4 (t, 2H, N-CH<sub>2</sub>-CH<sub>2</sub>-), 3.1 (t, 2H, CH<sub>2</sub>CN) and 2.45 (s, 3H, COMe). MS  $m/z$  (%): 383 (M-CN<sup>+</sup>, 11.1), 220 (100). Anal. calcd. for C<sub>20</sub>H<sub>17</sub>N<sub>4</sub>OBr (409): C, 58.6%; H, 4.1%; N, 13.69%; Br, 19.5%. Found C, 58.32%; H, 3.81%; N, 14.02%; Br, 20.0%.

## REFERENCES

- [1] E. M. Grivsky, S. Lee, C. W. Sigel, D. S. Dush, and C. A. Nichol, *J. Med. Chem.*, **23**, 327 (1980).
- [2] E. Kretzchmar, *Pharmazie*, **35**, 253 (1980).
- [3] K. Furukawa and T. Hasegawa, Canadian Patent 2151971 (1994); *Chem. Abstr.*, **124**, 289568c (1996).
- [4] J. W. Ellingboe; U.S. Patent 5466692; *Chem. Abstr.*, **124**, 176134q (1996).
- [5] J. Osselaere and C. Lapiere, *Eur. J. Chem.*, **9**, 305 (1974).
- [6] H. Parish, R. Gillion, N. Purcell, R. Brawne, R. Spirr, and H. White, *J. Med. Chem.*, **25**, 98 (1982).
- [7] K. Nishikawa, H. Shimakawa, Y. Inada, S. Kikucki, and Y. Oka, *Chem. Pharm. Bull.*, **24**, 2057 (1976).
- [8] A. C. Valentin, E. S. Vyacheslav, A. G. Ekaterina. M. D. Sergey, and I. M. Vladimir, *Collect. Czech. Chem. Commun.*, **70**(3), 350–360 (2005).
- [9] H. Moustafa, M. F. Shibl, and R. Hilal, *J. Phosphorus, SulfurSulfur, and Silicon and Related Elements*, **180**(2), 459–478 (2005).
- [10] M. R. Mahmoud, *Synth. Commun.*, **24**(14), 2057 (1994).
- [11] R. P. Pellicciari, B. Natalini, G. Costantino, M. R. Mahmoud, I. Mattoli, and B. M. Sedeghpour, *J. Med. Chem.*, **37**(5), 647 (1994).
- [12] M. R. Mahmoud and D. B. Guirguis, *J. Phosphorus, SulfurSulfur, and Silicon and Related Elements*, **102**, 279 (1995).
- [13] M. M. Kamel, M. T. Omar, M. R. Mahmoud, H. H. Fahmy, Z. M. Nofal, and N. S. Ismail, *Egypt J. Chem.*, **39**(6), 591 (1996).
- [14] M. R. Mahmoud and H. M. F. Madkour, *Synth. Commun.*, **26**(20), 3799 (1996).
- [15] M. R. Mahmoud, E. A. Soliman, G. A. Ibrahim, and A. M. Rabie, *J. Phosphorus, Sulfur, and silicon and Related Elements*, **105**(4), 1–10 (1998).

- [16] H. M. F. Madkour, M. R. Mahmoud, M. H. Nassar, and M. M. Habashy, *Sulfur Letters*, **21**(6), 253–261 (1998).
- [17] H. M. F. Madkour, M. R. Mahmoud, M. H. Nassar, and M. M. Habashy, *J. Chinese Chem. Soc.*, **47**, 937–941 (2000).
- [18] H. M. F. Madkour, M. R. Mahmoud, A. M. Sakr, and M. M. Habashy, *J. Sci. Pharm. (Austria)*, **69**, 33 (2001).
- [19] O. A. Fathalla, M. Kamel, M. F. El-Zahar, M. R. Mahmoud, and E. M. Mohei El-Deen, *Egypt J. Chem.*, **46**(1), 135–152 (2003).
- [20] M. A. I. Salem, E. A. Soliman, M. B. Smith, M. R. Mahmoud, and M. E. Azab, *J. Phosphorus, Sulfur, and Silicon and the Related Elements*, **179**, 61–66 (2004).