

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Novel Heterocyclic P-Ligands: Synthesis and Application in Pt(II) Complexes

György Keglevich^a; Andrea Kerényi^a; Melinda Sipos^a; Annamária Balassa^a; Tamás Körtvélyesi^b

^a Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, Budapest, Hungary ^b Department of Physical Chemistry, University of Szeged, Szeged, Hungary

To cite this Article Keglevich, György , Kerényi, Andrea , Sipos, Melinda , Balassa, Annamária and Körtvélyesi, Tamás(2008) 'Novel Heterocyclic P-Ligands: Synthesis and Application in Pt(II) Complexes', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 183: 2, 440 – 444

To link to this Article: DOI: 10.1080/10426500701735544

URL: <http://dx.doi.org/10.1080/10426500701735544>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Novel Heterocyclic P-Ligands: Synthesis and Application in Pt(II) Complexes

György Keglevich,¹ Andrea Kerényi,¹ Melinda Sipos,¹
Annamária Balassa,¹ and Tamás Körtvélyesi²

¹Department of Organic Chemistry and Technology, Budapest
University of Technology and Economics, Budapest, Hungary

²Department of Physical Chemistry, University of Szeged, Szeged,
Hungary

3-Diphenylphosphino-1,2,3,6-tetrahydro- and 1,2,3,4,5,6-hexahydro-phosphinines, as well as dibenzo[c,e][1,2]oxaphosphorines (L) were prepared and used in the synthesis of the corresponding cis chelate Pt(II) complexes and PtCl₂L₂ complexes, respectively. The latter family of compounds includes 7-membered ring Pt(II) complexes.

Keywords P-heterocycles; monodentate and bidentate P-ligands; Pt(II) complexes; ring Pt(II) complexes; stereospecific ³J(P-Pt) NMR couplings


INTRODUCTION

In this article, we summarize our recent results on P-ligands and their platinum complexes. It was found earlier that the interaction of arylphospholes and dichlorodibenzonitrile platinum formed a mixture of a monophosphole and a diphosphole complex, this latter with *trans* orientation of the hetero rings.¹ As a comparison, an analogous reaction of the Mathey-phosphole led, almost exclusively, to the *cis* diphosphole complex. The difference is due to the presence or lack of steric hindrance. Other P-heterocycles, such as phosphabicyclo[3.1.0]hexanes, 1,2-dihydrophosphinines and 1,4-dihydrophosphinines were converted to Rh(III) or Pd(II) complexes.²

This project was supported by the European Union and the Hungarian State (GVOP-3.2.2.-2004-07-0006/3.0) and jointly by the Hungarian Scientific Research Fund (OTKA, Grant No. T067679).

Address correspondence to György Keglevich, Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, H-1521 Budapest, Hungary. E-mail: keglevich@mail.bme.hu


New phosphinine derivatives with exocyclic P-functions were developed by us (Scheme 1). The Michael-addition of diphenylphosphine oxide to 1-phenyl-1,2-dihydrophosphinine oxide (**1**) afforded a 3-diphenylphosphinoxido-1,2,3,6-tetrahydrophosphinine oxide (**2**)³ that was converted to the corresponding 1,2,3,4,5,6-hexahydrophosphinine oxide (**3**).⁴ Both bis(phosphine oxides) were formed in a diastereoselective manner, whose stereostructures were evaluated by DFT calculations and/or stereospecific NMR couplings. The phosphinoxido-tetrahydrophosphinine oxide (**2**) was subjected to double deoxygenation to provide the corresponding bis(phosphine) with preserved stereochemistry that on reaction with the platinum precursor led to a *cis* chelate complex (**4**).⁵ The phosphinoxido-hexahydrophosphinine oxide (**3**) was taken in a similar sequence of reactions to furnish another *cis* chelate complex (**5**) as the final outcome.⁶



SCHEME 1


On a related topic, dibenzo-oxaphosphorines with a variety of P-substituents were synthesized and used in complexations. In the first place, aryl-dibenzo-oxaphosphorines (**7**) were made available from the P-chloro derivative (**6**), and then converted to the platinum complexes containing only one heterocyclic unit. It was interesting to find, on the basis of stereospecific Pt–P couplings, that in the case of phenyl substituent a *cis* complex (**8**), while in the case of triisopropylphenyl group a *trans* complex (**9**) was formed (Scheme 2).⁷

Then a few P-alkoxy and P-amino dibenzo-oxaphosphorines (**10**) were prepared and transformed to the corresponding platinum complexes (**11**) that in this particular case contained two heterocyclic P-ligands, moreover in position *cis*.⁸ In the next stage, α -phenylethylamino- and menthyl-dibenzo-oxaphosphorines (**12** and **14**)⁸ with optical activity and




SCHEME 2

the related complexes (**13** and **15**)⁹ were synthesized (Scheme 3). In the first case, the complex (**11**) was formed in a homo- and a heterochiral form, while in the other two cases, the complexes (**13** and **15**) were obtained in two homochiral forms and a heterochiral form.




SCHEME 3

The L-prolin methyl ester served as an additional reagent for the P-substitution of the chloro-dibenzooxaphosphorine (**6**). Complexation of the prolinino-dibenzooxaphosphorine (**16**) led to the *cis* form of the bis(dibenzooxaphosphorino) platinum complex (**17**) (Scheme 4).




SCHEME 4

Ring opening reaction accompanying the substitution of the starting dibenzooxaphosphorine (**6**) furnished a new P-ligand (**18**), which on reaction with the platinum precursor gave a bis(triarylphosphino) complex (**19**) (Scheme 5). In this case, the substituents were, however, in the *trans* disposition, as suggested by stereospecific Pt–P couplings.¹⁰



SCHEME 5


Phosphorylation of the 1-hydroxy-1'-diphenylphosphino-biphenyl (**18**) with chloro-dibenzooxaphosphorine offered a newer possibility for ligand modification. Complexation of the special bidentate P-ligand (**20**) so obtained, led to a special ring Pt-complex (**21**) (Scheme 6).¹⁰




SCHEME 6

Aiming at the synthesis of further bidental P-ligands, methylaminoethanol was reacted with two equivalents of chloro-dibenzooxaphosphorine (**6**). The reaction took place smoothly and the resulting bisdibenzooxaphosphorine (**22**) was utilized in the preparation of a ring platinum complex (**23**) (Scheme 7).

L-Ephedrine was taken in a similar sequence of reactions to afford, eventually, an analogous 7-ring platinum complex (**25**) (Scheme 8).



SCHEME 7



SCHEME 8

REFERENCES

- [1] Zs. Csók, Gy. Keglevich, Gy. Petöcz, and L. Kollár, *Inorg. Chem.*, **38**, 831 (1999).
- [2] I. L. Odinet, N. M. Vinogradova, K. A. Lyssenko, D. G. Golovanov, P. V. Petrovskii, T. A. Mastryukova, H. Szelke, N. Balázsi Szabó, and Gy. Keglevich, *J. Organomet. Chem.*, **690**, 704 (2005).
- [3] Gy. Keglevich, M. Sipos, D. Szieberth, L. Nyulászi, T. Imre, K. Ludányi, and L. Töke, *Tetrahedron*, **60**, 6619 (2004).
- [4] Gy. Keglevich, M. Sipos, T. Körtvélyesi, T. Imre, and L. Töke, *Tetrahedron Lett.*, **46**, 1655 (2005).
- [5] Gy. Keglevich, M. Sipos, D. Szieberth, Gy. Petöcz, and L. Kollár, *J. Organomet. Chem.*, **689**, 3158 (2004).
- [6] Gy. Keglevich, M. Sipos, V. Ujj, and T. Körtvélyesi, *Letters in Org. Chem.*, **2**, 608 (2005).
- [7] Gy. Keglevich, H. Szelke, A. Kerényi, T. Imre, K. Ludányi, J. Dukai, F. Nagy, and P. Arányi, *Heteroatom Chem.*, **15**, 459 (2004).
- [8] Gy. Keglevich, H. Szelke, A. Kerényi, V. Kudar, M. Hanusz, K. Simon, T. Imre, and K. Ludányi, *Tetrahedron Asym.*, **16**, 4015 (2005).
- [9] Gy. Keglevich, H. Szelke, A. Kerényi, and T. Imre *Transit Metal Chem.*, **31**, 306 (2006).
- [10] Gy. Keglevich, A. Kerényi, H. Szelke, K. Ludányi, and T. Körtvélyesi, *J. Organomet. Chem.*, **691**, 5038 (2006).