

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Synthesis of New Phosphorous Ylides Containing Urethane Derivatives

Esmat Tavakolinejad^a; Naeim Ghatali^a; Mohammad Reza Islami^a; Hojat Khabazzadeh^a

^a Department of Chemistry, Shahid Bahonar, University of Kerman, Kerman, Iran

To cite this Article Tavakolinejad, Esmat , Ghatali, Naeim , Islami, Mohammad Reza and Khabazzadeh, Hojat(2008) 'Synthesis of New Phosphorous Ylides Containing Urethane Derivatives', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 183: 9, 2301 — 2308

To link to this Article: DOI: 10.1080/10426500801963574

URL: <http://dx.doi.org/10.1080/10426500801963574>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of New Phosphorous Ylides Containing Urethane Derivatives

Esmat Tavakolinejad, Naeim Ghatali, Mohammad Reza Islami, and Hojat Khabazzadeh

Department of Chemistry, Shahid Bahonar, University of Kerman, Kerman, Iran

The readily available (DMAD& DEAD) and triphenylphosphine undergo facile reaction with several urethanes such as methyl-N-phenylcarbamate, ethyl-N-phenylcarbamate, phenyl-N-phenylcarbamate, phenyl-N-(2-methylphenyl) carbamate and phenyl -N(3-methylphenyl)carbamate, to give crystalline phosphorus ylides in good yields. The overall sequence from a 1:1:1 reaction provides a simple and efficient rout to functionalized urethanes.

Keywords Acetylenic esters; phosphorus compounds; urethanes


INTRODUCTION

The urethanes have attracted considerable attention since they have been extensively used in organic synthesis polyurethanes during the past 60 years.^{1–4} Although the properties and synthesis of these compounds have been widely studied, the closely related highly functionalized urethane derivatives containing phosphorous ylide moiety are unknown so far. Recently, our interest focused on the synthesis of phosphorous ylides that could be transformed into versatile compounds.^{5–8} Our aim of this study is to prepare new urethanes containing the ylide moiety from the reaction of carbamate derivatives with acetylenic esters in the presence of triphenylphosphine as a good nucleophile.^{9–12}

Received October 24, 2007; accepted December 18, 2007.

The authors express appreciation to Shahid Bahonar University of Kerman, Faculty Research Fund for its support of this investigation.

Address correspondence to Esmat Tavakolinejad, Naeim Ghatali, Mohammad Reza Islami and Hojat Khabazzadeh, Department of Chemistry, Shahid Bahonar, University of Kerman, Kerman 76169, Iran. E-mail: etavakoly@yahoo.com.



SCHEME 1

RESULTS AND DISCUSSION

The synthesis of compounds **4a-i** was accomplished in accordance with the reaction depicted in Scheme 1. The starting compounds, carbamate derivatives **3a-i** were prepared according to the procedure, which was reported in the literature. The reaction of compounds **3a-i** with dialkyl acetylenedicarboxylate in the presence of Ph_3P gave the stable phosphorus ylides **4a-i**.

A plausible mechanism of the reaction and formation of the final products is depicted in Scheme 1. The structures of compounds **4a-i** were deduced from their high-field ^1H , ^{13}C NMR, and IR spectral data. The ^1H and ^{13}C NMR spectroscopic data for compounds **4a-i** exhibit a mixture of two rotational isomers. The ylide moiety of these compounds is strongly conjugated with the adjacent carbonyl group and rotation about the partial double bond in **4-(E)** and **4-(Z)** geometrical isomer is low on the NMR time scale at ambient temperature. The structure of rotamers, which were found in ylides, have been previously established and reported in the literature.^{13,14}

The ^1H NMR spectrum of **4a** showed six sharp lines ($\delta = 2.83, 3.0, 3.42, 3.45, 3.7, 3.86$ ppm) due to the methoxy protons, along with signals for methine protons at $\delta = 4.92$ and 4.96 which appear as two doublets ($^3J_{\text{PH}} = 16.2\text{Hz}$) and ($^3J_{\text{PH}} = 18\text{Hz}$), respectively, for major and minor geometrical isomers. The aromatic region appeared as a multiplet at $7.26\text{--}7.59$ ppm. The ^{13}C NMR spectrum of **4a** is in agreement with the mixture of two rotamers. Although the presence of the ^{31}P nucleus complicates both the ^1H and ^{13}C NMR spectra of the compounds **4a-i**

but it does help us to obtain some valuable information by long-range spin-spin coupling constants ^{31}P with ^1H and ^{13}C nuclei.

The ^1H and ^{13}C NMR spectroscopic data for compound **4b** are similar to those of **4a**, except for the ester groups, which exhibited characteristic resonances with appropriate chemical shifts.

The NMR spectral data for compounds (**4c–i**) are consistent with the phosphorane structure.

The structural assignments made for the phosphorane **4a–i** based on the ^1H , ^{13}C NMR, and their IR spectra.

The carbonyl group region of the spectrum exhibited two absorption bands for each compound, the conjugation of one ester group with the negative charge is a plausible factor in reduction of the wave number of one carbonyl absorption band.

CONCLUSION

In conclusion, we have demonstrated that the readily available acetylenedicarboxylates in the presence of triphenylphosphine undergoes facile reaction with urethane, providing a convenient and rapid synthesis of urethane derivatives containing the ylide moiety.

EXPERIMENTAL

Dimethyl acetylenedicarboxylate, diethyl acetylenedicarboxylate, phenylisocyanate, o-cresol, and m-cresol were obtained from Merck Co. and were used without further purification.

Melting points were obtained on a Gallenkamp melting point apparatus and were uncorrected. IR spectra were measured on a Mattson 1000 FT-IR spectrometer. ^1H and ^{13}C NMR spectra were reported on a BRUKER DRX-500 AVANCE spectrometer at 500.13 and 125.77 MHz, respectively.

Dimethyl-2-[(methoxycarbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4a**

Dimethyl acetylenedicarboxylate or diethyl acetylenedicarboxylate (2 mmol, 0.24 ml) was added dropwise to a magnetically stirred solution of triphenylphosphine (2 mmol, 0.53 g) and urethane (2 mmol) in a 15 ml mixture of ethyl acetate-petroleum ether (4:1) at an ambient temperature. After the addition was complete the mixture was stirred for an additional 6 h and then filtered.

The filtrate was washed thoroughly with mixture of ethyl acetate-petroleum ether to give a white powder. (0.92g, m.p. 156–158°C, yield 83%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1756, 1645, and 1617 (C=O). m.p.

Major isomer, **4a**-(E) (57.2%), ^1H NMR: δ 3.09, 3.45, and 3.77 (9H, 3s, 3OCH₃), 4.92(1H, d, $^3J_{\text{PH}} = 16.2$ Hz, P=C–CH)*, 7.26–7.59 (40H, m, arm)*. ^{13}C NMR: δ 40.87 (d, $^1J_{\text{PC}} = 135.8$ Hz, P=C), 49.35, 52.03, and 52.34(3OCH₃), 61.12(d, $^2J_{\text{PC}} = 17.0$ Hz, P=C–CH), 126.65, and 127.35 (2CH)*, 126.27 (d, $^1J_{\text{PC}} = 91.9$ Hz, C^{ipso}), 128.53 (d, $^3J_{\text{PC}} = 12.4$ Hz, C^{meta}), 131.24 (1C)*, 132.04 (d, $^4J_{\text{PC}} = 4.0$ Hz, C^{para}), 133.79 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 139.14 and 155.78 (2C), 168.4 (C=O), 169.99 (d, $^2J_{\text{PC}} = 17.8$ Hz, C=O)*, 173.50 (d, $^3J_{\text{PC}} = 3.7$ Hz, C=O)*.

Minor isomer, **4a**-(Z) (42.8%), ^1H NMR: δ 2.87, 3.42, and 3.86 (9H, 3s, 3OCH₃), ^{13}C NMR: 38.65 (d, $^1J_{\text{PC}} = 136.6$ Hz, P=C), 48.70, 52.28 and 52.43(3OCH₃), 61.96 (d, $^2J_{\text{PC}} = 16.6$ Hz, P=C–CH), 126.13 (d, $^1J_{\text{PC}} = 96.8$ Hz, C^{ipso}), 128.63 (d, $^3J_{\text{PC}} = 12.4$ Hz, C^{meta}), 131.94 (d, $^4J_{\text{PC}} = 3.60$ Hz, C^{para}), 133.56 (d, $^2J_{\text{PC}} = 9.7$ Hz, C^{ortho}), 139.41, and 155.44 (2C), 168.51 (C=O).

Diethyl-2-[(methoxycarbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4b**

(0.87g, m.p. 170–173°C, yield 75%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1750, 1651, and 1621 (C=O).

Major isomer, **4b**-(E) (51.1%), ^1H NMR: δ 0.30 and 1.33 (6H, 2t, CH₃), 3.34 (3H, s, OCH₃), 3.39, and 4.21 (4H, 2q, OCH₂-), 4.90 (1H, d, $J_{\text{PH}} = 16.4$ Hz, P=C–CH)*, 7.23–7.62 (40H, m, arm)*. ^{13}C NMR: δ 13.9 and 14.3 (2CH₃), 40.72 (d, $^1J_{\text{PC}} = 134.7$ Hz, P=C), 52.40(OCH₃), 57.37 and 60.81 (2OCH₂-), 61.12(d, $^2J_{\text{PC}} = 17.0$ Hz, P=C–CH), 126.15 (d, $^1J_{\text{PC}} = 91.3$ Hz, C^{ipso}), 127.39, and 127.56 (2C)*, 128.44 (d, $^3J_{\text{PC}} = 12.2$ Hz, C^{meta}), 131.43 (1C)*, 132.04 (d, $^4J_{\text{PC}} = 2.6$ Hz, C^{para}), 133.67 (d, $^2J_{\text{PC}} = 9.7$ Hz, C^{ortho}), 139.51, and 155.81(2C), 167.97 (C=O), 168.82 (d, $^2J_{\text{PC}} = 18.0$ Hz, C=O)*, 172.68 (d, $^3J_{\text{PC}} = 13.6$ Hz, C=O)*.

Minor isomer, **4b**-(Z) (49.9%), ^1H NMR: δ 0.94, and 1.36 (6H, 2t, CH₃), 3.37 (3H, s, OCH₃), 3.43 and 4.26 (4H, 2q, OCH₂-), ^{13}C NMR: 14.28 and 14.45 (2CH₃), 38.42 (d, $^1J_{\text{PC}} = 127.8$ Hz, P=C), 52.40 (OCH₃), 57.91 and 60.12 (2OCH₂-), 61.96 (d, $^2J_{\text{PC}} = 16.6$ Hz, P=C–CH), 126.13 (d, $^1J_{\text{PC}} = 91.9$ Hz, C^{ipso}), 128.56 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.97 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.80 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 139.22, and 155.49 (2C), 167.86 (C=O).

*For two rotamers

Dimethyl-2-[(ethoxycarbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4c**

(0.99g, m.p. 150–151°C, yield 86%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1754, 1650, and 1619 (C=O).

Major isomer, **4c**-(E) (53.1%), ^1H NMR: δ 1.1 (CH_3)*, 2.88 and 3.77 (3H, 2s, OCH_3), 3.97 (4H, m, OCH_2 -)*, 4.90 (1H, d, $J_{\text{PH}} = 16.3$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 7.26–7.58 (40H, m, arm)*. ^{13}C NMR: δ 14.53 (CH_3)*, 40.77 (d, $^1J_{\text{PC}} = 135.1$ Hz, $\text{P}=\text{C}$), 49.40 (OCH_2 -), 52.03 (OCH_3), 61.03 (OCH_3), 61.95 (d, $^2J_{\text{PC}} = 16.5$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 125.93, 126.75 and 126.64 (C)*, 126.61 (d, $^1J_{\text{PC}} = 90.6$ Hz, C^{ipso}), 128.64 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.20 (1C)*, 132.03 (d, $^4J_{\text{PC}} = 2.6$ Hz, C^{para}), 133.78 (d, $^2J_{\text{PC}} = 9.9$ Hz, C^{ortho}), 139.25 and 155.36 (2C), 168.56 (C=O), 169.98 (d, $^2J_{\text{PC}} = 18.8$ Hz, C=O)*, 173.54 (d, $^3J_{\text{PC}} = 6.8$ Hz, C=O)*. Minor isomer, **4c**-(Z) (46.9%), ^1H NMR: δ 3.1 and 3.90 (3H, 2s, OCH_3), 3.43 and 4.26 (4H, 2q, OCH_2 -), ^{13}C NMR: 38.61 (d, $^1J_{\text{PC}} = 129.5$ Hz, $\text{P}=\text{C}$), 48.75 (OCH_2 -), 52.32 (OCH_3), 60.94 (OCH_3), 126.26 (d, $^1J_{\text{PC}} = 92.1$ Hz, C^{ipso}), 128.54 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.93 (d, $^4J_{\text{PC}} = 2.5$ Hz, C^{para}), 133.55 (d, $^2J_{\text{PC}} = 9.7$ Hz, C^{ortho}), 139.49 and 154.99 (2C), 168.45 (C=O).

Diethyl-2-[(ethoxycarbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4d**

(0.96 g, m.p. 155–156°C, yield 78%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1754, 1660, and 1625 (C=O).

Major isomer, **4d**-(E) (50.3%), ^1H NMR: δ 0.29, 0.95, and 1.33 (9H, 3t, CH_3), 3.33, 3.92, and 4.20 (6H, 3q, OCH_2 -), 4.91 (1H, d, $J_{\text{PH}} = 16.4$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 7.23–7.61 (40H, m, arm)*. ^{13}C NMR: δ 13.89, 14.30, and 14.57 (3 CH_3), 40.62 (d, $^1J_{\text{PC}} = 134.8$ Hz, $\text{P}=\text{C}$), 57.38, 60.92, and 61.14 (3 OCH_2 -), 61.89 (d, $^2J_{\text{PC}} = 16.6$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 125.96, 126.71, and 127.34 (3C)*, 126.51 (d, $^1J_{\text{PC}} = 91.9$ Hz, C^{ipso}), 128.55 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.43 (1C)*, 131.86 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.67 (d, $^2J_{\text{PC}} = 9.7$ Hz, C^{ortho}), 139.33 and 155.40 (2C), 168.00 (C=O), 169.28 (d, $^2J_{\text{PC}} = 18.8$ Hz, C=O)*, 172.82 (d, $^3J_{\text{PC}} = 4.9$ Hz, C=O)*.

Minor isomer, **4d**-(Z) (49.7%), ^1H NMR: δ 0.97, 0.98, and 1.34 (9H, 3t, CH_3), 3.68, 4.1, and 4.38 (6H, q, OCH_2 -), 3.43 and 4.26 (4H, 2q, OCH_2 -), ^{13}C NMR: 14.28, 14.39 and 14.45 (3 CH_3), 38.61 (d, $^1J_{\text{PC}} = 133.6$ Hz, $\text{P}=\text{C}$), 52.40 (OCH_3), 57.92, 60.87, and 60.97 (3 OCH_2 -), 126.84 (d, $^1J_{\text{PC}} = 91.3$ Hz, C^{ipso}), 128.43 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.94 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.79 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 139.61 and 155.06 (2C), 167.90 (C=O).

*For two rotamers

Dimethyl-2-[(phenoxy carbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4e**

(0.98g, m.p. 140–142°C, yield 77%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1755, 1660, and 1616 (C=O).

Major isomer, **4e**-(E) (53.5%), ^1H NMR: δ 2.91 and 3.8 (6H, 2s, OCH_3), 5.02 (1H, d, $J_{\text{PH}} = 19.9$ Hz, $\text{P}=\text{C}-\text{CH}$), 6.92–7.63 (50H, m, arm)*. ^{13}C NMR: δ 40.66 (d, $^1J_{\text{PC}} = 135.4$ Hz, $\text{P}=\text{C}$)*, 49.46 and 52.20 (2OCH_3), 61.69 (1H, d, $^2J_{\text{PC}} = 17.4$ Hz, $\text{P}=\text{C}-\text{CH}$), 121.42 and 121.46 (2C^{ortho} , OPh)*, 124.72, 127.50, 127.79, and 131.14 (4CH)*, 126.15 (d, $^1J_{\text{PC}} = 92.0$ Hz, C^{ipso}), 127.39 and 127.56 (4CH)* 128.75 (d, $^3J_{\text{PC}} = 12.2$ Hz, C^{meta}), 131.43 (1C)*, 132.18 (d, $^4J_{\text{PC}} = 2.6$ Hz, C^{para}), 133.78 (d, $^2J_{\text{PC}} = 9.9$ Hz, C^{ortho}), 139.51, 151.67, 153.65 and 155.81 (4C), 168.51 (C=O), 169.99 (d, $^2J_{\text{PC}} = 16.7$ Hz, $\text{C}=\text{O}$)*, 173.17 (d, $^3J_{\text{PC}} = 16.4$ Hz, $\text{C}=\text{O}$)*.

Minor isomer, **4e**-(Z) (46.5%), ^1H NMR: δ 3.1 and 3.9 (6H, 2s, OCH_3), 5.04 (1H, d, $J_{\text{PH}} = 17.7$), ^{13}C NMR: 14.28 and 14.45 (2CH_3), 48.84 and 52.50 (2OCH_3), 62.63 (d, $^2J_{\text{PC}} = 16.9$ Hz, $\text{P}=\text{C}-\text{CH}$), 126.47 (d, $^1J_{\text{PC}} = 87.7$ Hz, C^{ipso}), 128.65 (d, $^3J_{\text{PC}} = 11.8$ Hz, C^{meta}), 129.80, 139.15, 151.68, and 153.66 (4C), 132.08 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.58 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 139.22 and 155.49 (2C), 168.62 (C=O).

Diethyl-2-[(phenoxy carbonyl)anilino]-3-(1,1,1-triphenyl- λ 5-phosphonylidene)succinate **4f**

(0.96g, m.p. 160–161°C, yield 72%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1748, 1650, and 1617 (C=O).

Major isomer, **4f**-(E) (69.0%), ^1H NMR: δ 0.32 and 1.36 (6H, 2t, CH_3), 3.34 and 4.10 (4H, 2q, OCH_2 -), 5.02 (1H, d, $J_{\text{PH}} = 12.4$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 6.91–7.65 (50H, m, arm)*. ^{13}C NMR: δ 13.9 and 14.3 (2CH_3), 40.50 (d, $^1J_{\text{PC}} = 135.0$ Hz, $\text{P}=\text{C}$)*, 57.48 and 60.33 (2OCH_2 -), 60.99 (d, $^2J_{\text{PC}} = 16.6$ Hz, $\text{P}=\text{C}-\text{CH}$)*, 121.41 and 121.47 (C^{ortho} , OPh)*, 124.66, 126.44, 127.70, and 131.31 (4CH)*, 126.40 (d, $^1J_{\text{PC}} = 89.8$ Hz, C^{ipso}), 128.65 (d, $^3J_{\text{PC}} = 12.7$ Hz, C^{meta}), 131.43 (4CH)*, 132.00 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.77 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 139.24 and 151.75 (2C), 167.95 (C=O), 168.82 (d, $^2J_{\text{PC}} = 17.4$ Hz, $\text{C}=\text{O}$)*, 172.68 (d, $^3J_{\text{PC}} = 6.6$ Hz, $\text{C}=\text{O}$)*.

Minor isomer, **4f**-(Z) (31.0%), ^1H NMR: δ 1.25 and 2.04 (6H, 2t, CH_3), 3.71 and 4.43 (4H, 2q, OCH_2 -), ^{13}C NMR: 14.28 and 14.45 (2CH_3), 38.42 (d, $^1J_{\text{PC}} = 127.8$ Hz, $\text{P}=\text{C}$), 57.48 and 60.12 (2OCH_2 -), 61.96 (d, $^2J_{\text{PC}} = 16.6$ Hz, $\text{P}=\text{C}-\text{CH}$), 126.78 (d, $^1J_{\text{PC}} = 91.5$ Hz, C^{ipso}), 128.75 (d, $^3J_{\text{PC}} = 10.8$ Hz, C^{meta}), 132.08 (d, $^4J_{\text{PC}} = 2.6$ Hz, C^{para}), 133.79 (d, $^2J_{\text{PC}} = 9.9$, C^{ortho}), 138.95 and 152.62 (2C), 168.05 (C=O).

*For two rotamers

Dimethyl-2-[[(2-methylphenoxy)carbonyl]anilino]-3-(1,1-triphenyl- λ 5-phosphonylidene) Succinate 4g

(0.95g, m.p. 130–132°C, yield 70%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1754, 1662, and 1618 (C=O).

Major isomer, **4g**-(E) (56.7%), ^1H NMR: δ 2.06 (CH_3), 3.13 and 3.79 (6H, 2s, OCH_3), 4.99 (1H, d, $J_{\text{PH}} = 19.4$ Hz, $\text{P}=\text{C}-\text{CH}^*$), 6.91–7.63 (48H, m, arm)*. ^{13}C NMR: δ 16.17 (CH_3), 40.49 (d, $^1J_{\text{PC}} = 135.4$ Hz, $\text{P}=\text{C}^*$), 49.41 and 52.14 (OCH_3), 61.50 (d, $^2J_{\text{PC}} = 16.1$ Hz, $\text{P}=\text{C}-\text{CH}^*$), 121.89, 125.01, 126.37, 127.21, 127.45, 127.74, and 131.30 (9CH)*, 126.18 (d, $^1J_{\text{PC}} = 92.0$ Hz, C^{ipso}), 128.72 (d, $^3J_{\text{PC}} = 12.2$ Hz, C^{meta}), 139.00, and 150.11 (3C)*, 132.07 (d, $^4J_{\text{PC}} = 2.5$ Hz, C^{para}), 133.77 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 68.84 (C=O), 170.02 (d, $^2J_{\text{PC}} = 17.1$ Hz, $\text{C}=\text{O}^*$), 172.97 (d, $^3J_{\text{PC}} = 13.7$ Hz, $\text{C}=\text{O}^*$).

Minor isomer, **4g**-(Z) (43.3%), ^1H NMR: δ 2.03 (CH_3), 2.89 and 3.88 (6H, 2s, OCH_3), ^{13}C NMR: 16.25 (CH_3), 48.79 and 52.40 (OCH_3), 126.66 (d, $^1J_{\text{PC}} = 91.8$ Hz, C^{ipso}), 128.63 (d, $^3J_{\text{PC}} = 11.7$ Hz, C^{meta}), 132.05 (d, $^4J_{\text{PC}} = 2.5$ Hz, C^{para}), 133.59 (d, $^2J_{\text{PC}} = 9.7$ Hz, C^{ortho}), 139.22 and 155.49 (2C), 168.58 (C=O).

Diethyl-2-[[(2-methylphenoxy)carbonyl] anilino]-3-(1,1-triphenyl- λ 5-phosphonylidene)succinate 4h

(0.93g, m.p. 163–165°C, yield 66%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1745, 1653, and 1622 (C=O).

Major isomer, **4h**-(E) (63.7%), ^1H NMR: δ 0.30 and 1.25 (6H, 2t, CH_3), 2.06 (3H, m, CH_3^*), 3.34 and 4.11 (4H, 2q, OCH_2^-), 5.08 (1H, d, $J_{\text{PH}} = 17.8$ Hz, $\text{P}=\text{C}-\text{CH}^*$), 6.90–7.66 (48H, m, arm)*. ^{13}C NMR: δ 14.20, 14.32, and 16.32 (3 CH_3), 40.42 (d, $^1J_{\text{PC}} = 132.7$ Hz, $\text{P}=\text{C}$), 57.45 and 60.35 (2 OCH_2^-), 61.48 (d, $^2J_{\text{PC}} = 17.9$ Hz, $\text{P}=\text{C}-\text{CH}$), 121.89, 126.36, 126.60, 127.20, 129.86, 130.45, 130.57, and 131.51 (9CH)*, 124.96, 139.04 and 152.87 (3C)*, 126.41 (d, $^1J_{\text{PC}} = 92.0$ Hz, C^{ipso}), 128.65 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 131.43 (1C)*, 132.01 (d, $^4J_{\text{PC}} = 2.5$ Hz, C^{para}), 133.77 (d, $^2J_{\text{PC}} = 10.1$ Hz, C^{ortho}), 167.94 (C=O), 169.87 (d, $^2J_{\text{PC}} = 17.7$ Hz, $\text{C}=\text{O}^*$), 172.37 (d, $^3J_{\text{PC}} = 5.2$ Hz, $\text{C}=\text{O}^*$).

Minor isomer, **4h**-(Z) (36.3%), ^1H NMR: δ 1.37 and 1.39 (6H, 2t, CH_3), 3.72 and 4.38 (4H, 2q, OCH_2^-), ^{13}C NMR: 14.40, 14.49, and 16.23 (3 CH_3), 37.42 (d, $^1J_{\text{PC}} = 133.7$ Hz, $\text{P}=\text{C}$), 57.99 and 60.94 (2 OCH_2^-), 62.45 (d, $^2J_{\text{PC}} = 16.7$ Hz, $\text{P}=\text{C}-\text{CH}$), 126.86 (d, $^1J_{\text{PC}} = 91.8$ Hz, C^{ipso}), 128.54 (d, $^3J_{\text{PC}} = 12.3$ Hz, C^{meta}), 132.09 (d, $^4J_{\text{PC}} = 2.3$ Hz, C^{para}), 133.69 (d, $^2J_{\text{PC}} = 10.0$ Hz, C^{ortho}), 139.22 and 155.49 (2C), 168.04 (C=O).

*For two rotamers

Dimethyl-2-[[[3-methylphenoxy)carbonyl]anilin]-3-(1, 1,1-triphenyl- λ 5-phosphonylidene)succinate **4i**

(1.07g, m.p. 125–127°C, yield 79%); IR (KBr) ($\nu_{\max}$, cm^{-1}): 1749, 1651, and 1619 (C=O).

Major isomer, **4i**-(E) (53.7%), ^1H NMR: δ 2.27 (3H, s, CH_3)*, 3.13 and 3.80 (6H, 2s, OCH_3), 5.00 (1H, d, $J_{\text{PH}} = 19.9$ Hz, $\text{P}=\text{C}-\text{CH}$), 6.73–7.62 (48H, m, arm)*. ^{13}C NMR: δ 21.21 (CH_3), 40.67 (d, $^1J_{\text{PC}} = 133.2$ Hz, $\text{P}=\text{C}$), 48.83 and 52.19 (2OCH_3), 61.59 (d, $^2J_{\text{PC}} = 17.1$ Hz, $\text{P}=\text{C}-\text{CH}$), 118.36, 125.52, 126.41, 127.11, 127.47, 127.78, 131.13 and 151.33 (9CH)*, 122.07, 138.91 and 153.75 (3C)*, 126.41 (d, $^1J_{\text{PC}} = 92.0$ Hz, C^{ipso}), 128.73 (d, $^3J_{\text{PC}} = 12.2$ Hz, C^{meta}), 131.43 (1C)*, 132.16 (d, $^4J_{\text{PC}} = 2.5$ Hz, C^{para}), 133.78 (d, $^2J_{\text{PC}} = 9.8$ Hz, C^{ortho}), 168.50 (C=O), 170.00 (d, $^2J_{\text{PC}} = 19.2$ Hz, C=O)*, 173.14 (d, $^3J_{\text{PC}} = 4.2$ Hz, C=O)*.

Minor isomer, **4i**-(Z) (46.3%), ^1H NMR: δ 2.90 and 3.90 (6H, 2s, OCH_3), 5.03 (1H, d, $J_{\text{PH}} = 18.1$ Hz, $\text{P}=\text{C}-\text{CH}$), ^{13}C NMR: 37.39 (d, $^1J_{\text{PC}} = 136.0$ Hz, $\text{P}=\text{C}$), 49.47 and 52.50 (2OCH_3), 62.56 (d, $^2J_{\text{PC}} = 16.5$ Hz, $\text{P}=\text{C}-\text{CH}$), 126.57 (d, $^1J_{\text{PC}} = 91.5$ Hz, C^{ipso}), 128.64 (d, $^3J_{\text{PC}} = 11.9$ Hz, C^{meta}), 132.05 (d, $^4J_{\text{PC}} = 2.7$ Hz, C^{para}), 133.58 (d, $^2J_{\text{PC}} = 9.6$ Hz, C^{ortho}), 139.22 and 155.49 (2C), 168.60 (C=O).

REFERENCES

- [1] H. Yeganeh, S. Jamshidi, and P. Hayati, *Eur. Polym. J.*, **42**, 17439 (2002).
- [2] S. Dutta and N. Karak, *Prog. Org. Coat.*, **53**, 147 (2005).
- [3] L. Wu, B. You, and D. Li, *J. Appl. Polym. Sci.*, **84**, 1620 (2002).
- [4] J. Y. Kim, D. H. Shim, J. H. Ryu, G. H. Choi, and K. D. Suh, *J. Appl. Polym. Sci.*, **91**, 3549 (2004).
- [5] E. Tavakolinejad, K. Saidi, and M. R. Islami, *Phosphorus. Sulfur. Silicon. Relat. Elem.*, **180**, 1879 (2005).
- [6] I. Yavari and M. R. Islami, *J. Chem. Res(s)*, **3**, 166 (1998).
- [7] I. Yavari, M. R. Islami, and H. R. Bijanzadeh, *Tetrahedron*, **55**, 5547 (1999).
- [8] M. R. Islami, I. Yavari, A. M. Tikdari, L. Ebrahimi, S. Razee, and H. R. Bijanzadeh, *Russ. Chem. Bull.*, **51**, 121, 2244 (2002).
- [9] B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, **89**, 863 (1989).
- [10] I. Yavari and R. Baharifar, *Tetrahedron Lett.*, **39**, 1051 (1998).
- [11] M. R. Islami, Z. Hassani, H. Sheibani, B. Abdollahzadeh, and N. Etminan, *Tetrahedron*, **59**, 4993 (2003).
- [12] M. Maghsoodlou, N. Hazeri, S. M. H. Khorassani, L. Saghatforoush, M. K. Rofouei, and M. Rezaie, *Arkivoc.*, **117** (xiii) (2006).
- [13] M. R. Islami, J. Torghabeh, S. J. Fatemi, Z. Hassani, and A. Amiry, *Syn. Lett.*, **10**, 1707 (2004).
- [14] M. R. Islami, Z. Hassani, and K. Saidi, *Synth. Commun.*, **33**, 65 (2003).

*For two rotamers