

This article was downloaded by:

On: 27 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### (*N*-Isocyanimino)triphenylphosphorane-Mediated, One-Pot, Efficient Synthesis of Sterically Congested 1,1,1-Trifluoro-2-(5-aryl-1,3,4-oxadiazol-2-yl)-2-propanol Derivatives via Intramolecular Aza-Wittig Reaction

Ali Ramazani<sup>a</sup>; Nahid Shajari<sup>a</sup>; Amir Tofangchi Mahyari<sup>a</sup>; Mehdi Khoobi<sup>a</sup>; Yavar Ahmadi<sup>b</sup>; Ali Souldozi<sup>c</sup>

<sup>a</sup> Chemistry Department, Zanjan University, Zanjan, Iran <sup>b</sup> Member of Young Researchers Club, Islamic Azad University, Zanjan, Iran <sup>c</sup> Chemistry Department, Islamic Azad University-Urmia Branch, Urmia, Iran

Online publication date: 19 November 2010

**To cite this Article** Ramazani, Ali , Shajari, Nahid , Mahyari, Amir Tofangchi , Khoobi, Mehdi , Ahmadi, Yavar and Souldozi, Ali(2010) '(*N*-Isocyanimino)triphenylphosphorane-Mediated, One-Pot, Efficient Synthesis of Sterically Congested 1,1,1-Trifluoro-2-(5-aryl-1,3,4-oxadiazol-2-yl)-2-propanol Derivatives via Intramolecular Aza-Wittig Reaction', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185: 12, 2496 – 2502

**To link to this Article:** DOI: 10.1080/10426501003713098

**URL:** <http://dx.doi.org/10.1080/10426501003713098>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## (*N*-ISOCYANIMINO)TRIPHENYLPHOSPHORANE-MEDIATED, ONE-POT, EFFICIENT SYNTHESIS OF STERICALLY CONGESTED 1,1,1-TRIFLUORO-2-(5-ARYL-1,3,4-OXADIAZOL-2-YL)-2-PROPANOL DERIVATIVES VIA INTRAMOLECULAR AZA-WITTIG REACTION

Ali Ramazani,<sup>1</sup> Nahid Shajari,<sup>1</sup> Amir Tofangchi Mahyari,<sup>1</sup> Mehdi Khoobi,<sup>1</sup> Yavar Ahmadi,<sup>2</sup> and Ali Souldozi<sup>3</sup>

<sup>1</sup>Chemistry Department, Zanjan University, Zanjan, Iran

<sup>2</sup>Member of Young Researchers Club, Islamic Azad University, Zanjan Branch, Zanjan, Iran

<sup>3</sup>Chemistry Department, Islamic Azad University–Urmia Branch, Urmia, Iran

*Reactions of (N-isocyanimino)triphenylphosphorane with 1,1,1-trifluoroacetone in the presence of aromatic (or heteroaromatic) carboxylic acids (3-methylbenzoic acid, 1-naphthalenecarboxylic acid, 2-naphthalenecarboxylic acid, 2-furancarboxylic acid, and 2-thiophenecarboxylic acid) proceed smoothly at room temperature and in neutral conditions to afford sterically congested 1,1,1-trifluoro-2-(5-aryl-1,3,4-oxadiazol-2-yl)-2-propanol derivatives in high yields. The reaction proceeds smoothly and cleanly under mild conditions, and no side reactions were observed.*

*Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.*

**Keywords** Aromatic carboxylic acid; aza-Wittig reaction; heteroaromatic carboxylic acid; (*N*-isocyanimino)triphenylphosphorane; 1,3,4-oxadiazole; 1,1,1-trifluoroacetone

## INTRODUCTION


Organophosphorus compounds<sup>1–5</sup> have been extensively employed in organic synthesis as useful reagents, as well as ligands, in a number of transition metal catalysts.<sup>3</sup> Iminophosphoranes are a class of special type of zwitterions that bear a strongly nucleophilic, electron-rich nitrogen. The electron distribution around the P<sup>+</sup>–N<sup>–</sup> bond and its consequent chemical implications have been probed and assessed through theoretical, spectroscopic, and crystallographic investigations.<sup>6</sup> The proton affinity of these iminophosphoranes can be used as a molecular guide to assess their utility as synthetic reagents and their function as ligands in coordination and organometallic chemistry.<sup>6–14</sup>

Received 15 January 2010; accepted 18 February 2010.

This research was supported by the Zanjan University and the Iran National Science Foundation (INSF).

Address correspondence to Ali Ramazani, Chemistry Department, Zanjan University, PO Box 45195-313, Zanjan, Iran. E-mail: aliramazani@gmail.com

The intramolecular version of the *aza*-Wittig-type reaction has attracted considerable attention recently because of its high potential for the synthesis of a wide variety of nitrogen heterocycles, which can be attributed, in good measure, to the rapid progress in the preparation of functionalized iminophosphoranes. Several interesting heterocyclization reactions involving iminophosphoranes have been reviewed.<sup>6–12</sup> These compounds can easily be converted through the *aza*-Wittig reaction with isocyanates, carbon dioxide, or carbon disulfide into functionalized heterocumulenes, which exhibit a rich chemistry of unusual synthetic promise.<sup>6–12</sup> The nucleophilicity at the nitrogen is a factor of essential mechanistic importance in the use of these iminophosphoranes as *aza*-Wittig reagents. Iminophosphoranes are important reagents in synthetic organic chemistry, especially in the synthesis of naturally occurring products and compounds with biological and pharmacological activity.<sup>6–12</sup> However, the organic chemistry of (*N*-isocyanimino)triphenylphosphorane **3** remains almost unexplored.<sup>13,14</sup> (*N*-Isocyanimino)triphenylphosphorane **3** is expected to have synthetic potential because it provides a reaction system in which the iminophosphorane group can react with a reagent having a carbonyl functionality.<sup>13,14</sup> In recent years, we have established a one-pot method for the synthesis of organophosphorus compounds.<sup>15–24</sup> As part of our ongoing program to develop efficient and robust methods for the preparation of heterocyclic compounds,<sup>25–32</sup> we sought to develop a convenient preparation of 1,1,1-trifluoro-2-(5-aryl-1,3,4-oxadiazol-2-yl)-2-propanols **4a–f** from (*N*-isocyanimino)triphenylphosphorane **3**, 1,1,1-trifluoroacetone **2**, and aromatic (or heteroaromatic) carboxylic acids **1** (3-methylbenzoic acid, 1-naphthalenecarboxylic acid, 2-naphthalenecarboxylic acid, 2-furancarboxylic acid, and 2-thiophenecarboxylic acid) in excellent yields under neutral conditions (Scheme 1).



**Scheme 1** Three-component synthesis of sterically congested 2,5-disubstituted 1,3,4-oxadiazoles **4** (see Table I).

1,3,4-Oxadiazoles have attracted interest in medicinal chemistry as surrogates of carboxylic acids, esters, and carboxamides. They are an important class of heterocyclic compounds that have a wide range of pharmaceutical and biological activities including antimicrobial, antifungal, anti-inflammatory, and antihypertensive.<sup>33–39</sup> Several methods have been reported in the literature for the synthesis of 1,3,4-oxadiazoles. These protocols are multistep in nature.<sup>40–45</sup> The most general method involves the cyclization of diacylhydrazides with a variety of reagents, such as thionyl chloride, phosphorous oxychloride, or sulfuric acid, usually under harsh reaction conditions. Few reliable and operationally simple examples have been reported for the one-step synthesis of 1,3,4-oxadiazoles, especially from readily available carboxylic acids and acid hydrazides.<sup>46–50</sup>

## RESULTS AND DISCUSSION

In the last years, several synthetic methods have been reported for the preparation of (*N*-isocyanimino)triphenylphosphorane ( $\text{CNNPPh}_3$ ) **3** (Scheme 1).<sup>13,14</sup> There are

**Table I** Synthesis of sterically congested 2,5-disubstituted 1,3,4-oxadiazoles derivatives **4** (see Scheme 1)

4	Ar	Yield(%) <sup>a,b</sup>
<b>a</b>	3-MeC <sub>6</sub> H <sub>4</sub>	88
<b>b</b>	1-Naphthyl	86
<b>c</b>	2-Naphthyl	83
<b>d</b>	2-Furyl	87
<b>e</b>	2-Thienyl	80
<b>f</b>	2-BrC <sub>6</sub> H <sub>4</sub>	82


<sup>a</sup>We also used acetone and acetophenone instead of 1,1,1-trifluoroacetone **2** in this reaction, but no corresponding products **4** were observed.

<sup>b</sup>Isolated yields.

several reports for the use of (*N*-isocyanimino)triphenylphosphorane **3** in the synthesis of metal complexes.<sup>13,14</sup> However, the application of **3** in the synthesis of organic compounds is rare.<sup>25–31</sup> As part of our ongoing program to develop efficient and robust methods for the preparation of heterocyclic compounds,<sup>25–32</sup> we sought to develop a convenient preparation of sterically congested 2,5-disubstituted 1,3,4-oxadiazoles **4** from (*N*-isocyanimino)triphenylphosphorane **3**, 1,1,1-trifluoroacetone **2**, and aromatic (or heteroaromatic) carboxylic acids **1** (3-methylbenzoic acid, 1-naphthalenecarboxylic acid, 2-naphthalenecarboxylic acid, 2-furancarboxylic acid, and 2-thiophenecarboxylic acid) in excellent yields under neutral conditions (Scheme 1). The carboxylic acid derivative **1** with 1,1,1-trifluoroacetone **2** and (*N*-isocyanimino)triphenylphosphorane **3** in CH<sub>2</sub>Cl<sub>2</sub> react together in a 1:1:1 ratio at room temperature to produce sterically congested 2,5-disubstituted 1,3,4-oxadiazoles **4** and triphenylphosphine oxide **5** (Scheme 1 and Table I). The reaction proceeds smoothly and cleanly under mild conditions, and no side reactions were observed. We also used acetone and acetophenone instead of 1,1,1-trifluoroacetone **2** in this reaction, but no corresponding products **4** were observed. In the both cases, 2-aryl-1,3,4-oxadiazoles and triphenylphosphine oxide **5** were observed, as has been previously reported,<sup>26</sup> and the acetone and acetophenone were recovered unreacted at the end of reaction. As indicated in Table I, the reactions proceeded efficiently with an electron-poor ketone (1,1,1-trifluoroacetone) **2**, and electron-rich ketones such as acetone and acetophenone are not suitable starting materials in these reactions.

The structures of the products were deduced from their IR, mass, H NMR, and <sup>13</sup>C NMR spectra. The mass spectra of these compounds displayed molecular ion peaks at the appropriate *m/z* values. The H NMR spectrum of **4a** consisted of two singlets for the 2 CH<sub>3</sub> ( $\delta$  = 1.93 and 2.43), a broad singlet for a OH ( $\delta$  = 4.95–5.50, exchangeable by D<sub>2</sub>O), and two multiplets for the aromatic protons [ $\delta$  = 7.18–7.61 (m, 2H, arom) and  $\delta$  = 7.63–8.02 (m, 2H, arom)]. The <sup>1</sup>H decoupled <sup>13</sup>C NMR spectrum of **4a** showed 12 distinct resonances, partial assignment of these resonances is given in the Experimental section. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of compounds **4b–f** were similar to those of **4a**, except for the aromatic or heteroaromatic moieties, which exhibited characteristic signals with appropriate chemical shifts.

A mechanistic rationalization for this reaction is provided in Scheme 2. On the basis of the chemistry of isocyanides,<sup>19–25</sup> it is reasonable to assume that the first step may involve nucleophilic addition of the (*N*-isocyanimino)triphenylphosphorane **3** to the 1,1,1-trifluoroacetone **2**, which facilitates by its protonation with the acid **1**, leading to nitrilium intermediate **6**. This intermediate may be attacked by conjugate base of the acid **7** to form 1:1:1 adduct **8**. This adduct may undergo intramolecular aza-Wittig<sup>25–29</sup> reaction



**Scheme 2** The proposed mechanism for the formation of sterically congested 2,5-disubstituted 1,3,4-oxadiazole derivatives **4**.

of the iminophosphorane moiety with the ester carbonyl to afford the isolated sterically congested 2,5-disubstituted 1,3,4-oxadiazoles **4** by removal of triphenylphosphine oxide **5** from intermediate **9**.

We believe that the reported method offers a mild, simple, and efficient route for the preparation of sterically congested 2,5-disubstituted 1,3,4-oxadiazole derivatives. Its ease of workup, high yields, and fairly mild reaction conditions make it a useful addition to modern synthetic methodologies. Other aspects of this process are under investigation.

## EXPERIMENTAL

Starting materials and solvents were obtained from Merck (Germany) and Fluka (Switzerland) and were used without further purification. The methods used to follow the reactions were TLC and NMR. TLC and NMR indicated that there is no side product. Melting points were measured on an Electrothermal 9100 apparatus and are uncorrected. IR spectra were measured on a Jasco 6300 FTIR spectrometer.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were measured ( $\text{CDCl}_3$  solution) with a Bruker DRX-250 Avance spectrometer at 250.0 and 62.5 MHz, respectively. Elemental analyses were performed using a Heraeus CHN-O-Rapid analyzer. Mass spectra were recorded on a Finnigan-MATT 8430 mass spectrometer operating at an ionization potential of 20 eV. Flash chromatography columns were prepared from Merck silica gel powder.

### General Procedure for the Preparation of Compounds 4a–f (Exemplified for 4a)

To a magnetically stirred solution of (N-isocyanimino)triphenylphosphorane **3** (1 mmol) and 1,1,1-trifluoroacetone **2** (1 mmol) in  $\text{CH}_2\text{Cl}_2$  (7 mL), a solution of 3-methylbenzoic acid **1a** (1 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added dropwise at room temperature over 15 min. The mixture was stirred for 4 h. The solvent was removed under reduced pressure, and the viscous residue was purified by flash column chromatography [silica gel powder: petroleum ether–ethyl acetate (2:1)]. The solvent was removed under reduced pressure, and the products were obtained. The characterization data of the compounds are given below:

**1,1,1-Trifluoro-2-[5-(3-methylphenyl)-1,3,4-oxadiazol-2-yl]-2-propanol (4a)**

Yellow crystals; mp: 60.1–61.6°C; Yield: 88%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3292 (br), 1692, 1560, 1162, 1088, 724.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.93 and 2.42 (2 s, 6 H, 2  $\text{CH}_3$ ), 4.95–5.50 (br. s, 1 H, OH; exchanged by  $\text{D}_2\text{O}$  addition), 7.18–7.61 (m, 2 H, arom), 7.63–8.02 (m, 2 H, arom).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  21.19 and 22.27 (2  $\text{CH}_3$ ), 72.93 (q,  $^2J_{\text{CF}} = 32.1$  Hz, C- $\text{CF}_3$ ), 123.65 and 139.23 (2 C, arom), 124.64 (q,  $^1J_{\text{CF}} = 286.2$  Hz,  $\text{CF}_3$ ), 125.34, 128.63, 130.07 and 134.32 (4 CH, arom), 164.99 and 167.54 (2 C=N). Anal. Calcd for  $\text{C}_{12}\text{H}_{11}\text{F}_3\text{N}_2\text{O}_2$  (272.2): C, 52.94; H, 4.07; N, 10.29%. Found: C, 52.99; H, 4.04; N, 10.32. MS (EI):  $m/z$  (%) = 272 ( $\text{M}^+$ , 48), 203 (48), 161 (48), 159 (50), 136 (32), 119 (50), 91 (100), 77 (11), 65 (21) and 43 (28).

**1,1,1-Trifluoro-2-[5-(1-naphthyl)-1,3,4-oxadiazol-2-yl]-2-propanol (4b)**

White crystals; mp: 149.2–151.0°C; Yield: 86%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3472 (br), 1540, 1165, 1103, 772.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.99 (s, 3 H,  $\text{CH}_3$ ), 4.17 (s, 1 H, OH), 7.52–7.78 (m, 3 H, arom), 7.95 (d,  $^3J_{\text{HH}} = 8.0$  Hz, 1 H, arom), 8.08 (d,  $^3J_{\text{HH}} = 8.0$  Hz, 1 H, arom), 8.18 (d,  $^3J_{\text{HH}} = 7.2$  Hz, 1 H, arom), 9.13 (d,  $^3J_{\text{HH}} = 8.5$  Hz, 1 H, arom).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  20.46 ( $\text{CH}_3$ ), 72.38 (q,  $^2J_{\text{CF}} = 32.1$  Hz, C- $\text{CF}_3$ ), 119.89, 130.32 and 134.16 (3 C, arom), 124.01 (q,  $^1J_{\text{CF}} = 286.2$  Hz,  $\text{CF}_3$ ), 125.16, 126.15, 127.26, 128.83, 129.14, 129.27 and 133.69 (7 CH, arom), 163.63 and 166.87 (2 C=N). Anal. Calcd for  $\text{C}_{15}\text{H}_{11}\text{F}_3\text{N}_2\text{O}_2$  (308.2): C, 58.45; H, 3.60; N, 9.09%. Found: C, 58.41; H, 3.62; N, 9.03. MS (EI):  $m/z$  (%) = 308 ( $\text{M}^+$ , 22), 239 (4), 197 (14), 155 (50), 153 (100), 126 (75), 76 (16), 63 (18) and 43 (62).

**1,1,1-Trifluoro-2-[5-(2-naphthyl)-1,3,4-oxadiazol-2-yl]-2-propanol (4c)**

Yellow crystals; mp: 138.5–140.2°C; Yield: 83%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3274 (br), 1556, 1155, 1096, 763.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.99 (s, 3 H,  $\text{CH}_3$ ), 4.94–5.18 (br. s, 1 H, OH; exchanged by  $\text{D}_2\text{O}$  addition), 7.47–7.67 (m, 2 H, arom), 7.74–7.97 (m, 3 H, arom), 8.07 (d,  $^3J_{\text{HH}} = 8.7$  Hz, 1 H, arom), 8.52 (s, 1 H, arom).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  21.27 ( $\text{CH}_3$ ), 72.99 (q,  $^2J_{\text{CF}} = 32.1$  Hz, C- $\text{CF}_3$ ), 120.99, 133.63 and 135.91 (3 C, arom), 123.96, 128.26, 128.96, 129.37, 129.90 and 130.19 (7 CH, arom), 124.69 (q,  $^1J_{\text{CF}} = 286.2$  Hz,  $\text{CF}_3$ ), 165.04 and 167.61 (2 C=N). Anal. Calcd for  $\text{C}_{15}\text{H}_{11}\text{F}_3\text{N}_2\text{O}_2$  (308.2): C, 58.45; H, 3.60; N, 9.09%. Found: C, 58.51; H, 3.62; N, 9.05. MS (EI):  $m/z$  (%) = 308 ( $\text{M}^+$ , 55), 239 (12), 197 (29), 155 (100), 153 (84), 127 (82), 100 (11), 77 (11), 63 (10) and 43 (39).

**1,1,1-Trifluoro-2-[5-(2-furyl)-1,3,4-oxadiazol-2-yl]-2-propanol (4d)**

Yellow crystals; mp: 120.7 (dec) °C; Yield: 87%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3245 (br), 1642, 1526, 1155, 1096, 751.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.92 (s, 3 H,  $\text{CH}_3$ ), 4.51 (s, 1 H, OH; exchanged by  $\text{D}_2\text{O}$  addition), 6.62 (s, 1 H, furan), 7.25 (s, 1 H, furan), 7.66 (s, 1 H, furan).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  20.39 ( $\text{CH}_3$ ), 72.27 (q,  $^2J_{\text{CF}} = 32.1$  Hz, C- $\text{CF}_3$ ), 112.71, 115.76 and 146.73 (3 CH, furan), 123.86 (q,  $^1J_{\text{CF}} = 286.2$  Hz,  $\text{CF}_3$ ), 138.78 (C, furan), 159.42 and 163.38 (2 C=N). Anal. Calcd for  $\text{C}_9\text{H}_7\text{F}_3\text{N}_2\text{O}_3$  (248.2): C, 43.56; H, 2.84; N, 11.29%. Found: C, 43.62; H, 2.86; N, 11.24. MS (EI):  $m/z$  (%) = 248 ( $\text{M}^+$ , 69),

194 (32), 179 (91), 137 (100), 108 (39), 95 (99), 52 (31) and 43 (69). Sample NMR, IR, and mass spectra for **4d** are shown in Figures S1–S6 (Supplemental Materials, available online).

### 1,1,1-Trifluoro-2-[5-(2-thienyl)-1,3,4-oxadiazol-2-yl]-2-propanol (**4e**)

Yellow crystals; mp: 94.2–96.1 °C; Yield: 80%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3237 (br), 1603, 1163, 1108, 727.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.92 (s, 3 H,  $\text{CH}_3$ ), 4.41–4.61 (br. s, 1 H, OH; exchanged by  $\text{D}_2\text{O}$  addition), 7.14–7.24 (m, 1 H, thiophene), 7.60 (d,  $^3J_{\text{hh}} = 4.2$  Hz, 1 H, thiophene), 7.81 (s, 1 H, thiophene).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  21.11 ( $\text{CH}_3$ ), 72.89 (q,  $^2J_{\text{cf}} = 32.1$  Hz, C- $\text{CF}_3$ ), 124.54 (q,  $^1J_{\text{cf}} = 286.2$  Hz,  $\text{CF}_3$ ), 124.92 (C, thiophene), 129.30, 131.88 and 132.20 (3 CH, thiophene), 163.61 and 164.17 (2 C=N). Anal. Calcd for  $\text{C}_9\text{H}_7\text{F}_3\text{N}_2\text{O}_2\text{S}$  (264.2): C, 40.91; H, 2.67; N, 10.60%. Found: C, 40.99; H, 2.72; N, 10.56. MS (EI):  $m/z$  (%) = 264 ( $\text{M}^+$ , 4), 195 (4), 153 (7), 124 (8), 111 (100), 96 (7), 82 (19), 69 (24) and 42 (32).

### 2-[5-(2-Bromophenyl)-1,3,4-oxadiazol-2-yl]-1,1,1-trifluoro-2-propanol (**4f**)

Yellow crystals; mp: 83.5–85.1 °C; Yield: 82%. IR (KBr) ( $\nu_{\max}$ ,  $\text{cm}^{-1}$ ): 3431 (br), 1456, 1167, 1089, 728.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 250 MHz):  $\delta_{\text{H}}$  1.93 (s, 3 H,  $\text{CH}_3$ ), 4.08–4.32 (br. s, 1 H, OH; exchanged by  $\text{D}_2\text{O}$  addition), 7.35–8.02 (m, 4 H, arom).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 62.5 MHz):  $\delta_{\text{C}}$  20.36 ( $\text{CH}_3$ ), 72.36 (q,  $^2J_{\text{cf}} = 32.1$  Hz, C- $\text{CF}_3$ ), 122.15 and 124.76 (2 C, arom), 123.90 (q,  $^1J_{\text{cf}} = 286.2$  Hz,  $\text{CF}_3$ ), 128.06, 132.23, 133.48 and 134.97 (4 CH, arom), 164.62 and 165.78 (2 C=N). Anal. Calcd for  $\text{C}_{11}\text{H}_8\text{BrF}_3\text{N}_2\text{O}_2$  (337.1): C, 39.19; H, 2.39; N, 8.31%. Found: C, 39.22; H, 2.44; N, 8.22. MS (EI):  $m/z$  (%) = 338 ( $[\text{M}^+ + 2]$ , 8), 336 ( $\text{M}^+$ , 7), 304 (6), 269 (10), 225 (31), 183 (58), 155 (29), 102 (22), 89 (36), 75 (100), 62 (35) and 43 (100).

## REFERENCES

1. O. I. Kolodiazny, *Phosphorus Ylides: Chemistry and Applications in Organic Chemistry* (Wiley, New York, 1999).
2. A. Ramazani, A. R. Kazemizadeh, E. Ahmadi, N. Noshiranzadeh, and A. Souldozi, *Curr. Org. Chem.*, **12**, 59 (2008).
3. W. C. Kaska, *Coord. Chem. Rev.* **48**, 1 (1983).
4. A. Ramazani, A. R. Kazemizadeh, E. Ahmadi, K. Slepokura, and T. Lis, *Z. Naturforsch.*, **61b**, 1128 (2006).
5. D. E. C. Cobridge, *Phosphorus: An Outline of Chemistry, Biochemistry and Uses*, 5th ed. (Elsevier, Amsterdam, 1995).
6. P. Molina and M. J. Vilaplana, *Synthesis*, 1197 (1994).
7. F. Palacios, D. Aparicio, G. Rubiales, C. Alonso, and J. M. de los Santos, *Curr. Org. Chem.*, **13**, 808 (2009).
8. F. Palacios, D. Aparicio, G. Rubiales, C. Alonso, and J. M. de los Santos, *Curr. Org. Chem.*, **10**, 2371 (2006).
9. G. Hajós and I. Nagy, *Curr. Org. Chem.*, **12**, 39 (2008).
10. F. Palacios, C. Alonso, D. Aparicio, G. Rubiales, and J. M. de los Santos, *Tetrahedron*, **63**, 523 (2007).

11. F. P. Cossio, C. Alonso, B. Lecea, M. Ayerbe, G. Rubiales, and F. Palacios, *J. Org. Chem.*, **71**, 2839 (2006).
12. F. Palacios, E. Herrán, C. Alonso, G. Rubiales, B. Lecea, M. Ayerbe, and F. P. Cossío, *J. Org. Chem.*, **71**, 6020 (2006).
13. H. Stolzenberg, B. Weinberger, W. P. Fehlhammer, F. G. Pühlhofer, and R. Weiss, *Eur. J. Inorg. Chem.*, **21**, 4263 (2005).
14. T. W. Chiu, Y. H. Liu, K. M. Chi, Y. S. Wen, and K. L. Lu, *Inorg Chem.*, **44**, 6425 (2005).
15. I. Yavari and A. Ramazani, *Synth. Commun.*, **26**, 4495 (1996).
16. I. Yavari and A. Ramazani, *Phosphorus, Sulfur, and Silicon*, **130**, 73 (1997).
17. A. Ramazani and A. Bodaghi, *Tetrahedron Lett.*, **41**, 567 (2000).
18. P. Pakravan, A. Ramazani, N. Noshiranzadeh, and A. Sedrpoushan, *Phosphorus, Sulfur, and Silicon*, **182**, 545 (2007).
19. A. Ramazani, M. Rahimifard, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **182**, 1 (2007).
20. A. Ramazani, M. Rahimifard, N. Noshiranzadeh, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **182**, 413 (2007).
21. I. Yavari, A. Ramazani, and A. Yahya-Zadeh, *Synth. Commun.*, **26**, 4495 (1996).
22. A. Ramazani, E. Ahmadi, A. R. Kazemizadeh, L. Dolatyari, N. Noshiranzadeh, I. Eskandari, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **180**, 2419 (2005).
23. A. Ramazani and M. Mohammadi-Vala, *Phosphorus, Sulfur, and Silicon*, **176**, 223 (2001).
24. A. Ramazani, I. Amini, and A. Massoudi, *Phosphorus, Sulfur, and Silicon*, **181**, 2225 (2006).
25. A. Souldozi, A. Ramazani, N. Bouslimani, and R. Welter, *Tetrahedron Lett.*, **48**, 2617 (2007).
26. A. Souldozi and A. Ramazani, *Tetrahedron Lett.*, **48**, 1549 (2007).
27. A. Ramazani and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **184**, 3191 (2009).
28. A. Ramazani and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **184**, 2344 (2009).
29. A. Ramazani and A. Souldozi, *Arkivoc*, **xvi**, 235 (2008).
30. A. Ramazani, S. Salmanpour, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **185**, 97 (2010).
31. A. Souldozi, A. Ramazani, K. Ślepokura, and T. Lis, *Z. Naturforsch.*, **62b**, 835 (2007).
32. A. Ramazani, A. Morsali, B. Ganjeie, A. R. Kazemizadeh, E. Ahmadi, R. Kempe, and I. Hertle, *Z. Naturforsch.*, **60b**, 569 (2005).
33. W. R. Tully, C. R. Gardner, R. J. Gillespie, and R. Westwood, *J. Med. Chem.*, **34**, 2060 (1991).
34. C. Chen, C. H. Senanayake, T. J. Bill, R. D. Larsen, T. R. Verhoeven, and P. J. Reider, *J. Org. Chem.*, **59**, 3738 (1994).
35. B. S. Holla, R. Gonsalves, and S. Shenoy, *Eur. J. Med. Chem.*, **35**, 267 (2000).
36. M. J. Crimmin, P. J. Hanlon, N. H. Rogers, and G. Walker, *J. Chem. Soc., Perkin Trans. 1*, 2047 (1989).
37. U. V. Laddi, S. R. Desai, R. S. Bennur, and S. C. Bennur, *Indian J. Heterocycl. Chem.*, **11**, 319 (2002).
38. Q. Chen, X.-H. Zhu, L.-L. Jiang, Z.-M. Liu, and G.-F. Yang, *Eur. J. Med. Chem.*, **47**, 595 (2008).
39. Z.-M. Liu, G.-F. Yang, and X.-H. Qin, *J. Chem. Technol. Biotechnol.*, **76**, 1154 (2001).
40. I. R. Baxendale, S. V. Ley, and M. Martinelli, *Tetrahedron*, **61**, 5323 (2005).
41. S. Liras, M. P. Allen, and B. E. Segelstein, *Synth. Commun.*, **30**, 437 (2000).
42. B. J. Brown, I. R. Clemens, and J. K. Neesom, *Synlett*, 131 (2000).
43. F. T. Coppo, K. A. Evans, T. L. Graybill, and G. Burton, *Tetrahedron Lett.*, **45**, 3257 (2004).
44. C. T. Brain, J. M. Paul, Y. Loong, and P. J. Oakley, *Tetrahedron Lett.*, **40**, 3275 (1999).
45. C. T. Brain and S. A. Brunton, *Synlett*, 382 (2001).
46. V. K. Tandon and R. B. Chhor, *Synth. Commun.*, **31**, 1727 (2001).
47. S. H. Mashraqui, S. G. Ghadigaonkar, and R. S. Kenny, *Synth. Commun.*, **33**, 2541 (2003).
48. F. Bentiss, M. Lagrenee, and D. Barbry, *Synth. Commun.*, **31**, 935 (2001).
49. E. Jedlovská and J. Lesko, *Synth. Commun.*, **24**, 1879 (1994).
50. Y. Wang, D. R. Sauer, and S. W. Djuric, *Tetrahedron Lett.*, **47**, 105 (2006).