

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Regioselective Cycloadditions of Phosphonyl Nitrile Oxides with Vinylphosphonate and Phosphaalkyne

Li-li Shen^{ab}; Yong Ye^{ac}; Yong Luo^c; Lun-zu Liu^c

^a Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou, China ^b Department of Chemistry, Henan Institute of Education, Zhengzhou, China ^c State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin, China

Online publication date: 24 February 2010

To cite this Article Shen, Li-li, Ye, Yong, Luo, Yong and Liu, Lun-zu(2010) 'Regioselective Cycloadditions of Phosphonyl Nitrile Oxides with Vinylphosphonate and Phosphaalkyne', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185: 3, 680 – 687

To link to this Article: DOI: 10.1080/10426500902917651

URL: <http://dx.doi.org/10.1080/10426500902917651>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

REGIOSELECTIVE CYCLOADDITIONS OF PHOSPHONYL NITRILE OXIDES WITH VINYLPHOSPHONATE AND PHOSPHAALKYNE

Li-li Shen,^{1,2} Yong Ye,^{1,3} Yong Luo,³
and Lun-zu Liu³

¹Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou, China

²Department of Chemistry, Henan Institute of Education, Zhengzhou, China

³State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin, China

1,3-Dipolar cycloaddition reactions are important synthetic manipulations allowing the construction of five-membered heterocycles. In this article, we report the cycloaddition of phosphonyl nitrile oxides with vinylphosphonate and phosphalkyne to form the unexpected 2:1 cycloaddition product with excellent levels of regiocontrol product. The structures of title compounds were confirmed by ¹H NMR, ³¹P NMR, MS, and IR. The mechanism of the cycloaddition was explored using the density functional theory (DFT) method.

Keywords Cycloaddition; dihydroisoxazoles; nitrile oxides

INTRODUCTION

Cycloaddition reactions are important synthetic manipulations for construction of the five-member ring carbocycles and heterocycles.^{1–2} The reaction between nitrile oxides and alkenes is of considerable interest in organic synthesis, as the resulting heterocycles are versatile intermediates for the syntheses of natural products and biologically active compounds.³ Intensive investigations have been undertaken on 1,3-dipolar cycloaddition reactions between nitrile oxide and electron-deficient alkenes, and in most cases 4,5-dihydroisoxazoles were obtained. Although there are several examples of the 1,3-dipolar cycloaddition of electron-deficient nitrile oxide with unsaturated alkenes, the scope and generality of the nitrile oxides remain insufficient. In this article, we report the cycloaddition of phosphonyl nitrile oxides and the formation of an unexpected 2:1 cycloaddition product. It provides a direct route to obtain triphosphonyl-substituted dihydroisoxazolyl dihydroisoxazoles and diphosphonyl-substituted dihydroisoxazolyl dihydroisoxazoles with

Received 16 January 2009; accepted 23 March 2009.

The authors would like to thank financial supports from NNSFC (Nos. 20602032, 20972143).


Address correspondence to Yong Ye, Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou 450052, China. E-mail: yeyong03@mail.nankai.edu.cn

excellent regiocontrol product. We believe that this strategy provides a potentially facile route to a wide range of dihydroisoxazolyl dihydroisoxazoles-based targets.

RESULTS AND DISCUSSION

Cycloaddition of Phosphonyl Nitrile Oxides with Phosphaalkyne

The requisite nitrile oxides **1** were prepared according to the literature from the corresponding hydroximoyl bromides via base-induced dehydrobromination.⁴⁻⁶ We first investigated the 1,3-dipolar cycloaddition between nitrile oxide and phosphaalkyne **2**, as shown in Scheme 1. In our initial attempts, we failed to isolate nitrile oxide **1**. Therefore, without purification, nitrile oxide was directly used for the cycloaddition reaction. Tert-butyl phosphaalkyne **2** can react with nitrile oxides **1** to give heterocyclic compounds. However, we found that an unexpected 2:1 cycloaddition product **4** is achieved when threefold excess of tert-butyl phosphaalkyne reacts with phosphonyl nitrile oxides. Its regio isomer **4'** is not obtained. This is because of the steric bulk of the tert-butyl on phosphaalkyne **2** and the highly reactive dipole **1**, so only one regio isomer is obtained.



Scheme 1 Cycloaddition of phosphonyl nitrile oxides with phosphaalkyne.

When onefold excess of dibutoxyphosphonyl nitrile oxides **1b** was added, a 1:1 cycloaddition product **3b** was observed from ¹H NMR and ³¹P NMR spectra. From ³¹P NMR (Figure 1), a doublet at 86.3 ppm is observed, which indicates a typical bivalent phosphorus chemical displacement.⁷ It is a chemical displacement of phosphorus atom in isoxazole. Another doublet at 7.5 ppm is the chemical displacement of phosphorus atom in phosphonate. From ³¹P NMR, a coupling constant of 67.7 Hz can also be observed. In the case of a reverse dipole orientation as represented by the structure **3'**, the ^αP resonances would be expected at a considerably lower field ($\delta = \text{ca. } 160\text{--}180$).⁸

In addition, from the integration of the hydrogen in ¹H NMR, a 1:1 cycloaddition product is also observed. The 1:1 cycloaddition product **3** could not be isolated. After




Figure 1 ^{31}P NMR of compound **3**.

another one equivalent of phosphonyl nitrile oxide was added to the reaction, the expected product **4** is formed. We can draw the conclusion that the cycloaddition reaction is completed in two steps. First, a 1:1 cycloaddition product **3** is formed. Second, another phosphonyl nitrile oxide **1** adds to the $\text{P}=\text{C}$ double bond of **3**, so forming a 2:1 product **4**.

The structure of the 2:1 cycloaddition product **4a** is also confirmed by ^1H and ^{31}P NMR. Figure 2 shows a ^{31}P spectrum of compound **4a**. Since the middle-group phosphorus (βP) is attached to two chemically different phosphorus atoms (αP), spin-spin splitting into a 1:2:1 triplet is to be expected in δ 16.1–17.6. Coupling constants of 56.1 Hz can be observed. Likewise, the attachment of the two end-group phosphorus atoms to the lone middle-group phosphorus gives spin-spin splitting into 1:1 doublet at δ 1.05–1.75.

Cycloaddition of Phosphonyl Nitrile Oxides with Alkene

We also examined the scope of phosphonyl nitrile oxides with an electron-deficient alkene. With acrylonitrile, diisopropoxy phosphonyl nitrile oxides **1** showed analogous






Figure 2 ^{31}P NMR of compound **4a**.



Scheme 2 Cycloaddition of phosphonyl nitrile oxides with acrylonitrile.

behavior. Nitrile oxide **1** reacted regioselectively with acrylonitrile to give an unexpected 2:1 product **5** (shown in Scheme 2).

When reacted with trans-vinylphosphonate, phosphonyl nitrile oxides **1** also showed analogous behavior (shown in Scheme 3). The vinylphosphonate **6** was prepared according to the procedures we previously described.^{2h} When diisopropoxy phosphonyl nitrile oxides **1** reacted with β -substituted vinylphosphonate **6**, only the regioselective 2:1 addition product **7** was obtained. Meanwhile, when excessive β -substitute vinylphosphonate **6** was added, only the 2:1 cycloadduct was obtained.



A-F: R = C₆H₅, C₆H₁₃, *p*-O₂NC₆H₄, CO₂CH₃, 2,4-Cl₂C₆H₃, *m*-O₂NC₆H₄

Scheme 3 Cycloaddition of phosphonyl nitrile oxides with vinylphosphonate.

The Mechanism of Cycloaddition

The detailed mechanism has been explored using the density functional theory (DFT) method. Each structure was fully optimized with the B3LYP⁹⁻¹³ method using the 6-31G* basis set for C, O, and H atoms (Figure 3). Harmonic vibrational frequencies were calculated for each structure. The bond orders reported were Wiberg bond calculated by means of natural bond orbitals (NBO). The charges reported were the Mulliken atomic charges. All calculations were carried out with the Gaussian 03 program package. All relative energies discussed within this context are free energies at 298 K. In order to make the calculations easier, we used methyl instead of isopropyl.

A plausible reaction pathway has been proposed to account for the high reactivity and selectivity of diisopropoxy phosphonyl nitrile oxide. First, a regiospecific 1:1 product was formed. Second, a highly reactive diisopropoxy phosphonyl nitrile oxide can continue reacting with the C=N double bond and give the corresponding cycloadduct. From Figure 3




Figure 3 Potential energy surface of the reaction with the B3LYP method. All energies are electronic energies without ZPE (zero point energy) corrections with respect to the reactants. ts1 and ts2 stand for the transition state of the first step and second step.

we can see that the free energy barriers for the first and second steps are 11.7 and -14.6 kcal/mol, respectively. The free energy of the second step is quite low. So the cycloaddition stopped in this step.

CONCLUSIONS

In conclusion, the [3+2] cycloaddition reaction of nitrile oxides with alkynes and alkenes provides a direct route to triphosphonyl-substituted dihydroisoxazolyl dihydroisoxazoles and diphosphonyl-substituted dihydroisoxazolyl dihydroisoxazoles with excellent levels of regiocontrol product. Additionally, an unexpected 2:1 cycloaddition takes place in the presence of Et_3N when phosphonyl hydroximoyl bromides are employed to furnish the dihydroisoxazoles. We believe that this strategy provides a potentially facile route to a wide range of dihydroisoxazolyl dihydroisoxazoles-based targets. The DFT studies of the reactions between nitrile oxides and acrylonitrile reveal that the following mechanism is quite possible: It starts as a normal 1,3-dipolar cycloaddition reaction to produce a regioselective 1:1 product, then highly reactive diisopropyl phosphonyl nitrile oxide continue to react with the regioselective 1:1 product and give the corresponding cycloadduct. Further study is underway to expand the scope of this methodology, as well as to ascertain mechanistic details of the cycloaddition process.

EXPERIMENTAL

Elemental analyses were carried on a Yanaco CHN Corder MT-3 apparatus. ^1H , ^{13}C , and ^{31}P NMR spectra were measured by using a Bruker 300 spectrometer with TMS and

85% H_3PO_4 as the internal and external references, respectively, and with CDCl_3 as the solvent. IR spectra were recorded as KBr pellets on a Bruck spectrometer. Mass spectra were acquired in positive ion mode using a Bruker Esquire-LCTM ion trap spectrometer equipped with a gas nebulizer probe, capable of analyzing ions up to m/z 20000. Solvents used were purified and dried by standard procedures.

Procedure for the Synthesis of 4

To a rapidly stirred solution of tert-butyl phosphaacetylene **2** (0.2 mL, 1 M in $\text{Me}_3\text{SiOSiMe}_3$) and Et_3N (0.15 g) in dry ether (10 mL) under N_2 , a solution of compound **1** (0.288 g, 1 mmol) in dry ether (6 mL) was added dropwise at -10°C . The mixture was stirred at room temperature for 24 h. Then, the reaction mixture was filtered to remove triethylamine hydrobromides, and the solvent was evaporated in vacuum. The residue was chromatographed on a silica gel column with petroleum ether:ethyl acetate 1:1 (V:V) to give pure **4a–b** as an oil.

Similar to the synthesis of **4**, compound **3** was obtained as an oil. Yield 43.2%, ^1H NMR: 0.87–0.94 (m, 6H), 1.19–1.44 (m, 13H), 1.67–1.68 (m, 4H), 4.17 (m, 4H); ^{31}P NMR: 86.3 (d, $J = 67.7$ Hz, $^\alpha\text{P}$), 7.5 (d, $J = 67.7$ Hz, $^\beta\text{P}$).

4a: Yield 50.4%, anal. calcd. found (calcd): C 44.27 (44.36), H 7.32 (7.25), N 5.44 (5.44). ^1H NMR: 4.78 (m, 4H, CHMe_2), 1.26–1.33 (t, 24H, CHMe_2), 1.03 (s, 9H, Me); ^{13}C NMR: 20.1 (CH_3), 50.0 (CH), 26.3 (MeC), 36.6 (Me_3C), 130.0 (P–C), 156.0 (C=N); ^{31}P NMR: 1.4 ($^\alpha\text{P}$), 16.8 ($^\beta\text{P}$).

4b: Yield 76.4%, anal. calcd. found (calcd): C 48.46 (48.42), H 7.92 (7.95), N 4.88 (4.91). ^1H NMR: 4.66 (m, 4H, CH), 1.67 (m, 8H, CH_2), 1.28–1.36 (m, 12H, CH_3), 1.08 (s, 9H, CMe_3), 0.88–0.95 (t, 12H, Me), ^{31}P NMR: 1.7 ($^\alpha\text{P}$), 17.6 ($^\beta\text{P}$).

General Procedure for the Synthesis of 5 and 7

To stirred solution of compound **1** (0.432 g, 1.5 mmol) and alkene (15 mmol) in dry ether (10 mL), a solution of Et_3N (0.2 g, 2 mmol) in dry ether (10 mL) was added dropwise at -10°C . After 30 min, the solution was warmed to ambient temperature. The reaction mixture was stirred continuously for 3 days. After it was filtered, the solvent was evaporated in vacuo. The residue was chromatographed on a silica gel column with petroleum ether:ethyl acetate (1:3) to give pure **5** or **7** as an oil.

5: Yield 22%, anal. calcd. found (calcd): C 43.64 (43.69), H 6.63 (6.68), N 8.95 (8.99), ^1H NMR: 5.03–5.12 (dd, $J = 35$ Hz, 1H, CHCN), 4.53–4.61 (m, 4H, OCHMe_2), 3.53–3.59 (t, $J = 7.8$ Hz, 2H, CH_2CHCN), 0.99–1.23 (m, 24H, OCHMe_2); ^{13}C NMR: 22.0 (CH_3CH), 50.0 (CH_3CH), 70.0 (CH_2), 128.0 (CP), 131.0 (CH), 148.0 (PC=N), 151.0 (CN); ^{31}P NMR: 0.6 ($^\alpha\text{P}$), -3.4 ($^\beta\text{P}$).

7a: Yield 19.6%, anal. calcd. found (calcd): C 47.72 (47.71), H 6.79 (6.93), N 4.29 (4.28), ^1H NMR: 7.32 (m, 5H), 5.89–6.03 (dd, $J = 30$ Hz, 1H), 4.69 (m, 4H), 4.12–4.22 (m, 5H), 1.20–1.39 (m, 30H), ^{13}C NMR: 15.0 (CH_2CH_3), 22.3 (CH_3), 22.6 (CH_3), 30.6 (CH), 49.0 (CPh), 50.0 (CH), 56.0 (CH_2CH_3), 82.0 (CHP), 92.0 (PC–N), 160.0 (PC=N), 126.2, 128.5, 129.6, 143.7 (C_{arom}); ^{31}P NMR: 0.9 ($^\alpha\text{P}$), -3.2 ($^\beta\text{P}$), 18.5 ($^\gamma\text{P}$).

7b: Yield 31.4%, anal. calcd. found (calcd): C 46.98 (47.13), H 8.13 (8.06), N 4.28 (4.23). ^1H NMR: 4.65–4.75 (m, 5H), 3.99–4.10 (m, 5H), 1.08–1.41 (m, 40H), 0.73–0.76 (m, 3H), ^{13}C NMR: 14.2 (CH_2CH_3), 20.3 (CH_2), 22.3 (CH_2), 22.6 (CH_3), 27.4 (CH_2),

30.4 (CH₂), 32.4 (CH₂), 62.3 (CH₂CH₃), 80.1 (CHO), 82.5 (PC–O), 92.4 (PCN), 160.0 (PC=N); ³¹P NMR: 1.1 (αP), –3.1 (βP), 19.2 (γP).

7c: Yield 56.1%, anal. calcd. found(calcd): C 44.59 (44.64), H 6.35 (6.34), N 5.98 (6.01), ¹H NMR: 8.19–8.23 (d, J = 15 Hz, 2H), 7.51–7.55 (d, J = 22 Hz, 2H), 6.06 (dd, J = 35 Hz, 1H), 4.83 (m, 4H), 4.21 (m, 5H), 1.23–1.37 (m, 30H), ¹³C NMR: 14.1 (CH₂CH₃), 22.4 (CH₃), 22.5 (CH₃), 30.5 (CH), 62.4 (CH₂CH₃), 71.5 (CH), 83.3 (CHP), 92.0 (PC–O), 123.2, 127.5, 145.7, 154.6 (C_{arom}), 162.0 (PC=N); ³¹P NMR: 0.5 (αP), –3.3 (βP), 17.7 (γP).

7d: Yield 78.8%, anal. calcd. found(calcd): C 41.49 (41.51), H 6.80 (6.81), N 4.38 (4.40), ¹H NMR: 5.31–5.45 (dd, J = 18 Hz, 1H), 4.77 (m, 4H), 4.13–4.20 (m, 5H), 3.75–3.77 (s, 3H), 1.27–1.37 (m, 30H), ¹³C NMR: 14.5 (CH₂CH₃), 22.4 (CH₃), 32.5 (CH), 50.7 (OCH₃), 62.7 (CH₂CH₃), 72.1 (CHO), 79.2 (PCHO), 88.6 (PC–N), 162.5 (PC=N), 175.4 (C=O); ³¹P NMR: 0.03 (αP), –3.3 (βP), 17.7 (γP); IR: 3268 2957 1739 1649 1579 1490 1439 1204 924 769 cm^{–1}; ESI-MS: 637.3 [M+H]⁺.

7e: Yield 56.1%, anal. calcd. found(calcd): C 43.1 (43.16), H 5.96 (5.99), N 3.85 (3.87), ¹H NMR: 7.24–7.37 (m, 3H), 6.19–6.32 (dd, J = 27 Hz, 1H), 4.67–4.87 (m, 4H), 4.15–4.31 (m, 5H), 1.20–1.39 (m, 30H), ¹³C NMR: 14.6 (CH₂CH₃), 22.3 (CH₃), 22.5 (CH₃), 61.7 (CH₂CH₃), 72.2 (CHO), 82.8 (PCHO), 91.6 (PC–N), 162.0 (PC=N), 126.6, 128.8, 131.8, 132.7, 147.4 (C_{arom}); ³¹P NMR: 0.9 (αP), –3.2 (βP), 18.5 (γP).

7f: Yield 35.3%, anal. calcd. found(calcd): C 44.65 (44.64), H 6.37 (6.34), N 6.03 (6.01), ¹H NMR: 8.14–8.18 (d, J = 27 Hz, 2H), 7.54–7.66 (dd, J = 24 Hz, 2H), 5.95–6.09 (dd, J = 22 Hz, 1H), 4.69 (m, 4H), 4.14–4.36 (m, 5H), 1.21–1.35 (m, 30H), ¹³C NMR: 14.7 (CH₂CH₃), 22.3 (CH₃), 22.5 (CH₃), 30.2 (CH), 62.7 (CH₂CH₃), 72.0 (CHP), 85.4 (PCH–O), 90.3 (PCO), 161.0 (PC=N), 120.2, 122.5, 127.6, 129.3, 147.7, 149.2 (C_{arom}); ³¹P NMR: 0.5 (αP), –3.4 (βP), 17.6 (γP); ESI-MS: 700.5 [M+H]⁺.

REFERENCES

- For reviews, see: (a) R. Huisgen, *In 1,3-Dipolar Cycloaddition Chemistry*, A. Padwa, Ed. (Wiley, New York, 1984), p. 1; (b) A. R. S. Ferwanah and A. M. Awadallah, *Molecules*, **10**, 492 (2005); (c) K. Ruck-Braun, T. H. E. Freysoldt, and F. Wierschem, *Chem. Soc. Rev.*, **34**, 507 (2005); (d) R. P. Litvinovskaya and V. A. Khripach, *Uspekhi Khimii*, **70**, 464 (2001); (e) J. K. Gallos and A. E. Koumbis, *Curr. Org. Chem.*, **7**, 39 (2003); (f) I. N. N. Namboothiri and A. Hassner, *Top. Curr. Chem.*, **216**, 1 (2001); (g) R. Huisgen, *Angew. Chem. Int. Ed.*, **2**, 565 (1963); (h) T. M. V. D. P. E. Melo, *Curr. Org. Chem.*, **9**, 925 (2005); (i) R. Huisgen, *Angew. Chem. Int. Ed.*, **2**, 633 (1963); (j) I. N. N. Namboothiri, N. Rastogi, B. Ganguly, S. M. Mobin, and M. Cojocaru, *Tetrahedron*, **60**, 1453 (2004); (k) R. Huisgen, *Chem. Pharm. Bull.*, **48**, 757 (2000).
- For representative examples, see: (a) Z. X. Yu, P. Caramella, and K. N. Houk, *J. Am. Chem. Soc.*, **125**, 15420 (2003); (b) E. Coutouli-Argyropoulou, P. Lianis, M. Mitakou, A. Giannoulis, and J. Nowak, *Tetrahedron*, **62**, 1494 (2006); (c) H. Takikawa, Y. Hachisu, and J. W. Bode, *Angew. Chem. Int. Ed.*, **45**, 3492 (2006); (d) N. M. Fedou, P. J. Parsons, E. M. E. Viseux, and A. J. Whittle, *Org. Lett.*, **7**, 3179 (2005); (e) T. V. Hansen, P. Wu, and V. V. Fokin, *J. Org. Chem.*, **70**, 7761 (2005); (f) F. Himo, T. Lovell, R. Hilgraf, V. V. Rostovtsev, L. Noodleman, K. B. Sharpless, and V. V. Fokin, *J. Am. Chem. Soc.*, **127**, 210 (2005); (g) A. Dondoni, P. P. Giovannini, and A. Massi, *Org. Lett.*, **6**, 2929 (2004); (h) Y. Ye, Y. Zheng, G. Y. Xu, and L. Z. Liu, *Heteroatom Chem.*, **14**, 254 (2003); (i) D. Muri, N. Lohse-Fraefel, and E. M. Carreira, *Angew. Chem. Int. Ed.*, **44**, 4036 (2005).
- (a) A. I. Kotyatkina, V. N. Zhabinsky, and V. A. Khripach, *Uspekhi Khimii*, **70**, 730 (2001); (b) Z. X. Yu and K. N. Houk, *J. Am. Chem. Soc.*, **125**, 13825 (2003); (c) D. Giguere, R. Patnam,

- M. A. Bellefleur, C. St-Pierre, S. Sato, and R. Roy, *Chem. Commun.*, **22**, 2379 (2006); (d) M. Benlifa, S. Vidal, D. Gueyrard, P. G. Goekjian, M. Msaddek, and J. P. Praly, *Tetrahedron Lett.*, **47**, 6143 (2006).
4. K. N. Anisimov and A. N. Nesmeyanov, *Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya*, 610 (1954).
 5. K. N. Anisimov, N. E. Kolobova, and A. N. Nesmeyanov, *Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya*, 240 (1955).
 6. R. R. Shagidullin, V. A. Pavlov, B. I. Buzykin, N. V. Aristova, and L. F. Chertanova, *Zh.Obshch.Khim.*, **61**, 1590 (1991).
 7. M. M. Crutchfield, C. H. Dungan, J. H. Letcher, V. Mark, and J. R. Van Wazer, *Topics in Phosphorus Chemistry* (John Wiley & Sons Inc., New York, 1967), vol. 5.
 8. W. Rosch, Z. Facklown, and M. Regitz, *Tetrahedron*, **43**, 222 (1987).
 9. A. D. H. Becke, J. P. Morken, A. F. Hour, and Z. Xu, *J. Am. Chem. Soc.*, **114**, 6692 (1992).
 10. T. Ziegler, *Chem. Rev.*, **91**, 651 (1991).
 11. R. G. Parr and W. Yang, *Density Functional Theory of Atoms and Molecules* (Oxford University Press, New York, 1989).
 12. C. Lee, W. Yang, and R. Parr, *Phys. Rev. B*, **37**, 785 (1988).
 13. B3LYP calculations give relative energies of the various structural transition to be within ~5 kcal/mol of the actual energies: (a) A. Ricca and C. W. Bauschlicher, *Theor. Chim. Acta*, **92**, 123 (1995); (b) M. R. A. Blomberg, P. E. M. Siegbahn, and M. Svensson, *J. Chem. Phys.*, **104**, 9546 (1996); (c) A. Ricca and C. W. Bauschlicher, *J. Phys. Chem.*, **101**, 8949 (1997); (d) M. N. Glukhovstev, R. D. Bach, and C. J. Nagel, *J. Phys. Chem.*, **101**, 316 (1997); (e) B. D. Dunietz, Y. Cao, D. A. Whittington, S. J. Lippard, and R. A. Friesner, *J. Am. Chem. Soc.*, **122**, 2828 (2000).