

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Synthesis and Properties of Pentacoordinated Phenoxysilane and Carboxysilanes with Intramolecular Nitrogen-Silicon Coordination

Yuya Domoto^a; Kowichiro Saruhashi^b; Akihiro Fukushima^b; Shohei Sase^a; Kei Goto^a; Takayuki Kawashima^b

^a Interactive Research Center of Science, Graduate School of Science and Engineering, Tokyo Institute of Technology, Tokyo, Japan ^b Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo, Japan

Online publication date: 27 May 2010

To cite this Article Domoto, Yuya , Saruhashi, Kowichiro , Fukushima, Akihiro , Sase, Shohei , Goto, Kei and Kawashima, Takayuki(2010) 'Synthesis and Properties of Pentacoordinated Phenoxysilane and Carboxysilanes with Intramolecular Nitrogen-Silicon Coordination', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185: 5, 1221 – 1229

To link to this Article: DOI: 10.1080/10426501003773746

URL: <http://dx.doi.org/10.1080/10426501003773746>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS AND PROPERTIES OF PENTACOORDINATED PHENOXYLSILANE AND CARBOXYLSILANES WITH INTRAMOLECULAR NITROGEN–SILICON COORDINATION

Yuya Domoto,¹ Kowichiro Saruhashi,² Akihiro Fukushima,²
Shohei Sase,¹ Kei Goto,¹ and Takayuki Kawashima²

¹Interactive Research Center of Science, Graduate School of Science and Engineering, Tokyo Institute of Technology, Tokyo, Japan

²Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo, Japan

Reactions of a neutral pentacoordinated monohydrosilane bearing a dative N–Si bond with phenol and benzoic acid derivatives gave the corresponding pentacoordinated phenoxysilane and carboxysilanes, respectively. X-ray crystallographic analyses of these silanes revealed that the distance of the N–Si dative bond is shortened and the pentacoordination character of the silicon center becomes greater with the increase in the electron-withdrawing character of the apical substituent on silicon.

Keywords Carboxysilanes; intramolecular coordination; pentacoordinated silicon compounds; phenoxysilanes

INTRODUCTION

Highly coordinated silicon compounds are known to exhibit distinct difference in reactivities, structures, and properties from their tetracoordinated counterparts.¹ Among them, much attention has been paid to neutral pentacoordinated silicon compounds with an intramolecular N–Si dative bond for examination of their intriguing structures and reactivities that originate from the characteristics of the dative bond.² In this class of compounds,

Received 11 February 2009; accepted 24 February 2009.

Dedicated to Professor Naomichi Furukawa on the occasion of his 70th birthday.

This work was partly supported by Grants-in-Aid for The 21st Century COE Program for Frontiers in Fundamental Chemistry (T.K.), for The Global COE Program for Education and Research Center for Emergence of New Molecular Chemistry (K.G. and S.S.), and for Scientific Research (K.G. and T.K.) from the Ministry of Education, Culture, Sports, Science and Technology, Japan. K.G. acknowledges the support by Tokuyama Science Foundation. We also thank Tosoh Finechem Corporation and Shin-Etsu Chemical Co., Ltd. for the generous gifts of alkylolithiums and silicon compounds, respectively.

Address correspondence to Kei Goto, Interactive Research Center of Science, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1 O-okayama, Meguro-ku, Tokyo 152-8551, Japan. E-mail: goto@chem.titech.ac.jp and Takayuki Kawashima, Department of Chemistry, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. E-mail: takayuki@chem.s.u-tokyo.ac.jp





Figure 1 Schematic representation of pentacoordinated silanes with monocyclic (I), bicyclic (II), and tricyclic (III) motifs.

monocyclic,^{2a,b} bicyclic,^{2c,d} or tricyclic frameworks,^{2e,f} have been utilized to hold the nitrogen and silicon atoms in proximity (Figure 1), and it was found that the N–Si interaction is sensitive to the electronic character of the substituents on the silicon atom. To date, extensive structural investigations concerning pentacoordinated silicon compounds with a tricyclic motif such as silatranes have been undertaken, which have revealed that electron-withdrawing substituents on the silicon shorten the N–Si distance.³ For pentacoordinated silanes with the mono- and bicyclic frameworks, however, systematic study from this viewpoint is still missing in spite of extensive exploration of their reactivities. Recently, we have reported the synthesis of neutral pentacoordinated hydrosilanes with a bicyclic motif based on the dibenzosilazocine framework^{4a} and application to the construction of a tweezer-type cavitand.^{4b} This type of bicyclic motif has been successfully utilized for the investigation of the substituent effects on highly coordinated main group elements.⁵ Several compounds bearing the dibenzosilazocine framework have been reported by Carré et al.,^{2c} although crystallographic analysis has been performed only for dimethyl- and difluoro-substituted derivatives. In this article, we report the synthesis of neutral pentacoordinated phenoxy- and carboxysilanes by the reaction of a pentacoordinated hydrosilane with phenol and benzoic acid derivatives, respectively, along with their structural characterization.

RESULTS AND DISCUSSION

Treatment of neutral pentacoordinated hydrosilane **1**^{4a} with phenol in chloroform under reflux for 13 h gave the corresponding phenoxysilane **2** in 84% yield. The reaction of **1** with benzoic acid proceeded under milder conditions, and silyl benzoate **3** was formed after 18 h at room temperature, which was isolated in 66% yield. It is of note that salicylic acid reacts with **1** much faster than benzoic acid. The reaction was completed within 10 min at room temperature, affording silyl salicylate **4** in 87% yield. Compounds **2–4** (Scheme 1) were unambiguously characterized by NMR spectroscopy (¹H, ¹³C, ²⁹Si) and elemental analyses. The ²⁹Si NMR (CDCl₃) resonances of **2–4** appear at –40.3, –44.1, and –46.1 ppm, respectively, the high-field shifts of which are typical for pentacoordinated silicon compounds.⁶

While there have been extensive studies on pentacoordinated aryloxy- and carboxysilanes with intramolecular coordination of an amine unit, their structural properties have been hardly explored, and crystallographic analyses of the compounds bearing a monodentate aryloxy or carboxy ligand have been performed only for silatrane derivatives.⁷ The structures of phenoxysilane **2** and carboxysilanes **3** and **4** were established by X-ray crystallographic analyses. Figure 2 shows the ORTEP drawing of **2**. The selected bond lengths and angles of **2** are given in Table I.



Scheme 1

In the crystal structure of phenoxysilane **2**, there are two independent molecules in the asymmetric unit. The silicon atoms of both fragments adopt slightly distorted trigonal bipyramidal structures with the nitrogen and oxygen atoms at apical positions and three carbon atoms at equatorial positions. The N—Si distances of **2** [2.439(4) and 2.419(4) Å for




Figure 2 ORTEP drawing of **2** (50% probability). Hydrogen atoms are omitted for clarity.

Table I Selected bond lengths (Å) and angles (deg) for **2**

Bond lengths		Bond angles			
Si(1)—N(1)	2.439(4)	N(1)—Si(1)—O(1)	167.72(14)	N(2)—Si(2)—O(2)	168.02(15)
Si(1)—O(1)	1.696(3)	N(1)—Si(1)—C(1)	77.19(17)	N(2)—Si(2)—C(5)	77.46(16)
Si(1)—C(1)	1.880(5)	N(1)—Si(1)—C(2)	76.64(16)	N(2)—Si(2)—C(6)	76.82(17)
Si(1)—C(2)	1.890(5)	N(1)—Si(1)—C(3)	88.03(15)	N(2)—Si(2)—C(7)	87.33(16)
Si(1)—C(3)	1.873(4)	O(1)—Si(1)—C(1)	100.83(19)	O(2)—Si(2)—C(5)	94.08(18)
O(1)—C(4)	1.360(5)	O(1)—Si(1)—C(2)	94.71(18)	O(2)—Si(2)—C(6)	101.10(18)
Si(2)—N(2)	2.419(4)	O(1)—Si(1)—C(3)	103.73(16)	O(2)—Si(2)—C(7)	104.29(17)
Si(2)—O(2)	1.692(3)	C(1)—Si(1)—C(3)	113.3(2)	C(5)—Si(2)—C(7)	118.73(19)
Si(2)—C(5)	1.882(4)	C(1)—Si(1)—C(2)	122.22(19)	C(5)—Si(2)—C(6)	121.7(2)
Si(2)—C(6)	1.884(5)	C(2)—Si(1)—C(3)	116.2(2)	C(6)—Si(2)—C(7)	111.29(19)
Si(2)—C(7)	1.885(4)	Si(1)—O(1)—C(4)	138.0(3)	Si(2)—O(2)—C(8)	140.2(3)
O(2)—C(8)	1.352(5)				

each fragment] are much shorter than the sum of van der Waals radii of silicon and nitrogen (3.65 Å),⁸ proving strong interaction between the two atoms in the crystalline state, although they are longer than the values observed for silatranes bearing an aryloxy ligand (2.07–2.11 Å).^{7a,b} Carré et al. reported the crystal structure of a pentacoordinated difluorosilane with the same type of dibenzosilazocine framework,^{2c} the N—Si distance of which [2.430 Å (average)] is close to those of **2**. The Si—O bond lengths of **2** [1.696(3) and 1.692(3) Å for each fragment] are slightly longer than that of (2-bromophenoxy)triphenylsilane (1.645(4) Å),⁹ indicating the hypervalent character of the Si—O bond of **2**.

The ORTEP drawings of carboxysilanes **3** and **4** are shown in Figures 3 and 4, respectively. The selected bond lengths and angles of **3** and **4** are given in Table II. In silyl benzoate **3**, the N—Si distance is 2.315(2) Å, which is much shorter than that of **2**. Silyl salicylate **4** showed even shorter N—Si distance (2.2865(17) Å), although it is still longer than those of silatranes bearing an carboxy ligand (2.05–2.07 Å).^{7c,d} There have been several examples of pentacoordinated silanes with a salicylate ligand,¹⁰ but in all cases the salicylate serves as an *O,O*-bidentate ligand to form a cyclic structure. In contrast, a monodentate coordination of the salicylate ligand is clearly shown in the crystal structure of **4**, where the *o*-OH group is located in the far side of the silicon atom.

The pentacoordination characters (%TBP_a/°TBP_e)¹¹ and the N—Si bond distances of **2–4** are summarized in Table III along with their ²⁹Si NMR chemical shifts. The geometry around the silicon center becomes closer to the ideal TBP as the electron-withdrawing ability of the substituent at the apical position is higher.¹² The N—Si bond distance also depends on the electronic nature of the apical substituent and is shortened as the pentacoordination character becomes greater, as has been observed for the analogues with other main group element bearing the same type of framework.⁵ In the ²⁹Si NMR spectra, small but apparent high-field shifts were observed as the apical substituent becomes electron-withdrawing, showing the increase in the degree of pentacoordination. These results indicate that the correlation between pentacoordination character and electronic nature of the apical substituent is preserved also in solution.

In summary, we have synthesized neutral pentacoordinated phenoxy- and carboxysilanes with a N—Si dative bond bearing a bicyclic framework. We have also performed a systematic structural analysis of the phenoxy- and carboxysilanes and revealed that, in analogy with neutral pentacoordinated organosilanes with a tricyclic motif such as silatranes,




Figure 3 ORTEP drawing of **3** (50% probability). Hydrogen atoms are omitted for clarity.

Table II Selected bond lengths (Å) and angles (deg) for **3** and **4**

3		4	
Bond lengths			
Si(1)—N(1)	2.315(2)	Si(1)—N(1)	2.2865(17)
Si(1)—O(1)	1.768(2)	Si(1)—O(1)	1.7772(14)
Si(1)—C(1)	1.886(3)	Si(1)—C(1)	1.880(2)
Si(1)—C(2)	1.881(3)	Si(1)—C(2)	1.880(2)
Si(1)—C(3)	1.878(2)	Si(1)—C(3)	1.868(2)
O(1)—C(4)	1.335(3)	O(1)—C(4)	1.314(2)
C(4)—O(2)	1.219(3)	C(4)—O(2)	1.236(2)
Bond angles			
N(1)—Si(1)—O(1)	175.86(7)	N(1)—Si(1)—O(1)	171.25(7)
N(1)—Si(1)—C(1)	79.07(10)	N(1)—Si(1)—C(1)	79.49(8)
N(1)—Si(1)—C(2)	78.74(10)	N(1)—Si(1)—C(2)	79.61(8)
N(1)—Si(1)—C(3)	90.73(10)	N(1)—Si(1)—C(3)	89.37(8)
O(1)—Si(1)—C(1)	99.70(11)	O(1)—Si(1)—C(1)	93.45(8)
O(1)—Si(1)—C(2)	98.84(11)	O(1)—Si(1)—C(2)	100.03(8)
O(1)—Si(1)—C(3)	93.37(10)	O(1)—Si(1)—C(3)	98.55(8)
C(1)—Si(1)—C(3)	114.05(11)	C(1)—Si(1)—C(3)	117.64(10)
C(1)—Si(1)—C(2)	123.46(10)	C(1)—Si(1)—C(2)	121.50(9)
C(2)—Si(1)—C(3)	117.53(12)	C(2)—Si(1)—C(3)	116.07(9)
Si(1)—O(1)—C(4)	127.18(16)	Si(1)—O(1)—C(4)	127.82(13)
O(1)—C(4)—O(2)	123.3(3)	O(1)—C(4)—O(2)	123.07(18)




Figure 4 ORTEP drawing of **4** (50% probability). Hydrogen atoms, except for that of the hydroxy group, are omitted for clarity.

the dative N—Si bond distance is shortened with the increase in the electron-withdrawing character of the substituent at the apical position, and the pentacoordination character at the silicon center concomitantly becomes greater.

EXPERIMENTAL

All reactions were carried out under an argon atmosphere. Melting points were determined on a Yanaco micro melting point apparatus. All melting points were uncorrected. Chloroform was distilled from calcium hydride under an Ar atmosphere before use. ^1H , ^{13}C , and ^{29}Si NMR spectra were recorded on a Bruker DRX-500 spectrometer. ^1H and ^{13}C chemical shifts were referenced using a residual peak of CDCl_3 (^1H , δ_{H} 7.25, ^{13}C , δ_{C} 77.0) as an

Table III Selected structural parameters and ^{29}Si NMR (99 MHz, CDCl_3) data of **2**, **3**, and **4**

	N—Si/Å	%TBP _e / %TBP _a	δ_{Si}
2	2.439(4)	74/50	−40.3
3	2.315(2)	84/63	−44.1
4	2.2865(17)	85/62	−46.1

internal standard. ^{29}Si NMR chemical shifts are reported relative to Me_4Si (0 ppm) in ppm. IR spectra were recorded on JASCO V-650. Gel permeation liquid chromatography (GLPC) was performed by LC-918 and LC-9204 with JAIGEL 1H+2H columns (Japan Analytical Industry) with chloroform as solvent. Elemental analyses were performed by the Micro-analytical Laboratory of the Department of Chemistry, Faculty of Science, the University of Tokyo, or on a Perkin-Elmer PE2400 Series II CHNS/O Analyzer. Pentacoordinated hydrosilane **1** was prepared according to the reported procedure.^{4a}

Synthesis of Phenoxysilane 2

To the mixture of monohydrosilane **1** (116 mg, 0.325 mmol) and phenol (31 mg, 0.33 mmol), CHCl_3 (3 mL) was added. After stirring for 13 h at reflux temperature, the solvent was removed, and the residue was purified by GLPC to give phenoxysilane **2** as a colorless solid (123 mg, 0.272 mmol, 84%). Mp 78–80°C; ^1H NMR (500 MHz, CDCl_3) δ 0.65 (t, $^3J_{\text{HH}} = 7.3$ Hz, 3H), 0.83–0.87 (m, 2H), 1.17–1.23 (m, 2H), 1.44–1.46 (m, 2H), 3.68 (d, $^2J_{\text{HH}} = 14.7$ Hz, 2H), 3.74 (d, $^2J_{\text{HH}} = 14.7$ Hz, 2H), 6.70 (tt, $^3J_{\text{HH}} = 7.3$ Hz, $^4J_{\text{HH}} = 1.1$ Hz, 1H), 6.75 (dd, $^3J_{\text{HH}} = 8.7$ Hz, $^4J_{\text{HH}} = 1.1$ Hz, 2H), 6.98–7.02 (m, 2H), 7.14–7.21 (m, 5H), 7.26–7.28 (m, 2H), 7.34–7.40 (m, 4H), 8.24–8.26 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ 13.6 (q), 20.5 (t), 24.5 (t), 52.3 (t), 55.6 (t), 119.2 (d), 120.1 (d), 125.6 (d), 127.3 (d), 127.4 (d), 128.1 (d), 128.7 (d), 129.9 (d), 132.3 (d), 133.0 (s), 136.7 (s), 138.5 (d), 146.5 (s), 157.6 (s); $^{29}\text{Si}\{^1\text{H}\}$ NMR (99 MHz, CDCl_3) δ –40.3. Anal. Calcd. For $\text{C}_{30}\text{H}_{31}\text{NOSi}$: C, 80.13; H, 6.95; N 3.11. Found: C, 80.07; H, 6.97; N, 2.88.

Synthesis of Carboxysilane 3

To the mixture of monohydrosilane **1** (288 mg, 0.806 mmol) and benzoic acid (99 mg, 0.81 mmol), CHCl_3 (4 mL) was added. After stirring for 18 h at room temperature, the solvent was removed and the residue was purified by GLPC to give carboxysilane **3** as colorless solid (253 mg, 0.529 mmol, 66%). Mp 65–67°C; ^1H NMR (500 MHz, CDCl_3) δ 0.67 (t, $^3J_{\text{HH}} = 7.3$ Hz, 3H), 0.84–0.90 (m, 2H), 1.21–1.28 (m, 2H), 1.48–1.51 (m, 2H), 3.72 (d, $^2J_{\text{HH}} = 14.6$ Hz, 2H), 3.76 (d, $^2J_{\text{HH}} = 14.6$ Hz, 2H), 7.18–7.26 (m, 7H), 7.36–7.43 (m, 7H), 7.51 (tt, $^3J_{\text{HH}} = 7.4$ Hz, $^4J_{\text{HH}} = 1.4$ Hz, 1H), 8.10–8.15 (m, 4H); ^{13}C NMR (126 MHz, CDCl_3) δ 13.6 (q), 20.5 (t), 24.4 (t), 52.6 (t), 55.5 (t), 125.5 (d), 127.3 (d), 127.5 (d), 128.1 (d), 128.3 (d), 130.1 (d), 130.1 (d), 131.6 (s), 132.0 (d), 132.3 (d), 133.5 (s), 136.5 (s), 137.8 (d), 145.8 (s), 166.7 (s); $^{29}\text{Si}\{^1\text{H}\}$ NMR (99 MHz, CDCl_3) δ –44.1; IR (KBr) $\nu_{\text{C=O}}$ 1687 cm^{-1} . Anal. Calcd. For $\text{C}_{31}\text{H}_{31}\text{NO}_2\text{Si}$: C, 77.95; H, 6.54; N 2.93. Found: C, 77.74; H, 6.83; N, 2.66.

Synthesis of Carboxysilane 4

The carboxysilane **4** (374 mg, 0.758 mmol, 87%) was prepared from monohydrosilane **1** (312 mg, 0.873 mmol) and salicylic acid (122 mg, 0.883 mmol) by the same procedure as the synthesis of **3**, while the reaction time was 0.5 h. Mp 123–125°C; ^1H NMR (500 MHz, CDCl_3) δ 0.66 (t, $^3J_{\text{HH}} = 7.3$ Hz, 3H), 0.85–0.90 (m, 2H), 1.23–1.29 (m, 2H), 1.53–1.56 (m, 2H), 3.76 (d, $^2J_{\text{HH}} = 14.7$ Hz, 2H), 3.80 (d, $^2J_{\text{HH}} = 14.7$ Hz, 2H), 6.88 (dt, $^3J_{\text{HH}} = 7.9$ Hz, $^4J_{\text{HH}} = 1.0$ Hz, 1H), 6.92 (dd, $^3J_{\text{HH}} = 7.9$ Hz, $^4J_{\text{HH}} = 1.0$ Hz, 1H), 7.20–7.29 (m, 5H), 7.37–7.44 (m, 7H), 8.05–8.07 (m, 3H), 11.48 (s, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ 13.5 (q), 20.4 (t), 24.5 (t), 52.7 (t), 55.6 (t), 115.4 (s), 117.2 (d), 118.6 (d), 125.6 (d), 127.5 (d),

127.7 (d), 128.5 (d), 130.3 (d), 130.8 (d), 131.2 (s), 132.2 (d), 134.7 (d), 136.2 (s), 137.6 (d), 145.5 (s), 162.2 (s), 171.0 (s); $^{29}\text{Si}\{^1\text{H}\}$ NMR (99 MHz, CDCl_3) δ -46.1; IR (KBr) $\nu_{\text{C=O}}$ 1606, 1647 cm^{-1} . Anal. Calcd. For $\text{C}_{31}\text{H}_{31}\text{NO}_3\text{Si}$: C, 75.42; H, 6.33; N 2.84, Found C, 75.58; H, 6.57; N, 2.64.

X-Ray Crystallographic Analysis of 2, 3, and 4

Single crystals of **2**, **3**, and **4** were grown in their chloroform/hexane solutions. The intensity data were collected at 120 K on a Rigaku/MS Mercury CCD diffractometer with graphite-monochromated MoK_α radiation ($\lambda = 0.71069$ Å). Crystallographic and experimental data are listed in Table IV. The structures were solved by the direct method and refined by full-matrix least squares on F^2 using SHELXL 97.¹³ The nonhydrogen atoms were refined anisotropically. The hydrogen atoms were idealized by using the riding models except for the OH hydrogen of **4**, which was located in the difference Fourier map and refined isotropically. Crystallographic data for the structural analysis has been deposited with the Cambridge Crystallographic Data Centre, CCDC Nos. 721463, 721464, and 721465 for **2**, **3**, and **4** respectively. Copies of this information may be obtained free of charge from the director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK (fax: +44-1223-336-033; e-mail: deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>).

Table IV Crystallographic data for **2**, **3**, and **4**

	2	3	4
Formula	$\text{C}_{30}\text{H}_{31}\text{NOSi}$	$\text{C}_{31}\text{H}_{31}\text{NO}_2\text{Si}$	$\text{C}_{31}\text{H}_{31}\text{NO}_3\text{Si}$
Temperature/K	120(2)	123(2)	120(2)
Crystal system	Triclinic	Monoclinic	Monoclinic
Space group	$P1$	$P2_1/c$	$P2_1/c$
$a/\text{\AA}$	9.628(5)	9.081(6)	8.6593(18)
$b/\text{\AA}$	10.143(4)	18.032(10)	17.938(4)
$c/\text{\AA}$	13.930(6)	15.749(10)	17.321(4)
α/deg	66.272(14)		
β/deg	82.83(2)	95.935(13)	102.809(3)
γ/deg	88.67(2)		
$V/\text{\AA}^3$	1235.2(10)	2565(3)	2623.4(9)
Z	2	4	4
Calculated density/ g cm^{-3}	1.209	1.237	1.250
Reflections collected	8093	10995	17116
Unique	6056	3720	4614
R_{int}	0.0326	0.0447	0.0703
F_{000}	480	1016	1048
Limiting indices	$-11 \leq h \leq 11$ $-12 \leq k \leq 11$ $-16 \leq l \leq 15$	$-7 \leq h \leq 10$ $-19 \leq k \leq 20$ $-18 \leq l \leq 16$	$-10 \leq h \leq 7$ $-21 \leq k \leq 21$ $-20 \leq l \leq 20$
Restraints/parameters	3/597	0/317	0/330
Goodness of fit (F^2)	1.080	1.079	1.001
R indices ($I > 2\sigma(I)$)	$R1 = 0.0566$ $wR2 = 0.1298$	$R1 = 0.0467$ $wR2 = 0.1084$	$R1 = 0.0478$ $wR2 = 0.1132$
R indices (all data)	$R1 = 0.0613$ $wR2 = 0.1336$	$R1 = 0.0711$ $wR2 = 0.1243$	$R1 = 0.0594$ $wR2 = 0.1270$

REFERENCES

1. (a) C. Chuit, R. J. P. Corriu, C. Reye, and J. C. Young, *Chem. Rev.*, **93**, 1371 (1993); (b) R. R. Holmes, *Chem. Rev.*, **90**, 17 (1990); (c) R. R. Holmes, *Chem. Rev.*, **96**, 927 (1996).
2. (a) J. Boyer, C. Brelriere, R. J. P. Corriu, A. Kpoton, M. Poirier, and G. Royo, *J. Organomet. Chem.*, **311**, C39 (1986); (b) P. Arya, R. J. P. Corriu, K. Gupta, G. F. Lanneau, and Z. Yu, *J. Organomet. Chem.*, **399**, 11 (1990); (c) F. H. Carré, R. J. P. Corriu, G. F. Lanneau, P. Merle, F. Soulaïrol, and J. Yao, *Organometallics*, **16**, 3878 (1997); (d) W. F. Paton, E. R. Corey, and J. Y. Corey, *Acta Crystallogr., Sect. B: Struct. Sci.*, **33**, 3322 (1977); (e) S. N. Tanduar, M. G. Voronkov, and N. V. Alekseev, *Top. Curr. Chem.*, **131**, 99 (1986); (f) R. J. P. Corriu, *J. Organomet. Chem.*, **400**, 81 (1990).
3. (a) A. A. Kemme, J. J. Bleidelis, V. A. Pestunovich, V. P. Baryshok, and M. G. Voronkov, *Dokl. Akad. Nauk. SSSR*, **243**, 688 (1978); (b) M. G. Voronkov, V. M. D'yakov, and S. V. Kirpichenko, *J. Organomet. Chem.*, **233**, 1 (1982).
4. (a) K. Saruhashi, K. Goto, and T. Kawashima, *Chem. Heterocycl. Compd.*, **37**, 1394 (2001); (b) A. Yoshida, K. Goto, and T. Kawashima, *Bull. Chem. Soc. Jpn.*, **79**, 793 (2006).
5. (a) For examples, see: (a) K.-y. Akiba, K. Takee, K. Ohkata, and F. Iwasaki, *J. Am. Chem. Soc.*, **105**, 6965 (1983); (b) K.-y. Akiba, K. Takee, Y. Shimizu, and K. Ohkata, *J. Am. Chem. Soc.*, **108**, 6320 (1986); (c) H. Fujihara, H. Mima, M. Ikemori, and N. Furukawa, *J. Am. Chem. Soc.*, **113**, 6337 (1991); (d) S. Shimada, O. Yamazaki, T. Tanaka, Y. Suzuki, and M. Tanaka, *J. Organomet. Chem.*, **689**, 3012 (2004).
6. For example, see B. J. Helmer, R. West, R. J. P. Corriu, M. Poirier, G. Royo, and A. De Saxe, *J. Organomet. Chem.*, **251**, 295 (1983).
7. (a) L. Parkanyi, P. Hencsei, and L. Bihatsi, *J. Organomet. Chem.*, **233**, 315 (1982); (b) I. Kovacs, P. Hencsei, and L. Parkanyi, *Period. Polytech. Chem. Eng.*, **31**, 155 (1987); (c) L. Xiuyun, Z. Qitai, W. Shen, Z. Shude, S. Fuling, W. Jitao, X. Qinglan, and L. Jing, *Gaodeng Xuexiao Huaxue Xuebao*, **8**, 629 (1987); (d) L. Chen, Q. Xie, L. Sun, and H. Wang, *J. Organomet. Chem.*, **678**, 90 (2003).
8. L. Pauling, *The Nature of the Chemical Bond and the Structure of Molecules and Crystals: An Introduction to Modern Structural Chemistry*, 3rd ed. (Cornell University Press, Ithaca, NY, 1960).
9. A. Studer, M. Bossart, and T. Vasella, *Org. Lett.*, **2**, 985 (2000).
10. (a) R. Tacke, J. Heermann, M. Pülm, and I. Richter, *Organometallics*, **17**, 1663 (1998); (b) O. Seiler, C. Burschka, M. Penka, and R. Tacke, *Silicon Chem.*, **1**, 355 (2002).
11. K. Tamao, T. Hayashi, Y. Ito, and M. Shiro, *Organometallics*, **11**, 2099 (1992).
12. For example, R. J. Garant, L. M. Daniels, S. K. Das, M. N. Janakiraman, R. A. Jacobson, and J. G. Verkade, *J. Am. Chem. Soc.*, **113**, 5728 (1991).
13. G. M. Sheldrick, SHELXL-97. Computer Program for Crystal Structure Refinement, University of Göttingen, Göttingen, Germany (1997).