


This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Synthesis and Antiviral Activities of α -Aminophosphonate Derivatives Containing a Pyridazine Moiety

Jun Zhou^a; Hui-Tao Fan^a; Bao-An Song^a; Lin-Hong Jin^a; Pinaki S. Bhadury^a; De-Yu Hu^a; Song Yang^a

^a Center for Research and Development of Fine Chemicals, State Key Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering and Key Laboratory of Green Pesticide & Agricultural Bioengineering, Ministry of Education, Guizhou University, Guiyang, People's Republic of China

Online publication date: 13 January 2011

To cite this Article Zhou, Jun , Fan, Hui-Tao , Song, Bao-An , Jin, Lin-Hong , Bhadury, Pinaki S. , Hu, De-Yu and Yang, Song(2011) 'Synthesis and Antiviral Activities of α -Aminophosphonate Derivatives Containing a Pyridazine Moiety', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 186: 1, 81 – 87

To link to this Article: DOI: 10.1080/10426507.2010.482543

URL: <http://dx.doi.org/10.1080/10426507.2010.482543>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.


The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS AND ANTIVIRAL ACTIVITIES OF α -AMINOPHOSPHONATE DERIVATIVES CONTAINING A PYRIDAZINE MOIETY

Jun Zhou, Hui-Tao Fan, Bao-An Song, Lin-Hong Jin, Pinaki S. Bhadury, De-Yu Hu, and Song Yang

Center for Research and Development of Fine Chemicals, State Key Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering and Key Laboratory of Green Pesticide & Agricultural Bioengineering, Ministry of Education, Guizhou University, Guiyang, People's Republic of China

GRAPHICAL ABSTRACT



Abstract The title compounds (**4**) were obtained by the reaction of 2-phenyl-4-chloro-5-(4-aminophenyl)pyridazin-3(2H)-one, aldehydes, and O,O'-dialkyl phosphite under reflux conditions using dry toluene as a solvent. The preliminary bioassays indicate that the synthesized compounds have moderate to good against tobacco mosaic virus (TMV) activity. It was found that compound **4b** had a good inactivation effect *in vivo* against TMV, with an inhibition rate of 81.0%.

Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.

Keywords α -Aminophosphonate derivatives; antiviral activity; pyridazine moiety; synthesis

INTRODUCTION

Pyridazine derivatives are known for their wide range of biological activities as insecticides,¹ fungicides,² plant virucides,³ and as cardiotonics.⁴ They also serve as important

Received 2 February 2010; accepted 20 March 2010.

We are grateful for the generous financial support received from the National Key Program for Basic Research (No. 2010CB126105) and the National Natural Science Foundation of China (No 20872002).

Address correspondence to Bao-An Song, Center for Research and Development of Fine Chemicals, State Key Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering and Key Laboratory of Green Pesticide & Agricultural Bioengineering, Ministry of Education, Guizhou University, Guiyang 550025, People's Republic of China. E-mail: songbaoan22@yahoo.com

synthetic precursors to many active molecules due to the ease with which the four ring carbons can be functionalized. Another important class of compounds that belongs to the α -aminophosphonic acid group has been routinely employed as growth regulators,⁵ fungicides,⁶ plant virucides⁷ and herbicides.⁸ A great deal of research has been conducted in recent years on the development of suitable synthetic methodologies for biologically active α -aminophosphonates and their derivatives.^{9–12} Among the various synthetic methods,¹³ catalytic hydrophosphonylation of preformed imine by dialkyl phosphite is of particular interest, but this method suffers from lack of applicability to sensitive imines. To address the latter issue, and in our pursuit to prepare fluorinated heterocyclic α -aminophosphonates with potential antiviral activity against tobacco mosaic virus (TMV) *in vivo*, we had employed *in situ*-generated imine in a three-component, one-pot, Mannich-type reaction involving aldehyde, amine, and dialkyl phosphite.¹⁴ In this article, we report the preparation of new pyridazin-3(2*H*)-one derivatives bearing α -aminophosphonate moiety and their subsequent evaluation as anti-TMV agents.


RESULTS AND DISCUSSION

4,5-Dichloro-2-phenyl-3-pyridazone was synthesized by following a published reaction of mucochloric acid with phenylhydrazine. The reaction of compound **2** with 4-aminothiophenol and anhydrous potassium carbonate in acetone then afforded 2-phenyl-4-chloro-5-(4-aminophenylthio) pyridazin-3(2*H*)-one **3**, which was finally reacted in a three-component reaction with different aldehydes and dialkyl phosphites to provide title α -aminophosphonates **4** (Scheme 1). Synthesis of **4** was executed in different solvents under varying conditions using Lewis acid catalysts, ionic liquids, microwave irradiation, and conventional heating. Under optimized conditions, the best result was obtained when the Mannich reaction was carried out for 7 h in dry toluene at its reflux temperature. The products **4a–l** were obtained in good to high chemical yield (51–79%).

The chemical structures of compounds **3** and **4a–l** were confirmed by their spectral data and elemental analysis. In the IR spectrum of **3**, the characteristic stretching frequency of N–H appeared at 3437–3342 cm^{–1}, whereas pyridazinone C=O appeared at 1653 cm^{–1}. The chemical shifts in ¹H NMR spectrum showed a singlet at δ 7.09 due to the presence of an amino group and a multiplet in the region δ 6.65–7.53 for aromatic protons. The products **4a–l** exhibited in their IR spectra the characteristic absorption bands at 3267–3300 cm^{–1} (N–H) and at 1660–1670 cm^{–1} (C=O). The ¹H NMR spectral data consisted of aromatic signals (δ 6.62–7.54) and chemical shifts near δ 4.72–4.78 appearing as a doublet of doublets due to the CH moiety adjacent to a phosphorus atom. The ³¹P NMR spectral data showed the presence of a phosphorus atom and appeared around δ 19.85–23.39 as singlets or doublets.

EXPERIMENTAL

The melting points of the products were determined on a XT-4 binocular microscope (Beijing Tech Instrument Co., China) and were not corrected. The IR spectra were recorded on a Bruker Vector 22 spectrometer in KBr disks. ¹H, ¹³C, ³¹P, and ¹⁹F NMR (solvent CDCl₃ or D₃CCOCD₃) spectra were performed on a JEOL-ECX 500 NMR spectrometer at room temperature using TMS as internal standard. Elemental analysis was performed on an Elementar Vario-III CHN analyzer. Analytical TLC was performed on silica gel GF254. Column chromatographic purification was carried out using silica gel.



Scheme 1 Synthetic route to title compounds 4.

Dialkyl phosphites were prepared according to the method as described in the literature.¹⁵ Compound **2** was prepared according to the procedure in the literature.¹⁶ The antiviral activity assay and data for compounds **4a–l** can be found in the Supplemental Materials (available online).

2-Phenyl-4-chloro-5-(4-aminophenylthio)pyridazin-3(2H)-one **3**

A mixture of **2** (0.96 g, 4 mmol), 4-aminothiophenol (0.5 g, 4 mmol), and anhydrous potassium carbonate (0.83 g, 6 mmol) in dry acetone (30 mL) was stirred for 2 h at room temperature. After cooling, the reaction mixture was filtered. The solvent was evaporated under reduced pressure. The crude product was purified by chromatography on silica gel using a mixture of petroleum ether and ethyl acetate (V/V = 2:1) as the eluent to give the compound **3** (0.94 g, 71%), mp 164–166 °C; IR (ν cm⁻¹): 3437, 3342 (NH), 1653 (C=O), 1593, 1560, 1490; ¹H NMR (500 MHz, D₃CCOCD₃): δ 6.83–7.58 (m, 10H, Ar-H), 5.39 (s, 2H, NH₂); ¹³C NMR (125 MHz, D₃CCOCD₃): δ 154.54, 151.58, 145.00, 141.85, 137.44, 132.40, 128.52, 128.12, 126.13, 125.55, 115.53, 109.94; Anal. Calcd. for C₁₆H₁₂ClN₃OS: C, 58.27; H, 3.67; N, 12.74. Found: C, 58.48; H, 3.51; N, 12.58.

Preparation of Compound 4a–l

Compound **3** (0.20 g, 0.6 mmol) was first dissolved in dry toluene (10 mL). Aldehyde (0.6 mmol) and dialkyl phosphite (0.72 mmol) were added to the above solution, and then the mixture was refluxed for 7 h. The solvent was evaporated under reduced pressure. The crude product was purified by chromatography on silica gel using a mixture of petroleum ether and ethyl acetate (V/V = 2:1) as the eluent to afford the corresponding pure α -aminophosphonates.

O,O'-Diethyl-4-[[[2-phenyl-4-chloro-5-thio)pyridazin-3(2H)-one]](phenylamino)-(phenyl)methylphosphonate (4a). Mp 192–194 °C; yield 63%; IR (ν cm⁻¹): 3288 (NH), 2980, 1660 (C=O), 1593, 1519, 1490, 1234 (P=O), 1020 (P-O-C); ¹H NMR (500 MHz, CDCl₃): δ 6.65–7.53 (m, 15H, Ar-H), 5.21–5.25 (m, 1H, NH), 4.72 (dd, J = 24 Hz, 7.45 Hz, 1H, CH), 3.61–4.18 (m, 4H, 2OCH₂), 1.30 (t, J = 7.15 Hz, 3H, CH₃), 1.09 (t, J = 7.15 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 16.24, 16.28, 16.50, 16.55, 55.19, 56.38, 63.35, 63.39, 63.73, 63.79, 113.20, 115.05, 125.18, 127.34, 127.78, 127.82, 128.36, 128.43, 128.45, 128.78, 128.93, 133.16, 135.07, 137.42, 141.38, 144.78, 148.59, 148.71, 155.29; ³¹P NMR (200 MHz, CDCl₃): δ 22.48; Anal. Calcd for C₂₇H₂₇ClN₃O₄PS: C, 58.32; H, 4.89; N, 7.56; Found: C, 58.08; H, 4.58; N, 7.27.

O,O'-Dipropyl-4-[[[2-phenyl-4-chloro-5-thio)pyridazin-3(2H)-one]](phenylamino)-(phenyl)methylphosphonate (4b). Mp 196–199 °C; yield 58%; IR (ν cm⁻¹): 3292 (NH), 2968, 1664 (C=O), 1593, 1560, 1510, 1492, 1454, 1228 (P=O), 1004 (P-O-C); ¹H NMR (500 MHz, CDCl₃): δ 6.66–7.53 (m, 15H, Ar-H), 5.29–5.32 (m, 1H, NH), 4.74 (dd, J = 24 Hz, 7.45 Hz, 1H, CH), 3.50–4.05 (m, 4H, 2OCH₂), 1.64–1.71 (m, 2H, CH₂), 1.44–1.51 (m, 2H, CH₂), 0.91 (t, J = 7.45 Hz, 3H, CH₃), 0.78 (t, J = 7.45 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 9.94, 10.11, 23.74, 23.79, 23.95, 24.00, 68.79, 68.86, 69.02, 69.08, 112.97, 115.03, 125.19, 127.24, 127.82, 127.85, 128.37, 128.78, 128.90, 128.92, 133.18, 135.17, 137.40, 141.37, 144.86, 148.70, 148.80, 155.30; ³¹P NMR (200 MHz, CDCl₃): δ 22.36; Anal. Calcd for C₂₉H₃₁ClN₃O₄PS: C, 59.64; H, 5.35; N, 7.19; Found: C, 59.68; H, 5.27; N, 7.38.

O,O'-Diisopropyl-4-[[[2-phenyl-4-chloro-5-thio)pyridazin-3(2H)-one]](phenylamino)-(phenyl)methylphosphonate (4c). Mp 182–184 °C; yield 57%; IR (ν cm⁻¹): 3278 (NH), 2978, 1662 (C=O), 1593, 1560, 1516, 1490, 1228 (P=O), 999 (P-O-C); ¹H NMR (500 MHz, CDCl₃): δ 6.65–7.54 (m, 15H, Ar-H), 5.23 (t, J = 8.3 Hz, 1H, NH), 4.66–4.72 (m, 2H, NCH+CH), 4.42–4.46 (m, 1H, CH), 1.33 (d, J = 6.3 Hz, 3H, CH₃), 1.25 (d, J = 6.3 Hz, 3H, CH₃), 1.24 (d, J = 6.3 Hz, 3H, CH₃), 0.90 (d, J = 6.3 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 23.19, 23.24, 23.93, 23.96, 24.24, 24.27, 24.32, 24.34, 55.59, 56.80, 72.16, 72.22, 72.65, 72.70, 112.89, 114.99, 125.20, 127.25, 127.96, 128.01, 128.27, 128.38, 128.79, 133.20, 135.41, 137.41, 141.37, 144.87, 148.83, 148.93, 155.31; ³¹P NMR (200 MHz, CDCl₃): δ 20.67; Anal. Calcd for C₂₉H₃₁ClN₃O₄PS: C, 59.64; H, 5.35; N, 7.19; Found: C, 59.47; H, 5.17; N, 6.91.

O,O'-Diethyl-4-[[[2-phenyl-4-chloro-5-thio)pyridazin-3(2H)-one]](phenylamino)-(2-fluorophenyl)methylphosphonate (4d). Mp 206–208 °C; yield 66%; IR (ν cm⁻¹): 3298 (NH), 2978, 1668 (C=O), 1635, 1593, 1558, 1512, 1490, 1456, 1246 (P=O), 1026 (P-O-C); ¹H NMR (500 MHz, CDCl₃): δ 6.69–7.53 (m, 14H, Ar-H), 5.31–5.38 (m, 1H, NH), 5.15 (dd, J = 24 Hz, 8.6 Hz, 1H, CH), 3.68–4.26 (m, 4H, 2OCH₂), 1.33 (t, J = 7.15 Hz, 3H, CH₃), 1.08 (t, J = 6.85 Hz, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃): δ 16.16, 16.21, 16.51, 16.56, 47.31, 48.52, 63.58, 63.63, 63.80, 63.86, 113.50, 114.79, 115.44, 115.62, 122.77, 122.88, 124.92, 125.19, 127.35, 128.38, 128.79, 130.05,

130.07, 130.12, 130.13, 133.15, 137.52, 141.36, 144.74, 148.18, 148.30, 155.30; ^{31}P NMR (200 MHz, CDCl_3): δ 21.79 ($J = 4.6$ Hz); ^{19}F NMR (470 MHz, CDCl_3): δ -118.04; Anal. Calcd for $\text{C}_{27}\text{H}_{26}\text{ClFN}_3\text{O}_4\text{PS}$: C, 56.50; H, 4.57; N, 7.32; Found: C, 56.81; H, 4.21; N, 7.14.

***O,O'*-Dipropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(2-fluorophenyl)methylphosphonate (4e).** Mp 201–203 °C; yield 67%; IR (ν cm^{-1}): 3286 (NH), 2968, 1668 (C=O), 1595, 1560, 1512, 1490, 1456, 1228 (P=O), 1004 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.67–7.52 (m, 14H, Ar-H), 5.32–5.35 (m, 1H, NH), 5.15 (dd, $J = 24.6$ Hz, 8.55 Hz, 1H, CH), 3.55–4.11 (m, 4H, 2OCH_2), 1.41–1.72 (m, 4H, 2CH_2), 0.91 (t, $J = 7.45$ Hz, 3H, CH_3), 0.75 (t, $J = 7.45$ Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 9.84, 10.09, 23.70, 23.74, 23.95, 24.00, 47.17, 48.41, 69.04, 69.10, 113.38, 114.78, 115.41, 115.58, 122.91, 123.02, 124.90, 125.19, 127.32, 128.37, 128.79, 130.02, 130.07, 133.16, 137.50, 141.36, 144.78, 148.26, 148.38, 155.30, 159.82, 161.73; ^{31}P NMR (200 MHz, CDCl_3): δ 21.74 ($J = 4.6$ Hz); ^{19}F NMR (470 MHz, CDCl_3): δ -118.17; Anal. Calcd for $\text{C}_{29}\text{H}_{30}\text{ClFN}_3\text{O}_4\text{PS}$: C, 57.85; H, 5.02; N, 6.98; Found: C, 57.78; H, 5.02; N, 6.49.

***O,O'*-Diisopropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(2-fluorophenyl)methylphosphonate (4f).** Mp 228–230 °C; yield 59%; IR (ν cm^{-1}): 3284 (NH), 2978, 1668 (C=O), 1595, 1560, 1508, 1490, 1456, 1228 (P=O), 993 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.68–7.51 (m, 14H, Ar-H), 5.23–5.38 (m, 1H, NH), 5.07 (dd, $J = 24.6$ Hz, $J = 8.6$ Hz, H, CH), 4.77–4.80 (m, 1H, CH), 4.41–4.45 (m, 1H, CH), 1.34 (d, $J = 6.3$ Hz, 3H, CH_3), 1.29 (d, $J = 6.3$ Hz, 3H, CH_3), 1.23 (d, $J = 6.3$ Hz, 3H, CH_3), 0.85 (d, $J = 6.3$ Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 23.05, 23.10, 23.96, 24.00, 24.26, 47.57, 48.80, 72.38, 72.44, 72.76, 72.82, 113.18, 114.71, 115.33, 115.51, 123.23, 123.34, 124.80, 125.20, 127.29, 128.38, 128.79, 129.85, 129.90, 133.18, 137.50, 141.37, 144.81, 148.44, 148.55, 155.30, 159.89, 161.91; ^{31}P NMR (200 MHz, CDCl_3): δ 19.99 ($J = 4.58$ Hz); ^{19}F NMR (470 MHz, CDCl_3): δ -118.01; Anal. Calcd for $\text{C}_{29}\text{H}_{30}\text{ClFN}_3\text{O}_4\text{PS}$: C, 57.85; H, 5.02; N, 6.98; Found: C, 57.98; H, 5.24; N, 6.72.

***O,O'*-Diethyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(2-methoxyphenyl)methylphosphonate (4g).** Mp 205–207 °C; yield 79%; IR (ν cm^{-1}): 3280 (NH), 2980, 1664 (C=O), 1635, 1593, 1558, 1514, 1490, 1458, 1246 (P=O), 1026 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.67–7.53 (m, 14H, Ar-H), 5.36–5.42 (m, 2H, NCH+NH), 3.56–4.22 (m, 7H, $2\text{OCH}_2 + \text{OCH}_3$), 1.32 (t, $J = 7.45$ Hz, 3H, CH_3), 1.02 (t, $J = 7.45$ Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 16.19, 16.24, 16.55, 16.60, 47.08, 48.31, 55.87, 63.16, 63.21, 63.59, 63.64, 110.61, 112.53, 114.77, 121.25, 123.60, 125.19, 127.12, 128.16, 128.36, 128.78, 129.50, 133.25, 137.40, 141.36, 145.03, 148.69, 148.80, 155.32, 157.18, 157.23; ^{31}P NMR (200 MHz, CDCl_3): δ 23.39; Anal. Calcd for $\text{C}_{28}\text{H}_{29}\text{ClN}_3\text{O}_5\text{PS}$: C, 57.39; H, 4.99; N, 7.17; Found: C, 57.68; H, 5.01; N, 7.00.

***O,O'*-Dipropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(2-methoxyphenyl)methylphosphonate (4h).** Mp 193–195 °C; yield 72%; IR (ν cm^{-1}): 3296 (NH), 2966, 1670 (C=O), 1635, 1593, 1560, 1519, 1490, 1456, 1230 (P=O), 997 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.67–7.53 (m, 14H, Ar-H), 5.37–5.43 (m, 1H, NCH+NH), 3.43–4.10 (m, 7H, $2\text{OCH}_2 + \text{OCH}_3$), 1.36–1.73 (m, 4H, 2CH_2), 0.92 (t, $J = 7.45$ Hz, 3H, CH_3), 0.73 (t, $J = 7.45$ Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 9.88, 10.14, 23.69, 23.73, 23.96, 24.01, 47.00, 48.23, 55.81, 68.60, 68.66,

68.86, 68.92, 110.55, 112.49, 114.74, 121.24, 123.71, 125.19, 127.12, 128.20, 128.24, 128.36, 128.78, 129.45, 133.24, 137.40, 137.40, 141.37, 145.03, 148.70, 148.81, 155.32, 157.11, 157.16; ^{31}P NMR (200 MHz, CDCl_3): δ 23.35; Anal. Calcd for $\text{C}_{30}\text{H}_{33}\text{ClN}_3\text{O}_5\text{PS}$: C, 58.68; H, 5.42; N, 6.84; Found: C, 58.71; H, 5.22; N, 6.82.

***O,O'*-Diisopropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(2-methoxyphenyl)methylphosphonate (4i).** Mp 222–224 °C; yield 51%; IR (ν cm^{-1}): 3300 (NH), 2976, 1664 (C=O), 1595, 1560, 1519, 1490, 1458, 1228 (P=O), 1002 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.67–7.53(m, 14H, Ar-H), 5.29–5.35 (m, 1H, NCH+NH), 4.74–4.80 (m, 1H, OCH), 4.34–4.40 (m, 1H, OCH), 3.93 (s, 3H, OCH_3), 1.34 (d, J = 6.3 Hz, 3H, CH_3), 1.28 (d, J = 6.3 Hz, 3H, CH_3), 1.21 (d, J = 6.3 Hz, 6H, 2CH_3), 0.76 (d, J = 6.3 Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 22.92, 22.97, 23.99, 24.03, 24.25, 24.28, 24.39, 24.41, 47.39, 48.63, 55.78, 71.83, 71.88, 72.37, 72.43, 110.57, 112.36, 114.70, 121.19, 124.13, 125.20, 127.14, 128.17, 128.21, 128.34, 128.78, 129.28, 133.25, 137.39, 141.39, 145.04, 148.87, 148.97, 155.32, 157.28; ^{31}P NMR (200 MHz, CDCl_3): δ 21.78; Anal. Calcd for $\text{C}_{30}\text{H}_{33}\text{ClN}_3\text{O}_5\text{PS}$: C, 58.68; H, 5.42; N, 6.84; Found: C, 58.85; H, 5.12; N, 6.60.

***O,O'*-Diethyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(4-chlorophenyl)methylphosphonate (4j).** Mp 206–208 °C; yield 58%; IR (ν cm^{-1}): 3282 (NH), 2983, 1662 (C=O), 1593, 1521, 1489, 1234 (P=O), 1016 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.62–7.53 (m, 14H, Ar-H), 5.24–4.27 (m, 1H, NH), 4.70 (dd, J = 24 Hz, 6.85 Hz, 1H, NCH), 3.71–4.81 (m, 4H, 2OCH_2), 1.15 (t, J = 6.9 Hz, 3H, CH_3), 1.30 (t, J = 6.9 Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 16.32, 16.37, 16.52, 16.57, 54.61, 55.83, 63.57, 63.62, 63.80, 63.84, 113.56, 115.04, 125.18, 127.39, 128.41, 128.81, 129.09, 129.12, 129.16, 133.15, 133.72, 134.29, 137.48, 141.34, 144.70, 148.33, 148.45, 155.30; ^{31}P NMR (200 MHz, CDCl_3): δ 21.80; Anal. Calcd for $\text{C}_{27}\text{H}_{26}\text{Cl}_2\text{N}_3\text{O}_4\text{PS}$: C, 54.92; H, 4.44; N, 7.12; Found: C, 55.07; H, 4.44; N, 7.08.

***O,O'*-Dipropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(4-chlorophenyl phenyl)methylphosphonate (4k).** Mp 156–158 °C; yield 67%; IR (ν cm^{-1}): 3288 (NH), 2970, 1666 (C=O), 1593, 1516, 1489, 1232 (P=O), 999 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.63–7.53 (m, 14H, Ar-H), 5.20–4.39 (m, 1H, NH), 4.70 (dd, J = 24 Hz, 6.85 Hz, 1H, NCH), 3.60–4.08 (m, 4H, 2OCH_2), 1.49–1.56 (m, 4H, 2CH_2), 0.91 (t, J = 7.45 Hz, 3H, CH_3), 0.81 (t, J = 7.45 Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 9.96, 10.09, 23.80, 23.84, 23.95, 24.00, 54.57, 55.78, 68.98, 69.05, 69.12, 113.50, 115.02, 125.18, 127.37, 128.40, 128.80, 129.13, 133.14, 133.86, 134.25, 137.47, 141.35, 144.71, 148.39, 148.49, 155.29; ^{31}P NMR (200 MHz, CDCl_3): δ 21.70; Anal. Calcd for $\text{C}_{29}\text{H}_{30}\text{Cl}_2\text{N}_3\text{O}_4\text{PS}$: C, 56.31; H, 4.89; N, 6.79; Found: C, 56.39; H, 4.52; N, 6.49.

***O,O'*-Diisopropyl-4{[(2-phenyl-4-chloro-5-thio)pyridazin-3(2*H*)-one]}(phenylamino)-(4-chlorophenyl phenyl)methylphosphonate (4l).** Mp 198–200 °C; yield 63%; IR (ν cm^{-1}): 3267 (NH), 2978, 1664 (C=O), 1593, 1516, 1489, 1234 (P=O), 993 (P-O-C); ^1H NMR (500 MHz, CDCl_3): δ 6.62–7.53 (m, 14H, Ar-H), 5.21–5.22 (m, 1H, NH), 4.48–4.74 (m, 3H, NCH+2CH), 1.33 (d, J = 6.3 Hz, 3H, CH_3), 1.28 (d, J = 6.3 Hz, 3H, CH_3), 1.24 (d, J = 6.3 Hz, 3H, CH_3), 0.98 (d, J = 6.3 Hz, 3H, CH_3); ^{13}C NMR (125 MHz, CDCl_3): δ 23.40, 23.44, 23.93, 23.96, 24.26, 55.08, 56.29, 72.42, 72.48, 72.66, 72.72, 113.31, 114.98, 125.19, 127.34, 128.40, 128.80, 128.96, 129.25, 129.29, 133.16, 134.07, 134.15, 137.46, 141.35, 144.76, 148.59, 148.70, 155.30; ^{31}P NMR (200 MHz, CDCl_3): δ 19.85; Anal. Calcd for $\text{C}_{29}\text{H}_{30}\text{Cl}_2\text{N}_3\text{O}_4\text{PS}$: C, 56.31; H, 4.89; N, 6.79; Found: C, 56.18; H, 4.72; N, 6.67.

REFERENCES

1. Hoshino, K.; Koike, Y. JP Patent 2005263694 (2005); *Chem. Abstr.* **2007**, *143*, 320612147.
2. Wu, J.; Song, B. A.; Chen, H. J.; Bhadury, P. S.; Hu, D. Y. *Molecules* **2009**, *14*, 3676.
3. Zou, X. J.; Jin, G. Y.; Yang, Z. *Chin. J. Appl. Chem.* **2001**, *18*, 599.
4. Okujima, H.; Narimatsu, A.; Kobayashi, M.; Furuya, R.; Kitada, Y. *Jpn. Kokai Tokkyo JP* **63**, 215, 672; *Chem. Abstr.* **1989**, *110*, 75541.
5. Kafarski, P.; Lejczak, B.; Tyka, R.; Koba, L.; Pliszcak, E.; Wiecczorek, P. J. *Plant Growth Regul.* **1995**, *14*, 199.
6. Maier, L.; Diel, P. J. *Phosphorus, Sulfur Silicon Relat. Elem.* **1991**, *57*, 57.
7. Chen, R. Y.; Mao, L. J. *Phosphorus, Sulfur Silicon Relat. Elem.* **1994**, *89*, 97.
8. Shi, D. Q.; Chen, R. Y. *Chin. J. Appl. Chem.* **2002**, *19*, 780.
9. Jin, L. H.; Song, B. A.; Zhang, G. P.; Xu, R. Q.; Zhang, S. M.; Gao, X. W.; Hu, D. Y.; Yang, S. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 1537.
10. Deng, S. L.; Baglin, I.; Nour, M.; Flekhter, O.; Vita, C.; Cave, C. *Phosphorus, Sulfur Silicon Relat. Elem.* **2007**, *182*, 951.
11. Pan, W. D.; Ansiaux, C.; Vincent, S. P. *Tetrahedron Lett.* **2007**, *48*, 4353.
12. Manjula, A.; Vittal, B. R.; Neelakantan, P. *Synth. Commun.* **2003**, *33*, 2963.
13. (a) Vahdat, S. M.; Baharfar, R.; Tajbakhsh, M.; Heydari, A.; Baghbanian, S. M.; Khaksar, S. *Tetrahedron* **2008**, *49*, 6501; (b) Yang, S.; Gao, X. W.; Diao, C. L.; Song, B. A.; Jin, L. H.; Xu, G. F.; Zhang, G. P.; Wang, W.; Hu, D. Y.; Xue, W.; Zhou, X.; Lu, P. *Chin. J. Chem.* **2006**, *24*, 1581; (c) Reddy, Y. T.; Reddy, P. N.; Kumar, B. S.; Rajput, P.; Sreenivasulu, N.; Rajitha, B. *Phosphorus, Sulfur Silicon Relat. Elem.* **2007**, *182*, 161.
14. Chen, M. H.; Chen, Z.; Song, B. A.; Bhadury, P. S.; Yang, S.; Cai, X. J.; Hu, D. Y.; Xue, W.; Zheng, S. J. *J. Agric. Food. Chem.* **2009**, *57*, 1383.
15. Huang, R. Q.; Wang, H. L.; Zhou, J. *Preparation of Organic Intermediate*; Chemical Industry Press of China: Beijing, China, 2001; pp. 224–225.
16. Mowry, D. T. *J. Am. Chem. Soc.* **1953**, *75*, 1909.