

Effects of ACTH Peptide Fragments on Memory Formation

JAMES F. FLOOD AND MURRAY E. JARVIK

Department of Psychiatry, University of California, Los Angeles, CA 90024

AND

EDWARD L. BENNETT AND ANN E. ORME

Laboratory of Chemical Biodynamics, University of California, Berkeley, CA 94720

FLOOD, J. F., M. E. JARVIK, E. L. BENNETT AND A. E. ORME. *Effects of ACTH peptide fragments on memory formation*. PHARMAC. BIOCHEM. BEHAV. 5: SUPPL. 1, 41-51, 1976. The effects of peptides derived from ACTH on the formation of long-term memory have been investigated in male mice. Post-training administration of ACTH 4-10-L-Phe-7 (ACTH-L) improved retention for both passive and active avoidance tasks. Administration of ACTH 4-10-D-Phe-7 (ACTH-D) impaired retention for both tasks. The optimum dose for ACTH-L was about 0.3 mg/kg; the optimum dose for ACTH-D was in the range of 1.0-3.0 mg/kg. Using the passive avoidance task, it was shown that either drug had to be administered within 60 min of training to be highly effective. Amnesia produced by anisomycin (Ani), an inhibitor of protein synthesis, was lessened by ACTH-L and increased by ACTH-D. ACTH-D opposed the memory-facilitating effects of ACTH-L. Using intact mice, ACTH-L or ACTH-D did not significantly change the incorporation of valine into protein, nor did these peptides influence the inhibition of protein synthesis caused by anisomycin. The results show that ACTH may play a major role in memory processing, perhaps by facilitating essential protein synthesis at sites specific for the memory being established.

Active avoidance	Passive avoidance	Arousal	Anisomycin	Protein synthesis
Inhibition of protein synthesis	ACTH 4-10-L-Phe-7	ACTH 4-10-D-Phe-7	Peptides	

SINCE the early 1960's a great deal of research has been done on the role of the pituitary-adrenocortical axis in animal learning where footshock was the main motivation. More recently attention has turned to the effects of ACTH and related compounds on retrieval of stored information through studies of the effect of ACTH peptides on extinction of a habit.

De Wied and his co-workers have suggested that ACTH 4-10-L-Phe-7 (ACTH-L) is the portion of ACTH which affects behavior and that this portion has no adrenocorticotrophic activity. In addition ACTH 4-10-D-Phe-7 (ACTH-D) opposes the activity of ACTH-L [1] and increases the rate of extinction for shuttlebox training [1]. Greven and De Wied [19] studied different peptide fragments of ACTH to determine the shortest possible sequence of amino acids that would have the same effect on extinction as the full sequence of ACTH. ACTH-L was reported to be as effective as the naturally occurring ACTH in delaying extinction of shuttlebox performance, jump-pole response, and passive avoidance. ACTH-D facilitated shuttlebox and jump-pole extinction but delayed extinction for passive avoidance. The ACTH 4-10 peptides possess almost none of the adrenocorticotrophic effects of the parent molecule ACTH [3]. Consistent with the above findings it has been reported [24] that ACTH-L administered prior to the retention test could alleviate the amnesia caused by CO₂ or electroconvulsive shock.

Greven and de Wied [19] suggested that the ACTH

peptide acts on membranes of specific target cells and by inducing conformational changes, stimulates cyclic AMP production. This in turn stimulates protein synthesis and leads to the establishment of new synaptic junctions. Gispen *et al.* [16] and Walter [30] have outlined rather similar postulated mechanisms for actions of ACTH peptides on receptors leading to membrane changes through the cyclic nucleotide system. Greengard [18] has recently discussed the multiple roles that cyclic nucleotides may play in synaptic function.

The research that we have undertaken focuses on the effects of the ACTH peptides on memory formation as opposed to the experimental designs used to study its effects on learning and retrieval. For this reason, we have investigated the effects of ACTH 4-10-L-Phe-7 and ACTH 4-10-D-Phe-7 either alone or in combination with anisomycin, a protein synthesis inhibitor, on retention and protein synthesis. A number of studies have now shown that anisomycin is an effective inhibitor of memory formation [9, 10, 12-14, 27, 28]. In all of the experiments in this report, the ACTH peptide fragments were injected within 4 hr after training and thus 1 week prior to the retention test. This makes it unlikely that the effects of the peptides on retention are due to influences on the acquisition of the habit or on retrieval. Both active and passive avoidance tasks were used.

The effects of ACTH and derivative peptides have been studied in many laboratories under a variety of conditions.

Macromolecular effects of ACTH and its analogues in the brain including effects on protein synthesis have recently been reviewed or reported [4, 7, 8, 21]. Most of the studies cited investigated the effects of either ACTH or its analogues using either hypophysectomized or adrenalectomized mice or rats, or have used in vitro systems. However, no study was noted in which the compounds, mode of injection, and time parameters appeared to be relevant for our behavioral studies. The main objectives of our biochemical experiments were twofold: (1) to determine any gross effects of ACTH 4-10 peptides on cerebral protein synthesis within hours after a single subcutaneous administration, and (2) to determine if some unanticipated interaction occurred between anisomycin and either peptide to greatly modify the inhibition of protein synthesis.

MATERIALS AND PROCEDURES

BEHAVIORAL EXPERIMENTS

For the behavioral experiments, CD-1 male mice from Charles Rivers Breeding Laboratories, Wilmington, MA were obtained at 6 weeks of age. After 1 week in the laboratory, the mice were individually housed in small cages 24 hr prior to training. After training the mice were returned to individual cages until the retention test was given one week later. The mice were trained on a one-trial step-through passive avoidance or on a T-maze active avoidance task. Mice were tested and trained between 0800 and 1400 hr.

Passive Avoidance

The passive avoidance training and apparatus have been described previously [11]. The apparatus consists of a 44 cm long alley divided into a small, black start box and a longer white shock compartment. The two compartments are separated by a panel which contains a mouse hole. Entry into the white compartment was prevented until the appropriate time by a translucent guillotine door. The shock was administered by a high voltage, constant current 18 pole shock scrambler through a brass floor grid in the white box. The footshock intensity is given in each experimental description.

The training trial consisted of placing a mouse in the black start box for 20 sec, then illuminating the white shock box and the mouse hole for an additional 20 sec. Next, the guillotine door was removed while the mouse was facing away from it. The latency-to-enter was determined from the time the mouse oriented toward the mouse hole until it had entered completely the white compartment. The shock was turned on when the mouse was half-way down the alley (about 5 sec after entry), and was left on until the mouse escaped from the shock box and was returned to its own cage.

The retention test followed the same procedure as for training except that no footshock was given. Mice not entering into the white compartment within 180 sec were removed and given a score of 180 sec. Amnesia was defined as entering into the white shock compartment in 20 sec or less. Retention was defined as not entering the shock compartment within 20 sec. The time from shock onset until the animal returned to the black start box was called the latency-to-escape.

T-Maze Active Avoidance

The T-maze training and apparatus have been previously

described [9]. The training apparatus consisted of a black Plexiglas start alley with a start box at one end and two goal boxes at the other. A brass floor grid ran throughout the entire maze. Each goal box was fitted with a clear Plexiglas liner, the bottom of which went below the shock grid. This liner was used to remove the animal from the goal box. The start box was separated from the rest of the start alley by a black Plexiglas guillotine door which prevented the animal from moving down the start alley until the trial started. Animals were not permitted to explore the maze prior to training. The conditioned stimulus (CS) was a door bell type buzzer. The intertrial interval was about 40 sec; 0.37 mA shock was used in all active avoidance tasks.

A training trial consisted of placing the mouse in the start box, then raising the guillotine door and simultaneously sounding the buzzer. Mice not moving to the correct goal box within 5 sec were shocked until they did so. The side preference was determined on the first training trial by forcing all mice to go to the side opposite to their first response. On subsequent trials, the correct goal box was the non-preferred side for each mouse. At the end of each trial, the mouse was removed to its home cage by carefully removing the liner and placing it into the mouse cage. As training proceeded a mouse could make one of two types of responses: (a) a response latency longer than 5 sec was an escape, (b) a response latency less than or equal to 5 sec was an avoidance. The retention test given 1 week after training consisted of retraining to a criterion of 1 avoidance response. Mice requiring more than 3 trials to make the first avoidance response were scored as amnesic.

Injections

When pretraining injections of saline or anisomycin (2-p-methoxyphenyl-3-acetoxy-4-hydroxy-pyrrolidine) (20 mg/kg) were employed, they were administered under very light ether anesthesia. All ACTH peptide or control injections were given without ether. All injections were given subcutaneously in a volume of about 0.35 ml. Solutions of the ACTH fragments used for the behavioral studies were prepared in saline after first dissolving in a small quantity of 0.01 M HCl.

BIOCHEMICAL EXPERIMENTS

The effects of the ACTH peptides, either alone or in combination with anisomycin, on protein synthesis was determined by measuring the incorporation of [^{14}C (U)]-L-valine into the trichloroacetic acid fraction of either whole brain in the first experiment, or into brain stem and rest of brain separately in the second. For the first experiment, mice were sacrificed 75, 100, and 135 min after subcutaneous administration of 0.5 mg of anisomycin. ACTH-D, ACTH-L, or saline was administered 45 min after the anisomycin as had been done in the behavioral experiments. [^{14}C]-L-valine was administered 15 min prior to sacrifice. The design of Experiment 2 was similar except mice were sacrificed 135 or 180 min after anisomycin injection. In the biochemical experiments (but not in the behavioral experiments), mannitol had been used to stabilize the peptide solutions. Therefore, in addition to the usual saline controls, some mice were administered mannitol. Experimental procedures for determining protein synthesis have been described in further detail [11]. Duplicate fractionations and determinations of radioactivity were determined for

each brain sample. Four intact male Swiss mice, which were offspring of mice obtained from Charles River Breeding Laboratories, were used for each data point in the first experiment and 2 mice per data point in the second experiment.

BEHAVIORAL EXPERIMENTS AND RESULTS

EFFECTS OF ACTH 4-10-L-Phe-7 ON RETENTION EXPERIMENTS 1-3

Experiment 1: Dose Response

The purpose of the experiment was to determine the dose range that might facilitate retention. ACTH 4-10-L-Phe-7 (ACTH-L) was administered 30 min after passive avoidance training. A low shock level of 0.30 mA was chosen so that controls would show poor performance and thus improvement of retention could be measured. As a further means of setting the proper level of training, we retained only those mice with latencies-to-enter by latencies-to-escape of $2 \text{ sec} \times 1 \text{ sec}$, 2×2 , 2×3 , 3×1 , or 3×2 (values rounded to the nearest second). These latency values are very important since they determine the degree of learning [11]. ACTH-L was administered at 0, 0.1, 0.3, 1.0 and 3.0 mg/kg. The 0.1 and 0.3 mg/kg doses are values similar to those used by de Wied and his associates.

Results

The results summarized in Fig. 1 show that for this passive avoidance task, the 0.1, 0.3 and 1.0 mg/kg doses of ACTH-L facilitated retention significantly and equally, that is, these groups had all showed about 80% retention, whereas the group receiving no ACTH-L showed only 27% (Fig. 1). There is some indication that too much of the peptide produces less than optimal facilitation of memory formation since the 3.0 mg/kg showed 50% retention.

Experiment 2: Interaction of Dose and Time of Administration

The purpose of this experiment was to determine if (a) ACTH-L could produce a better dose response curve with a decrease in training strength from that used in Experiment 1, and (b) if the effect of ACTH-L was dependent on the time of drug administration after training. The mice were trained (shock intensity 0.30 mA) and injected in the same manner as in Experiment 1 except that the latency-to-enter had to be 2 sec and the latencies-to-escape 1 or 2 sec. This shifted the distribution of training values toward those producing somewhat poorer retention; that is the controls were amnesic. Thus the test should be more sensitive to the effects of ACTH-L. The N per group was 20.

Results

The dose response portion of the ACTH-L curve over which retention improved was between 0.03 and 0.1 mg/kg. The dose of 0.3 mg/kg seemed to produce the optimal level of improvement in retention (Table 1). However, the effective dose range was still 0.1 to 1.0 mg/kg. The effect of ACTH-L on retention was dependent on the time of its administration. Administration of the peptide at 30 or 60 min after training had similar effects on retention. But ACTH-L was less effective in improving retention when administered 90 min after training and an additional loss of effectiveness was observed when administration occurred

EFFECTS OF DOSE OF ACTH-L ON RETENTION PASSIVE AVOIDANCE TRAINING




FIG. 1. The effects of dose of ACTH-L on retention for passive avoidance training. The ACTH was administered 30 min after training. Retention was tested one week later. N/group was between 23 and 28.

240 min after training (Table 1). The decrease in effectiveness of the higher ACTH-L doses was found again. In addition, this modified training procedure was an improvement over the procedure previously used since it made it possible to demonstrate that a dose of 0.3 mg/kg was optimal.

Experiment 3: ACTH-L on Retention using T-Maze Active Avoidance

The purpose of this experiment was to test if ACTH-L would have the same effect on memory formation for an active avoidance task as it did in the passive avoidance task. The mice were given 4 training trials on the T-maze using a muffled door bell buzzer as CS. ACTH-L was administered within 1 min after training at doses of 0.3 or 3.0 mg/kg. A saline-injected group served as control. The N per group was 10.

Results

Administration of ACTH-L immediately after active avoidance training improved retention scores both as measured by the percentage of mice requiring more than 3 trials to make the first avoidance response i.e., amnesic mice (Fig. 2B), and by the number of trials for the first avoidance (Fig. 2A). Again as in the passive avoidance task, the higher dose was less effective than the lower dose of ACTH-L.

TABLE 1
AMNESIA AS A FUNCTION OF TIME AND DOSE OF ACTH-L
PASSIVE AVOIDANCE TASK

Time of Injection after training (min)	Dosage in mg/kg					
	0.0	0.03	0.1	0.3	1.0	3.0
	% Amnesic Mice*					
30	80	60	30	15	35	55
60	80	60	25	20	50	60
90	85	75	50	40	50	75
240	80	80	75	65	75	75

*N = 20/group.

EFFECTS OF ACTH 4-10-D-Phe-7 ON RETENTION
EXPERIMENTS 4-5

Experiment 4: Interaction of Dose and Time of Administration of ACTH-D

The purpose of the experiment was twofold: (1) to determine what doses of ACTH 4-10-D-Phe-7 (ACTH-D) would affect retention test scores and (2) to determine the influence of time of administration on the effect. The mice were trained on one-trial passive avoidance. In Experiment 2 it was necessary that the control animals would be only minimally trained so that facilitation of memory could be

demonstrated. In this experiment control animals need to be well trained so that impairment of memory formation by ACTH-D could be shown. One way of increasing training strength is to raise the level of shock intensity. For this reason shock intensity was 0.33 mA rather than 0.30 mA as in Experiment 2. ACTH-D was administered 30, 60, 90, or 240 min after training at doses of 0.3, 1.0 or 3.0 mg/kg. Saline was administered as a control for the stress of the injection. We retained only animals with latencies-to-enter of 2 sec and latencies-to-escape of 2 sec (when rounded off to the nearest second). The N per group was 20.




FIG. 2. Effect of dose of ACTH-L on active avoidance training. ACTH-L was administered immediately after training. A, (left) Mean number of trials to first avoidance; B, (right) Percentage of mice requiring more than 3 trials to make first active avoidance response.

Results

All three doses of ACTH-D were effective at reducing retention scores and thus increasing amnesia when administered 30 or 60 min after training. There is a trend under these conditions of training for the high dose of ACTH-D to be more effective than the low dose (Table 2). As was found with ACTH-L in Experiment 2, ACTH-D exerted a greater effect when injected 30 or 60 min than when injected 90 or 240 min after training.

TABLE 2

AMNESIA AS A FUNCTION OF TIME AND DOSE OF ACTH-D
PASSIVE AVOIDANCE TASK

Time of Injection after training (min)	Dosage in mg/kg			
	0.0	0.3	1.0	3.0
		% Amnesic Mice*		
30	20	55	65	75
60	15	55	70	70
90	15	30	35	40
240	20	10	30	45

*N = 20/group.

Experiment 5: ACTH-D Effect on T-Maze Active Avoidance

The purpose of the experiment was to determine if ACTH-D would have the same effect on retention for an active avoidance task as it did in a passive avoidance task. In Experiment 3 we examined the facilitating effect of ACTH-L on an active avoidance task, therefore it was necessary for controls to be minimally trained so only 4 training trials were given. In this experiment we are looking for impairment of memory, therefore well-trained controls were needed. For this reason the mice were given 5 training trials with a loud door bell buzzer as CS. In order to control the degree of learning a training performance criterion was imposed. Animals were not included if they made an avoidance response during original training or if they failed to make one correct escape response. ACTH-D was administered within 1 min after training at doses of 0.0, 0.3, 1.0 or 3.0 mg/kg. The retention test, as in all experiments of this series, was one week after training. The N was 20 per group.

Results

ACTH-D at 1.0 or 3.0 mg/kg significantly impaired retention; that is the percentage of amnesic mice increased (Table 3). As with Experiment 4, a significant difference was not found between 1.0 and 3.0 mg/kg doses although there is indication of greater effect with the larger dose (compare Tables 2 and 3).

The mice which made avoidance responses during the original training were analyzed separately to see if ACTH-D would still have an effect (Table 4). Interestingly, this seems to have brought out the differences in dose response. ACTH-D at 3.0 mg/kg caused more amnesia. Owing to the small N's no differences were significant but the group given 3.0 mg/kg differed from the control group at approximately $p < 0.06$.

TABLE 3

AMNESIA AS A FUNCTION OF DOSE OF ACTH-D
ACTIVE AVOIDANCE TASK

Dose in mg/kg	0.0	0.3	1.0	3.0
% amnesic mice*	20%	40%	60%	65%
p value from 0.0		NS	0.05	0.02

*N = 20/group. Mice not making the first avoidance response in 3 trials or less were scored as amnesic.

TABLE 4

AMNESIA AS A FUNCTION OF DOSE OF ACTH-D
SELECTED ANIMALS*—ACTIVE AVOIDANCE TASK

Dose in mg/kg	0.0	0.3	1.0	3.0
N	10	8	7	8
% Amnesia	10%	12%	14%	50%

*Only those animals which made an avoidance during the original training were used. Amnesic mice defined as in Table 3.

EFFECTS OF ACTH ON AMNESIA PRODUCED BY ANISOMYCIN: EXPERIMENTS 6-9

Experiment 6: Effects of ACTH-L

Protein synthesis inhibition by anisomycin (Ani) was used to cause amnesia in this experiment. In some of the mice ACTH-L was administered shortly after training (30 min) to determine if the ACTH-L would block the amnesia. Other groups received more injections of Ani to test if longer durations of inhibition of protein synthesis would block the anti-amnesic effect of ACTH-L.

Mice were trained on passive avoidance at a footshock of 0.36 mA. This is a relatively high footshock and produces strong training. Ani was administered 15 min prior to training and again at 1-3/4 hr after training. Some groups received a third injection of Ani 3-3/4 hr after training. Saline-injected mice served as controls. Saline, 0.3 or 3.0 mg/kg of ACTH-L was administered 30 min after training. N per group was 20.

Results

As expected, two or three successive injections of Ani caused significantly greater amnesia than was found in the saline control (80 and 85% vs 15%). An injection of ACTH-L at the dose of 3.0 mg/kg significantly reduced this amnesia in mice given 2 successive injections of Ani but not in those given 3 successive injections of Ani (Fig. 3). The effect of 0.3 mg/kg ACTH-L, while significant, did not cause such a large drop in the percent amnesia. ACTH-L did block the amnesic effect of two successive injections of Ani but increasing the number of injections (and thus the duration of inhibition) blocked the anti-amnesic effects of the ACTH-L.

The high dose of ACTH was more effective at reducing the level of amnesia than the lower dose. However, an additional injection of Ani which prolongs inhibition of protein synthesis by 2 hr at 80% or more blocked the anti-amnesic effect of the ACTH-L.




FIG. 3. Effects of Anisomycin (A) and Dose of ACTH-L on retention for a passive avoidance task. A was administered 15 min prior to training and subsequently at 2 hr intervals as indicated. ACTH or saline (S) was administered 30 min after training. ACTH-L overcame the amnesic effects of 2 doses of A, but not of 3 doses. The 3.0 mg/kg dose of ACTH was more effective than 0.3 mg/kg.

Experiment 7: Effects of Number of Injections of ACTH-L

The purpose of this experiment was to determine if additional injections of ACTH-L would block the effect of additional injections of Ani as observed in Experiment 6. The conditions of training were the same as in Experiment 6. The injection groups are given in Fig. 4. Ani or saline were administered 15 min prior to training and 1-3/4 hr after training. Some groups received a 3rd injection 3-3/4 hr after training. ACTH-L or its control injection (saline) was administered 1/2 hr or 1/2 h and 2-1/2 hr after training. ACTH-L was administered at a dose of 3.0 mg/kg. N per group was 10.

As in experiment 6, two injections of Ani caused significantly more amnesia than did injections of saline. ACTH-L blocked the amnesia in groups receiving only two successive injections of Ani. Neither a single or double injection of ACTH-L blocked the amnesia induced by three successive injections of Ani (Fig. 4). These data, like the time dependent results of Experiment 2, suggest that ACTH-L alters memory formation only when administered within the first hour after training.

Experiment 8: Effects of ACTH-D

Under appropriately high conditions of training the amnesic effect of high levels of protein synthesis inhibition




FIG. 4. Effect of number of injections of ACTH-L on amnesia produced by anisomycin: Anisomycin (A) was administered 15 min prior to training on a passive avoidance task and subsequently at 2 hr intervals. ACTH-L or saline (S) was administered 30 min after training and subsequently at 2 hr intervals as shown. ACTH-L did not overcome the amnesia produced by 6 hr of high inhibition of protein synthesis from 3 injections of anisomycin.

by anisomycin can be reduced so that one can test if another substance will increase amnesia.

Animals were trained on the passive avoidance task. They received two injections of either Ani or saline 15 min prior to training and 1-3/4 hr after training. ACTH-D was administered 30 min after training at 0.0, 0.3, 0.1 or 3.0 mg/kg. The latency-to-enter and escape was 2 sec by 2 sec. The footshock intensity was 0.34 mA. The N per group was 20.

Results

Two successive injections of Ani alone did not cause




FIG. 5. Effect of ACTH-D and anisomycin on amnesia. With strong training, for a passive avoidance task, anisomycin administered 15 min prior and 1- $\frac{1}{4}$ hr after training caused only a low percentage of amnesic mice. ACTH-D increased the amnesia; high doses were more effective than low doses.

significant amnesia. The groups receiving ACTH-D all differ significantly from the group receiving only [A(ACTH-D 0)A] (Fig. 5). The 3.0 and 1.0 mg/kg doses differ at $p < 0.001$. The groups receiving ACTH-D did not differ significantly from each other but a strong indication of the weaker effect of the low ACTH-D dose versus the high is present. The doses 1.0 and 3.0 mg/kg in combination with Ani induced the greatest retention deficit or amnesia.

OPPOSING EFFECTS OF ACTH 4-10-L- AND-D-Phe-7 ON RETENTION

Experiment 9A

The results of Experiments 1 through 8 indicated that

ACTH-L facilitated memory formation while ACTH-D impaired memory formation. As a further test of the opposing effects of these peptide fragments, we administered ACTH-L at its optimal dose (0.3 mg/kg) and combined this with various doses of ACTH-D (0.3 to 6.0 mg/kg) to determine how much ACTH-D is required to block the facilitating effects of ACTH-L.

The mice were trained on passive avoidance at a footshock intensity of 0.30 mA. Animals had to have latencies-to-enter of 2 sec and latencies-to-escape of 2 sec to be retained. The ACTH-L, ACTH-L + ACTH-D (combined in one injection), or saline were administered 30 min after training. The doses of the peptides and the N's per group are given in Fig. 6A.

Results

ACTH-L given alone significantly enhanced retention when compared with the saline controls (Fig. 6A). A dose of ACTH-D at 3.0 or 6.0 mg/kg was sufficient to block the effect of ACTH-L. It appeared that 6.0 mg/kg of ACTH-D completely blocked the effect of ACTH-L (0.3 mg/kg).

Experiment 9B

As a further check on whether such a high dose of ACTH-D was required to block the ACTH-L effect on retention, Experiment 9A was repeated except that the footshock was reduced slightly to 0.28 mA. This was done to increase the percentage of amnesic mice in the control group. The groups used are shown in Fig. 6B. Other conditions were as in experiment 9A.

Results

ACTH-L again enhanced retention. In fact the percent difference was similar under the two training conditions (a difference of 45% for Experiment 9A, 58% for Experiment 9B: ACTH-L versus Saline). Under less vigorous training only the 6.0 mg/kg dose of ACTH-D completely blocked the effect of ACTH-L (Fig. 6B).

Experiment 9C

Experiments 9A and 9B showed that large doses of ACTH-D blocked the retention-enhancing effect of the ACTH-L. The results were interpreted to indicate that ACTH-L and ACTH-D have the same site of action for effecting memory formation but have opposing action. An alternate suggestion made at this Conference was that the apparent opposing effects of ACTH-D and ACTH-L were due to the increasing total dose of ACTH. It may be

TABLE 5

EFFECT OF ACTH-L, ACTH-D, AND ANISOMYCIN ON [14 C]-VALINE INCORPORATION IN WHOLE BRAIN
% INHIBITION OF VALINE INCORPORATION

	75		100		135	
	Total ACTH (min)		Total ACTH (min)		Total ACTH (min)	
	15		15		15	
	Sal	Ani	Sal	Ani	Sal	Ani
ACTH none	0	90	0	85	0	73
ACTH-L 2.0 mg/kg	5	90	13	85	10	76
ACTH-D 0.5 mg/kg	7	90	6	88	-3	69

OPPOSING EFFECTS OF ACTH-L AND ACTH-D PASSIVE AVOIDANCE TRAINING




FIG. 6. Interaction of ACTH-D and ACTH-L on retention for a passive avoidance task. Under either condition of strong training (high footshock) (left) or weak training (right), ACTH-D blocked the enhanced retention produced by ACTH-L. Drugs were administered 18 sec after training. N's/group and significance of difference from the control groups are shown above the data bars.

recalled that in Experiment 1, higher dosages of ACTH-L were less effective than lower doses. The purpose of this experiment was to determine if increasing the total amount of ACTH necessarily resulted in increased or sustained high levels of amnesia. The mice were well-trained on passive avoidance using 0.33 mA as in Experiment 4. The experimental groups of mice are given in Table 7. The ACTH or saline was administered 30 min after passive avoidance

training. Testing was one week after training. The N per group was 10.

Results

The results show that simply increasing the total amount of ACTH did not result in amnesia. A high percentage of amnesia resulted only when ACTH-D was administered at

TABLE 6
EFFECT OF ACTH-L, ACTH-D AND ANISOMYCIN ON [¹⁴C]-VALINE INCORPORATION
% INHIBITION OF VALINE INCORPORATION

Sal or Ani time (min)	Rest of Brain				Brain Stem			
	135		180		135		180	
	90		135		90		135	
Valine Incorporation (min)	15		15		15		15	
	Sal	Ani	Sal	Ani	Sal	Ani	Sal	Ani
ACTH added								
none	0	68	0	53	0	53	0	38
ACTH-L 0.5 mg/kg	13	71	-13	61	2	64	-1	43
2.0 mg/kg	-1	68			-10	58		
ACTH-D 0.5 mg/kg	-11	76	4	63	-20	61	8	44
Mannitol 333.3 mg/kg	-11		12	63	5			49

(-) denotes stimulation.

TABLE 7

AMNESIA AS A FUNCTION OF DOSE OF ACTH-D AND ACTH-L
PASSIVE AVOIDANCE TASK

ACTH-D (mg/kg)	ACTH-L (mg/kg)	% Amnesic Mice*
9.0	0	80%
7.0	0	90%
6.0	0	80%
6.0	1.0	50%
6.0	3.0	30%
0	0	30%

*N = 10/ group.

dosages of 6.0 to 9.0 mg/kg and in the absence of ACTH-L. When ACTH-L was added to the ACTH-D, amnesia was reduced or blocked (Table 7). For example, those mice receiving ACTH-D, 6.0 mg/kg plus ACTH-L, 3.0 mg/kg received as much total peptide as mice administered ACTH-D, 9.0 mg/kg. Combining the peptides resulted in only 30% amnesia, while 9.0 mg/kg ACTH-D alone yielded 80% amnesia. When compared to the group given 7.0 mg/kg of ACTH-D, the group given 6.0 mg/kg of ACTH-D plus 1 mg/kg of ACTH-L also showed decreased amnesia (90% vs 50%). The results show that with respect to memory formation, the action of ACTH-L and ACTH-D are opposing rather than additive. In addition, it is likely that they have the same site of action.

BIOCHEMICAL RESULTS

The results of Biochemical Experiments 1 and 2 indicate that ACTH-L or ACTH-D, when administered subcutaneously to intact mice, did not alter cerebral protein synthesis by more than 10% within 2-1/4 hr (Tables 5 and 6). Furthermore, these peptides did not significantly modify the inhibition caused by anisomycin. Mannitol, which had been used to solubilize the ACTH peptides for the biochemical experiments, by itself caused a slight increase of valine incorporation. It should be emphasized that mannitol was not used in the behavioral experiments since it introduced side effects there also.

DISCUSSION

ACTH Effects on Memory Formation

The finding of primary importance in these experiments was that ACTH-L, a peptide fragment of ACTH, administered shortly after training can influence subsequent retention test performance and thus aid memory formation (Experiments 1-3). ACTH-D, containing the unnatural isomer of phenylalanine, impairs memory formation (Experiments 4-5). These effects do not appear to be proactive since it was found that the effect of the ACTH peptides was time dependent. Such peptides can have an effect on retention test performance when administered shortly prior to testing [2, 5, 6, 19, 29] and for the most part these effects are consistent with those found in our studies.

The results confirm previous reports that ACTH 4-10-L-Phe-7 and ACTH 4-10-D-Phe-7 have opposite effects on retention for an active avoidance task [1, 19, 29] and for an appetitive task [15]. Gray and Garrud [17] have reported

that ACTH-L and ACTH-D given during acquisition had opposite effects during the non-rewarded extinction test: the L-peptide blocked the effect of the non-reward, the D-peptide accelerated the extinction. ACTH-L consistently enhanced retention in active and passive avoidance tasks (Experiments 1, 2, 3). ACTH-D impaired retention in active and passive avoidance (Experiments 4, 5). However, Greven and de Wied [19] reported that ACTH 1-10-L-Phe-7 did not show the anticipated reversal of the effects of ACTH 1-10-L-Phe-7 for a passive avoidance task. Consistent with the opposing actions of the two peptides was the finding that ACTH-L acted as an anti-amnesic agent (Experiment 6) and ACTH-D potentiated amnesia caused by brain protein synthesis inhibition (Experiment 8).

Effective Dose Range of the ACTH Peptides

The effective dose range of the peptides was similar to that reported for extinction test studies [19, 24, 29]. ACTH-L was effective over a dose range of 0.1 to 1.0 mg/kg or 3 to 30 μ g per 35 gm mouse. The optimal dose was 0.3 mg/kg or 10 μ g per mouse. The ACTH-D effective dose range was somewhat higher than reported for extinction studies. Doses from 0.3 to 3.0 mg/kg were effective and 1.0 mg/kg or 30 μ g/mouse seemed to be the most effective dose. Extinction studies have shown that the effective dose range of the L- and D-ACTH peptide fragments is about the same. However, when administered different ratios of the peptides after training, ACTH-D seemed to have significantly less affinity for the receptor site than the L form, since it took a ratio of ACTH-D to ACTH-L of at least 10 to 1 to block ACTH-L facilitation (Experiment 9A and 9B). It is also possible different receptors are involved with the ACTH-D receptor being less active. A third possibility would be differences in effective concentration of the two drugs at the active sites. De Wied has presented evidence that different receptor sites exist for the two forms of the peptide [6].

Anti-amnesic Studies

We have shown that anisomycin administered at 2 hr intervals will maintain a high level of inhibition of protein synthesis [12]. There is an interaction between the strength of training as determined by several factors including the shock intensity, duration of shock, etc., and the duration of inhibition required to produce amnesia [11-13]. We have recently shown that a number of stimulants and depressants, which act by a variety of mechanisms, can modify the amnesia produced by anisomycin (Flood *et al.*, submitted to *Behavioral Biology*). In the present series of experiments, it was found that ACTH-L and ACTH-D can also modify anisomycin-induced amnesia. ACTH 4-10-L-Phe-7 was found to block anisomycin-induced amnesia when the peptide was injected 30 min after training. However, extending the duration of inhibition by an additional 2 hr by a 3rd injection of Ani re-established the amnesia (Experiment 6). An additional injection of the peptide was not able to block amnesia when three successive anisomycin injections were used (Experiment 7).

ACTH-D increased a partial amnesia caused by anisomycin after strong training. We would like to caution against an interpretation that the anti-amnesic effect of ACTH-L suggests that anisomycin causes amnesia by interfering with ACTH function or vice versa. In this series of experiments, we were unable to show that ACTH-L or

ACTH-D had more than a slight generalized effect on protein synthesis as measured in large brain areas. In addition, neither ACTH-L nor ACTH-D appeared to change significantly the overall inhibition of protein synthesis produced by anisomycin. In interpreting the effects of ACTH peptides on memory other results that we have obtained should also be considered: (a) ACTH-L improves retention in poorly trained mice; (b) stimulants including d-amphetamine, strychnine, picrotoxin, caffeine, nicotine, and bicuculline all blocked amnesia induced by anisomycin and the amnesia could be regained by extending the duration of inhibition of protein synthesis (Flood *et al.*, *ibid.*, and unpublished observations); (c) dexamethasone and hydrocortisone could block amnesia induced by inhibition of protein synthesis; (d) depressants, including sodium phenobarbital, chloral hydrate, chlorpromazine, meprobamate, potentiated anisomycin-induced amnesia (Flood *et al.*, *ibid.*).

Previous work with the peptides and the question of permanence of amnesia has employed pretesting injections of ACTH-L to alleviate amnesia [23,24]. We have attempted to obtain similar effects using anisomycin to impair retention, but as yet we have not yet been successful in overcoming amnesia. A possible reason for the difference in results is that many studies compare trained treated and untreated animals. This comparison, while showing a loss of retention due to some amnesic treatment, does not necessarily imply that the loss of memory was complete or even substantial. In our studies, treated and untreated mice were compared against a performance criterion that represents the performance of naive mice. Treatments that can alleviate amnesia using pretesting injections may only be successful if the degree of memory loss is substantially less than complete. We recently reported that d-amphetamine reversed amnesia in anisomycin-injected mice which had long amnesic latencies in a passive avoidance task but had no effect on mice that originally had very short test latencies [14]. Presumably the amnesia was more severe in the case of those mice with very short test latencies and increased arousal was not an aid to recall. In addition, amphetamine had the same effect of improving recall in poorly trained saline-injected mice. Thus anti-amnesic effects are not necessarily restricted to animals receiving amnesic treatments. It might be said that anti-amnesic treatments have the ability to improve the recall of stored memories that are poorly stored either because of drug interference with consolidation or because of weak training as in the case of poorly trained undrugged animals.

Biochemical Effects of ACTH Peptides

While it is generally stated that ACTH and selected analogues stimulate protein synthesis, the clearest evidence for this stimulation comes from studies of hypophysectomized animals. In this system, either single or chronic injections have been shown to produce large increases in protein synthesis [20, 25, 26]. It should be noted that protein synthesis is depressed in these animals, and the effect of the peptides has been to restore protein synthesis to a more normal value. Another commonly used system to investigate the effects of ACTH analogues has been tissue slices of brain derived from both normal and operated animals. Here also stimulation of protein synthesis to a normal value was observed in the operated animal by *in vitro* addition of ACTH [22]. A surprising paucity of data appears to exist on the acute effects of ACTH and its analogues on cerebral protein synthesis. Rees *et al.* [21] have recently shown that ACTH peptides administered intracerebrally causes a slight stimulation of cerebral protein synthesis within 90 min after administration. Dunn *et al.* [8] have reported that a small but significant increased incorporation of lysine into brain protein within 15-25 min after administration of ACTH-L. Our studies are in general agreement with those of Rees *et al.* [21] in that any effects we found were small. It is clear that further research needs to be done to determine the effects of ACTH derivatives on protein synthesis at specific sites (micro effects).

In our experiments ACTH peptides had the ability to modify memory processing. ACTH-L facilitated memory processing and ACTH-D impaired processing. The effects were time-dependent and were observed in both passive and active avoidance tests. However, until more is known of the mechanics of action of these ACTH peptides, their role in memory processing will be uncertain.

ACKNOWLEDGEMENTS

We wish to express our appreciation to N. Belcher of Pfizer Pharmaceuticals for their generous gift of anisomycin, to Dr. H. Van Riezen of Organon International, Science development group, Holland for all ACTH peptides, and to Gary E. Smith and Daniel Vidal for skilled assistance for the behavioral experiments. The behavioral research was supported by NIMH grant NH 26608-01 to M.E. Jarvik, M.D., and the biochemical research by the U. S. Energy Research and Development Administration through the Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory.

REFERENCES

1. Bohus, B. and D. de Wied. Inhibitory and facilitatory effect of two related peptides on extinction of avoidance behavior. *Science* 153: 318-320, 1966.
2. Bohus, B., W. H. Gispen and D. de Wied. Effect of lysine vasopressin and ACTH₄₋₁₀ on conditioned avoidance behavior of hypophysectomized rats. *Neuroendocrinology* 11: 137-143, 1973.
3. De Wied, D. Effects of peptide hormones on behavior. In: *Frontiers in Neuroendocrinology*, edited by W. F. Ganong and L. Martini. London: Oxford University Press, 1969, pp. 97-140.
4. De Wied, D. Pituitary-adrenal system hormones and behavior. In: *The Neurosciences Third Study Program*, edited by F. O. Schmitt and F. G. Worden. Cambridge, Mass.: MIT Press, 1974, pp. 653-666.
5. De Wied, D. and W. de Jong. Drug effects and hypothalamic-anterior pituitary function. *A. Rev. Pharmac.* 14: 389-412, 1974.
6. De Wied, D., A. Witter and H. M. Greven. Behaviorally active analogues. *Biochem. Pharmac.* 24: 1463-1468, 1975.
7. Dunn, A. J. Biochemical correlates of training experiences: A discussion of the evidence. In: *Neural Mechanisms of Learning and Memory*, edited by M. R. Rosenzweig and E. L. Bennett. Cambridge, Mass.: MIT Press, 1976, pp. 316-320.
8. Dunn, A. J., P. Iuvone and D. H. Rees. Neurochemical responses of mice to ACTH and lysine vasopressin. *Pharmac. Biochem. Behav.* 5: Suppl. 1, 139-145, 1976.
9. Flood, J. F., E. L. Bennett, A. E. Orme and M. R. Rosenzweig. Effects of protein synthesis inhibition on memory for active avoidance training. *Physiol. Behav.* 14: 177-184, 1975.

10. Flood, J. F., E. L. Bennett, A. E. Orme and M. R. Rosenzweig. Relation of memory formation to controlled amounts of brain protein synthesis. *Physiol. Behav.* 15: 97-102, 1975.
11. Flood, J. F., E. L. Bennett, M. R. Rosenzweig and A. E. Orme. Influence of training strength on amnesia induced by pre-training injections of cycloheximide. *Physiol. Behav.* 9: 589-600, 1972.
12. Flood, J. F., E. L. Bennett, M. R. Rosenzweig and A. E. Orme. The influence of duration of protein synthesis inhibition on memory. *Physiol. Behav.* 10: 555-562, 1973.
13. Flood, J. F., E. L. Bennett, M. R. Rosenzweig and A. E. Orme. Comparison of the effects of anisomycin on memory across six strains of mice. *Behav. Biol.* 10: 147-160, 1974.
14. Flood, J. F. and M. E. Jarvik. Drug influences on learning and memory. In: *Neural Mechanisms of Learning and Memory*, edited by M. R. Rosenzweig and E. L. Bennett. Cambridge, Mass.: MIT Press, 1976, pp. 483-507.
15. Garrud, P., J. A. Gray and D. de Wied. Pituitary-adrenal hormones and extinction of rewarded behavior in the rat. *Physiol. Behav.* 12: 109-119, 1974.
16. Gispen, W. H., M. E. A. Reith, P. Schotman, V. M. Wiegant, H. Zwijs and D. de Wied. The CNS and ACTH-like peptides: Neurochemical responses and interaction with opiates. In: *Neuropeptide Influences on Brain and Behavior Mechanisms*, edited by L. H. Miller, C. A. Sandman and A. J. Kastin. New York: Raven Press, in press.
17. Gray, J. A. and P. Garrud. Adreno-pituitary hormones and frustrative non-reward. In: *Neuropeptide Influences on Brain and Behavior Mechanisms*, edited by L. H. Miller, C. A. Sandman and A. J. Kastin. New York: Raven Press, in press.
18. Greengard, P. Possible role for cyclic nucleotides and phosphorylated membrane proteins in postsynaptic actions of neurotransmitters. *Nature* 260: 101-108, 1976.
19. Greven, H. M. and D. de Wied. The influence of peptides derived from corticotrophin (ACTH) on performance: Structure activity studies. In: *Progress in Brain Research* V. 39, "Drug Effects on Neuroendocrine Regulation," edited by F. Zimmermann, W. H. Gispen, B. H. Marks and D. de Wied. Amsterdam: Elsevier Scientific Publishing Co., 1973, pp. 429-442.
20. Reading, H. W. and A. J. Dewar. Effects of ACTH₄₋₁₀ on cerebral RNA and protein metabolism in the rat. Abst. 3rd Intern. Meeting of the Intern. Soc. for Neurochemistry, Budapest, 1971, p. 199.
21. Rees, H., A. Dunn and P. M. Iuvone. Behavioral and biochemical responses of mice to the intraventricular administration of ACTH analogs and lysine vasopressin. *Life Sci.* 18: 1333-1340, 1976.
22. Reith, M. E., P. Schotman and W. H. Gispen. Hypophysectomy, ACTH₁₋₁₀ and *in vitro* protein synthesis in rat brain stem slices. *Brain Res.* 81: 571-575, 1974.
23. Riger, H. and H. van Riezen. The anti-amnesic effect of ACTH₄₋₁₀: Its independence of the nature of the amnesic agent and the behavioral test. *Physiol. Behav.* 14: 563-566, 1975.
24. Riger, H., H. van Riezen and D. de Wied. The effects of ACTH- and vasopressin-analogues on CO₂-induced retrograde amnesia in rats. *Physiol. Behav.* 13: 381-388, 1974.
25. Schotman, P. and W. H. Gispen. Analogues of ACTH, conditioned avoidance behavior and metabolism in brain of rat. In: *Central Nervous System: Studies on Metabolic Regulation and Function*, edited by E. Genazzani and H. Herkin. New York: Springer-Verlag, 1974, pp. 231-235.
26. Schotman, P., W. H. Gispen, H. S. Jansz and D. de Wied. Effect of ACTH analogues on macromolecule metabolism in the brain stem of hypophysectomized rats. *Brain Res.* 46: 349-362, 1972.
27. Squire, L. R. and S. H. Barondes. Anisomycin, like other inhibitors of cerebral protein synthesis, impairs "long-term" memory of a discrimination task. *Brain Res.* 66: 301-308, 1974.
28. Squire, L. R. and H. P. Davis. Cerebral protein synthesis inhibition and discrimination training: Effects of extent and duration of inhibition. *Behav. Biol.* 13: 49-57, 1975.
29. van Wimersma Greidanus, Tj. B. and D. de Wied. Effects of systemic intracerebral administration of two opposite acting ACTH-related peptides on extinction of conditioned avoidance behavior. *Neuroendocrinology* 7: 291-301, 1971.
30. Walter, R. Biochemistry of the neuropeptides. In: *Neuropeptide Influences on Brain and Behavior Mechanisms*, edited by L. H. Miller, C. A. Sandman and A. J. Kastin. New York: Raven Press, in press.