

Search for bottom squarks in pp collisions at $\sqrt{s}=1.8$ TeV

B. Abbott,⁴³ M. Abolins,⁴⁰ V. Abramov,¹⁸ B. S. Acharya,¹¹ I. Adam,⁴² D. L. Adams,⁵² M. Adams,²⁷ S. Ahn,²⁶ V. Akimov,¹⁶ G. A. Alves,² N. Amos,³⁹ E. W. Anderson,³³ M. M. Baarmand,⁴⁵ V. V. Babintsev,¹⁸ L. Babukhadia,¹⁹ A. Baden,³⁶ B. Baldin,²⁶ S. Banerjee,¹¹ J. Bantly,⁴⁹ E. Barberis,²⁰ P. Baringer,³⁴ J. F. Bartlett,²⁶ A. Belyaev,¹⁷ S. B. Beri,⁹ I. Bertram,²⁹ V. A. Bezzubov,¹⁸ P. C. Bhat,²⁶ V. Bhatnagar,⁹ M. Bhattacharjee,⁴⁵ N. Biswas,³¹ G. Blazey,²⁸ S. Blessing,²⁴ P. Bloom,²¹ A. Boehnlein,²⁶ N. I. Bojko,¹⁸ F. Borchering,²⁶ C. Boswell,²³ A. Brandt,²⁶ R. Breedon,²¹ G. Briskin,⁴⁹ R. Brock,⁴⁰ A. Bross,²⁶ D. Buchholz,²⁹ V. S. Burtovoi,¹⁸ J. M. Butler,³⁷ W. Carvalho,² D. Casey,⁴⁰ Z. Casilum,⁴⁵ H. Castilla-Valdez,¹⁴ D. Chakraborty,⁴⁵ S. V. Chekulaev,¹⁸ W. Chen,⁴⁵ S. Choi,¹³ S. Chopra,²⁴ B. C. Choudhary,²³ J. H. Christenson,²⁶ M. Chung,²⁷ D. Claes,⁴¹ A. R. Clark,²⁰ W. G. Cobau,³⁶ J. Cochran,²³ L. Coney,³¹ W. E. Cooper,²⁶ D. Coppage,³⁴ C. Cretsinger,⁴⁴ D. Cullen-Vidal,⁴⁹ M. A. C. Cummings,²⁸ D. Cutts,⁴⁹ O. I. Dahl,²⁰ K. Davis,¹⁹ K. De,⁵⁰ K. Del Signore,³⁹ M. Demarteau,²⁶ D. Denisov,²⁶ S. P. Denisov,¹⁸ H. T. Diehl,²⁶ M. Diesburg,²⁶ G. Di Loreto,⁴⁰ P. Draper,⁵⁰ Y. Ducros,⁸ L. V. Dudko,¹⁷ S. R. Dugad,¹¹ A. Dyshkant,¹⁸ D. Edmunds,⁴⁰ J. Ellison,²³ V. D. Elvira,⁴⁵ R. Engelmann,⁴⁵ S. Eno,³⁶ G. Eppley,⁵² P. Ermolov,¹⁷ O. V. Eroshin,¹⁸ V. N. Evdokimov,¹⁸ T. Fahland,²² M. K. Fatyga,⁴⁴ S. Feher,²⁶ D. Fein,¹⁹ T. Ferbel,⁴⁴ H. E. Fisk,²⁶ Y. Fisyak,⁴⁶ E. Flattum,²⁶ G. E. Forden,¹⁹ M. Fortner,²⁸ K. C. Frame,⁴⁰ S. Fuess,²⁶ E. Gallas,⁵⁰ A. N. Galyaev,¹⁸ P. Gartung,²³ V. Gavrilov,¹⁶ T. L. Geld,⁴⁰ R. J. Genik II,⁴⁰ K. Genser,²⁶ C. E. Gerber,²⁶ Y. Gershtein,⁴⁹ B. Gibbard,⁴⁶ B. Gobbi,²⁹ B. Gómez,⁵ G. Gómez,³⁶ P. I. Goncharov,¹⁸ J. L. González Solís,¹⁴ H. Gordon,⁴⁶ L. T. Goss,⁵¹ K. Gounder,²³ A. Goussiou,⁴⁵ N. Graf,⁴⁶ P. D. Grannis,⁴⁵ D. R. Green,²⁶ H. Greenlee,²⁶ S. Grinstein,¹ P. Grudberg,²⁰ S. Grünendahl,²⁶ G. Guglielmo,⁴⁸ J. A. Guida,¹⁹ J. M. Guida,⁴⁹ A. Gupta,¹¹ S. N. Gurzhiev,¹⁸ G. Gutierrez,²⁶ P. Gutierrez,⁴⁸ N. J. Hadley,³⁶ H. Haggerty,²⁶ S. Hagopian,²⁴ V. Hagopian,²⁴ K. S. Hahn,⁴⁴ R. E. Hall,²² P. Hanlet,³⁸ S. Hansen,²⁶ J. M. Hauptman,³³ C. Hebert,³⁴ D. Hedin,²⁸ A. P. Heinson,²³ U. Heintz,³⁷ R. Hernández-Montoya,¹⁴ T. Heuring,²⁴ R. Hirosky,²⁷ J. D. Hobbs,⁴⁵ B. Hoeneisen,⁶ J. S. Hoftun,⁴⁹ F. Hsieh,³⁹ Tong Hu,³⁰ A. S. Ito,²⁶ S. A. Jerger,⁴⁰ R. Jesik,³⁰ T. Joffe-Minor,²⁹ K. Johns,¹⁹ M. Johnson,²⁶ A. Jonckheere,²⁶ M. Jones,²⁵ H. Jöstlein,²⁶ S. Y. Jun,²⁹ C. K. Jung,⁴⁵ S. Kahn,⁴⁶ D. Karmanov,¹⁷ D. Karmgard,²⁴ R. Kehoe,³¹ S. K. Kim,¹³ B. Klima,²⁶ C. Klopfenstein,²¹ W. Ko,²¹ J. M. Kohli,⁹ D. Koltick,³² A. V. Kostitskiy,¹⁸ J. Kotcher,⁴⁶ A. V. Kotwal,⁴² A. V. Kozelov,¹⁸ E. A. Kozlovsky,¹⁸ J. Krane,⁴¹ M. R. Krishnaswamy,¹¹ S. Krzywdzinski,²⁶ S. Kuleshov,¹⁶ Y. Kulik,⁴⁵ S. Kunori,³⁶ F. Landry,⁴⁰ G. Landsberg,⁴⁹ B. Lauer,³³ A. Leflat,¹⁷ J. Li,⁵⁰ Q. Z. Li,²⁶ J. G. R. Lima,³ D. Lincoln,²⁶ S. L. Linn,²⁴ J. Linnemann,⁴⁰ R. Lipton,²⁶ A. Lucotte,⁴⁵ L. Lueking,²⁶ A. L. Lyon,³⁶ A. K. A. Maciel,²⁸ R. J. Madaras,²⁰ R. Madden,²⁴ L. Magaña-Mendoza,¹⁴ V. Manankov,¹⁷ S. Mani,²¹ H. S. Mao,⁴ R. Markeloff,²⁸ T. Marshall,³⁰ M. I. Martin,²⁶ K. M. Mauritz,³³ B. May,²⁹ A. A. Mayorov,¹⁸ R. McCarthy,⁴⁵ J. McDonald,²⁴ T. McKibben,²⁷ J. McKinley,⁴⁰ T. McMahon,⁴⁷ H. L. Melanson,²⁶ M. Merkin,¹⁷ K. W. Merritt,²⁶ C. Miao,⁴⁹ H. Miettinen,⁵² A. Mincer,⁴³ C. S. Mishra,²⁶ N. Mokhov,²⁶ N. K. Mondal,¹¹ H. E. Montgomery,²⁶ P. Mooney,⁵ M. Mostafa,¹ H. da Motta,² C. Murphy,²⁷ F. Nang,¹⁹ M. Narain,³⁷ V. S. Narasimham,¹¹ A. Narayanan,¹⁹ H. A. Neal,³⁹ J. P. Negret,⁵ P. Nemethy,⁴³ D. Norman,⁵¹ L. Oesch,³⁹ V. Oguri,³ N. Oshima,²⁶ D. Owen,⁴⁰ P. Padley,⁵² A. Para,²⁶ N. Parashar,³⁸ Y. M. Park,¹² R. Partridge,⁴⁹ N. Parua,⁷ M. Paterno,⁴⁴ B. Pawlik,¹⁵ J. Perkins,⁵⁰ M. Peters,²⁵ R. Piegai,¹ H. Piekarczyk,²⁴ Y. Pischalnikov,³² B. G. Pope,⁴⁰ H. B. Prosper,²⁴ S. Protopopescu,⁴⁶ J. Qian,³⁹ P. Z. Quintas,²⁶ R. Raja,²⁶ S. Rajagopalan,⁴⁶ O. Ramirez,²⁷ S. Reucroft,³⁸ M. Rijssenbeek,⁴⁵ T. Rockwell,⁴⁰ M. Roco,²⁶ P. Rubinov,²⁹ R. Ruchti,³¹ J. Rutherford,¹⁹ A. Sánchez-Hernández,¹⁴ A. Santoro,² L. Sawyer,³⁵ R. D. Schamberger,⁴⁵ H. Schellman,²⁹ J. Sculli,⁴³ E. Shabalina,¹⁷ C. Shaffer,²⁴ H. C. Shankar,¹¹ R. K. Shivpuri,¹⁰ D. Shpakov,⁴⁵ M. Shupe,¹⁹ H. Singh,²³ J. B. Singh,⁹ V. Sirotenko,²⁸ E. Smith,⁴⁸ R. P. Smith,²⁶ R. Snihur,²⁹ G. R. Snow,⁴¹ J. Snow,⁴⁷ S. Snyder,⁴⁶ J. Solomon,²⁷ M. Sosebee,⁵⁰ N. Sotnikova,¹⁷ M. Souza,² G. Steinbrück,⁴⁸ R. W. Stephens,⁵⁰ M. L. Stevenson,²⁰ F. Stichelbaut,⁴⁶ D. Stoker,²² V. Stolin,¹⁶ D. A. Stoyanova,¹⁸ M. Strauss,⁴⁸ K. Streets,⁴³ M. Strovink,²⁰ A. Sznajder,² P. Tamburello,³⁶ J. Tarazi,²² M. Tartaglia,²⁶ T. L. T. Thomas,²⁹ J. Thompson,³⁶ T. G. Trippe,²⁰ P. M. Tuts,⁴² V. Vaniev,¹⁸ N. Varelas,²⁷ E. W. Varnes,²⁰ A. A. Volkov,¹⁸ A. P. Vorobiev,¹⁸ H. D. Wahl,²⁴ G. Wang,²⁴ J. Warchol,³¹ G. Watts,⁴⁹ M. Wayne,³¹ H. Weerts,⁴⁰ A. White,⁵⁰ J. T. White,⁵¹ J. A. Wightman,³³ S. Willis,²⁸ S. J. Wimpenny,²³ J. V. D. Wirjawan,⁵¹ J. Womersley,²⁶ D. R. Wood,³⁸ R. Yamada,²⁶ P. Yamin,⁴⁶ T. Yasuda,³⁸ P. Yepes,⁵² K. Yip,²⁶ C. Yoshikawa,²⁵ S. Youssef,²⁴ J. Yu,²⁶ Y. Yu,¹³ B. Zhang,⁴ Z. Zhou,³³ Z. H. Zhu,⁴⁴ M. Zielinski,⁴⁴ D. Zieminska,³⁰ A. Zieminski,³⁰ V. Zutshi,⁴⁴ E. G. Zverev,¹⁷ and A. Zylberstein⁸

(DØ Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Institute of High Energy Physics, Beijing, People's Republic of China⁵Universidad de los Andes, Bogotá, Colombia⁶Universidad San Francisco de Quito, Quito, Ecuador⁷Institut des Sciences Nucléaires, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France⁸DAPNIA/Service de Physique des Particules, CEA, Saclay, France

- ⁹Panjab University, Chandigarh, India
¹⁰Delhi University, Delhi, India
¹¹Tata Institute of Fundamental Research, Mumbai, India
¹²Kyungshung University, Pusan, Korea
¹³Seoul National University, Seoul, Korea
¹⁴CINVESTAV, Mexico City, Mexico
¹⁵Institute of Nuclear Physics, Kraków, Poland
¹⁶Institute for Theoretical and Experimental Physics, Moscow, Russia
¹⁷Moscow State University, Moscow, Russia
¹⁸Institute for High Energy Physics, Protvino, Russia
¹⁹University of Arizona, Tucson, Arizona 85721
²⁰Lawrence Berkeley National Laboratory, Berkeley, California 94720
and University of California, Berkeley, California 94720
²¹University of California, Davis, California 95616
²²University of California, Irvine, California 92697
²³University of California, Riverside, California 92521
²⁴Florida State University, Tallahassee, Florida 32306
²⁵University of Hawaii, Honolulu, Hawaii 96822
²⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510
²⁷University of Illinois at Chicago, Chicago, Illinois 60607
²⁸Northern Illinois University, DeKalb, Illinois 60115
²⁹Northwestern University, Evanston, Illinois 60208
³⁰Indiana University, Bloomington, Indiana 47405
³¹University of Notre Dame, Notre Dame, Indiana 46556
³²Purdue University, West Lafayette, Indiana 47907
³³Iowa State University, Ames, Iowa 50011
³⁴University of Kansas, Lawrence, Kansas 66045
³⁵Louisiana Tech University, Ruston, Louisiana 71272
³⁶University of Maryland, College Park, Maryland 20742
³⁷Boston University, Boston, Massachusetts 02215
³⁸Northeastern University, Boston, Massachusetts 02115
³⁹University of Michigan, Ann Arbor, Michigan 48109
⁴⁰Michigan State University, East Lansing, Michigan 48824
⁴¹University of Nebraska, Lincoln, Nebraska 68588
⁴²Columbia University, New York, New York 10027
⁴³New York University, New York, New York 10003
⁴⁴University of Rochester, Rochester, New York 14627
⁴⁵State University of New York, Stony Brook, New York 11794
⁴⁶Brookhaven National Laboratory, Upton, New York 11973
⁴⁷Langston University, Langston, Oklahoma 73050
⁴⁸University of Oklahoma, Norman, Oklahoma 73019
⁴⁹Brown University, Providence, Rhode Island 02912
⁵⁰University of Texas, Arlington, Texas 76019
⁵¹Texas A&M University, College Station, Texas 77843
⁵²Rice University, Houston, Texas 77005

(Received 12 March 1999; published 23 June 1999)

We report on a search for bottom squarks ($\tilde{b}$) produced in $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV using the DØ detector at Fermilab. Bottom squarks are assumed to be produced in pairs and to decay to the lightest supersymmetric particle (LSP) and a b quark with a branching fraction of 100%. The LSP is assumed to be the lightest neutralino and stable. We set limits on the production cross section as a function of $\tilde{b}$ mass and LSP mass. [S0556-2821(99)50313-1]

PACS number(s): 14.80.Ly, 13.85.Rm

Supersymmetry (SUSY) is a hypothetical fundamental space-time symmetry relating bosons and fermions [1]. Supersymmetric extensions to the standard model (SM) feature as yet undiscovered supersymmetric partners for every SM

particle. The scalar quarks (squarks) $\tilde{q}_L$ and $\tilde{q}_R$ are the partners of the left-handed and right-handed quarks, respectively. These are weak eigenstates, and can mix to form the mass eigenstates, with $\tilde{q}_1 = \tilde{q}_L \cos \theta + \tilde{q}_R \sin \theta$ for the lighter squark,




FIG. 1. The expected distributions of E_T for $m_{\tilde{b}}$ values of 70 (a) and 100 (b) GeV/c^2 , for the indicated values of m_{LSP} [7].

and the orthogonal combination for the heavier squark $\tilde{q}_2$. In most SUSY models, the masses of the squarks are approximately degenerate, but in some models, the lighter top and bottom squarks could have a lower mass than the other squarks because of the high mass values of the top and bottom quarks. In particular, lighter bottom squarks could arise for large values of $\tan\beta$, the ratio of the vacuum expectation values of the two Higgs fields in the minimal supersymmetric standard model.

We report the results of a mixing-independent search for bottom squarks produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. Squarks are produced in pairs by QCD processes with the production cross section depending on the mass of the squark, but not on the mixing angle θ . We search for events where both squarks decay to the lightest neutralino $\tilde{\chi}_1^0$ via $\tilde{b} \rightarrow \tilde{\chi}_1^0 + b$ and assume that the $\tilde{\chi}_1^0$ is the lightest supersymmetric particle (LSP) and stable. This should be the dominant decay channel provided that the mass of the squark ($m_{\tilde{b}}$) is larger than the combined masses of the b quark and LSP (m_{LSP}); therefore we assume its branching fraction is 100%. This yields a final state consisting of two b quarks and two unobserved stable particles resulting in missing transverse energy (E_T) in the detector. In this paper, we give limits on the squark pair production cross section for different values of $m_{\tilde{b}}$ and m_{LSP} . Limits on the cross section are used to exclude a region in the $(m_{\text{LSP}}, m_{\tilde{b}})$ plane. Limits [2] from the CERN e^+e^- collider (LEP) experiments depend on the Z/γ -to-squark coupling, which is a function of the mixing angle. For maximal coupling, the LEP exclusion region can extend to the kinematic maximum; for example, to about 85 GeV/c^2 at $\sqrt{s} = 183$ GeV.

The data used for our analysis were collected during 1992–1996 by the DØ detector [3] at the Fermilab Tevatron Collider. The DØ detector is composed of three major sys-

TABLE I. Total efficiencies for different $m_{\tilde{b}}$ and m_{LSP} values for the four channels, and 95% C.L. limits on the production cross section obtained by combining all channels.

$m_{\tilde{b}}$ (GeV/c^2)	m_{LSP} (GeV/c^2)	E_T + jets	Total efficiency ($\times 10^{-3}$)			σ limit (pb)
			dimuon	single muon low- p_T	high- p_T	
70	30	18	0.13	2.2	0.3	32
70	50	4	0.02	0.6	0.1	245
85	40	29	0.20	3.9	0.6	18.8
85	60	11	0.04	1.0	0.1	84
100	20	43	0.50	9.5	1.9	9.3
100	40	34	0.27	7.0	1.3	12.6
100	50	30	0.30	5.8	1.0	14.7
115	40	51	0.54	10.9	2.0	8.0

tems: an inner detector for tracking charged particles, a uranium-liquid argon calorimeter for measuring electromagnetic and hadronic energies, and a muon spectrometer consisting of a magnetized iron toroid and three layers of drift tubes. The detector measures jets with an energy resolution of approximately $\sigma/E = 0.8/\sqrt{E}$ (E in GeV) and muons with a momentum resolution of $\sigma/p = \{[0.18(p-2)/p]^2 + (0.003p)^2\}^{1/2}$ (p in GeV/c). E_T is determined by summing the calorimeter and muon transverse energies, and is measured with a resolution of $\sigma = 1.08 \text{ GeV} + 0.019(\sum |E_T|)$ [4].

Four channels are combined to set limits on the production of bottom squarks. The first required a E_T and jets topology. This channel was previously used to set limits on the mass of the top squark, which was assumed to decay $\tilde{t} \rightarrow \tilde{\chi}_1^0 + c$ [5]. The other three channels, in addition, required that at least one jet has an associated muon, thereby tagging b quark decay, and were used to set limits on a charge 1/3 third generation leptoquark for the decay $LQ \rightarrow \nu_\tau + b$ [6]. We use identical data samples and event selections for the bottom squark limits presented in this paper. For all channels, the presence of significant E_T is used to identify the non-interacting LSPs. Figure 1 shows the expected E_T distribution for two values of $m_{\tilde{b}}$ and different m_{LSP} [7]. Our requirement that $E_T > 35$ –40 GeV reduces the acceptance for small values of the mass difference $m_{\tilde{b}} - m_{\text{LSP}}$. Backgrounds arise from events where neutrinos produce significant E_T ; for example, in W + jets events, where $W \rightarrow l\nu$.

Events for the E_T + jets channel were collected using a trigger that required $E_T > 35$ GeV. The offline analysis required two jets ($E_T^{\text{jet}} > 30$ GeV), $E_T > 40$ GeV, and no isolated electrons or muons. Events had to have only one primary vertex to assure an unambiguous calculation of E_T . To eliminate QCD backgrounds, additional cuts were made on the angles between the two jets, and between jets and the direction of the E_T . Data with an integrated luminosity of 7.4 pb^{-1} , satisfying the above selection criteria, yielded three candidate events. Background was estimated to be 3.5 ± 1.2 events, with 3.0 ± 0.9 events from W boson decays and




FIG. 2. The 95% C.L. exclusion contour in the $(m_{\text{LSP}}, m_{\tilde{b}})$ plane. Also shown are the results from the ALEPH experiment at LEP for minimal ($\theta=68^\circ$) and maximal ($\theta=0^\circ$) coupling [2].

0.5 ± 0.3 events from Z boson decays [5].

The trigger for the muon channels required either two low- p_T muons ($p_T^\mu > 3.0$ GeV/ c), or a single low- p_T muon and a jet with $E_T > 10$ GeV, or a high- p_T muon ($p_T^\mu > 15$ GeV/ c) and a jet with $E_T > 15$ GeV. Integrated luminosities of 60.1 pb $^{-1}$, 19.5 pb $^{-1}$, and 92.4 pb $^{-1}$, respectively, were collected using the three muon triggers. The offline analysis used muons in the pseudorapidity range $|\eta_\mu| < 1.0$ and $p_T^\mu > 3.5$ GeV/ c , while jets were required to have $E_T > 10$ GeV. For events with two muons, each muon had to be associated with its own jet. In single muon events, the muon was required to be associated with a jet, and an additional jet with $E_T > 25$ GeV was also required. To remove QCD backgrounds, events were selected with $\cancel{E}_T > 35$ GeV and an azimuthal angular separation between the $\cancel{E}_T$ and the nearest jet of > 0.7 radians. For the single muon channels, backgrounds from W boson decays were reduced by cuts on muon-jet correlations, while background from top quark production was minimized by cuts on the scalar sum of jet E_T . After imposition of all selection criteria, two events remained in the data.

We considered background contributions to the muon channels from $t\bar{t}$ and W and Z boson decays [6]. Top quark events have multiple b quarks and $\cancel{E}_T$, and we estimated that 1.4 ± 0.5 $t\bar{t}$ events remained in our sample. W and Z events have $\cancel{E}_T$ from $W \rightarrow l\nu$ or $Z \rightarrow \nu\nu$. They can also have muons near jets that can mimic b quark decays when a prompt muon overlaps a jet, or a jet fragments into a muon via a c quark or a π/K decay. We estimated there were 1.0 ± 0.4 W boson events and 0.1 ± 0.1 Z boson events in the sample. The total background for the muon channels was therefore 2.5 ± 0.6 events.

Combining the four channels yields five events, with a total estimated background of 6.0 ± 1.3 events. We set limits on the cross section by combining the detection efficiencies and integrated luminosities for the different channels. We calculate the detection efficiency using Monte Carlo (MC) generated acceptances [7], multiplied by trigger and reconstruction efficiencies obtained from data [5,6]. The total efficiencies for different squark and neutralino masses are summarized in Table I. Using a muon to tag b quark decays reduces the efficiency for those channels, but their higher integrated luminosities yield a sensitivity comparable to that of the $\cancel{E}_T + \text{jets}$ channel. Including systematic errors and statistics for the MC, the total uncertainty on the combined efficiency varies between 8.6% and 29%, depending on the assumed masses. The jet energy scale dominates the systematic error for $m_{\tilde{b}} = 70$ GeV/ c^2 , while uncertainties on the muon trigger and reconstruction efficiency dominate at higher squark masses. The 95% confidence level (C.L.) upper limits on the pair production cross section are determined using Bayesian methods, and include the systematic uncertainty on the efficiency and a 5.3% uncertainty in the integrated luminosity. The resulting upper limits are given in Table I for different values of $m_{\tilde{b}}$ and m_{LSP} .

We use the program PROSPINO [8] to calculate the bottom squark pair production cross section as a function of $m_{\tilde{b}}$. The cross section is evaluated assuming a renormalization scale $\mu = m_{\tilde{b}}$. The program includes next-to-leading order diagrams, and uses CTEQ4M parton distribution functions [9]. For any given $m_{\tilde{b}}$, we determine the value of m_{LSP} where our 95% C.L. limit intersects the theoretical cross section. The excluded region in the $(m_{\text{LSP}}, m_{\tilde{b}})$ plane is shown in Fig. 2. We exclude values of $m_{\tilde{b}}$ below 115 GeV/ c^2 for $m_{\text{LSP}} < 20$ GeV/ c^2 . For $m_{\tilde{b}} = 85$ GeV/ c^2 , we exclude the region with $m_{\text{LSP}} < 47$ GeV/ c^2 . Also shown are limits [2] from ALEPH for $\sqrt{s} = 181\text{--}184$ GeV. For most allowable values of m_{LSP} , they exclude the region with $m_{\tilde{b}} < 83$ GeV/ c^2 , assuming maximal coupling ($\theta=0^\circ$) [10].

In conclusion, we observe five candidate events consistent with the final state $b\bar{b} + \cancel{E}_T$. We estimate that 6.0 ± 1.3 events are expected from $t\bar{t}$ and W and Z boson production, and find no excess of events that can be attributed to bottom squark production. We interpret our result as an excluded region in the $(m_{\text{LSP}}, m_{\tilde{b}})$ plane. This result is independent of the mixing between $\tilde{b}_L$ and $\tilde{b}_r$.

We thank S.P. Martin and M. Spira for their assistance. We thank the Fermilab and collaborating institution staffs for contributions to this work and acknowledge support from the Department of Energy and National Science Foundation (USA), Commissariat à l'Energie Atomique (France), Ministry for Science and Technology and Ministry for Atomic Energy (Russia), CAPES and CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), and CONICET and UBACyT (Argentina).

- [1] See, e.g., S.P. Martin, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998), hep-ph/9709356, and references therein.
- [2] ALEPH Collaboration, R. Barate *et al.*, Phys. Lett. B **434**, 189 (1998); OPAL Collaboration, K. Ackerstaff *et al.*, Eur. Phys. J. C **6**, 225 (1999); DELPHI Collaboration, P. Abreu *et al.*, *ibid.* **6**, 385 (1999); L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. B **445**, 428 (1999).
- [3] DØ Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res. A **338**, 185 (1994).
- [4] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. D **52**, 4877 (1995).
- [5] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **76**, 2222 (1996).
- [6] DØ Collaboration, B. Abbott *et al.*, Phys. Rev. Lett. **81**, 38 (1998).
- [7] Monte Carlo samples were generated with ISAJET. F. Paige and S. Protopopescu, BNL Report No. BNL38034, 1986, release v6.49. The simulation of the detector, trigger, and offline selections used GEANT. R. Brun and F. Carminati, CERN Program Library Long Writeup W5013, 1993.
- [8] W. Beenakker, M. Kramer, T. Plehn, M. Spira and P.M. Zerwas, Nucl. Phys. **B515**, 3 (1998); M. Spira (private communication).
- [9] H.L. Lai *et al.*, Phys. Rev. D **55**, 1280 (1997).
- [10] Preliminary ALEPH results for $\sqrt{s}=189$ GeV extend the excluded region to $m_{\tilde{b}} < 90$ GeV/ c^2 for maximal coupling ($\theta = 0^\circ$) and 75–80 GeV/ c^2 for minimal coupling ($\theta = 68^\circ$). M. Berggren, presented at the DPF99 Meeting at UCLA, 1999.