

## Permutation symmetry for neutrino and charged-lepton mass matrices

Ernest Ma

Physics Department, University of California, Riverside, California 92521

(Received 3 September 1999; published 11 January 2000)

Permutation symmetry  $S_3$  is applied to obtain two equal Majorana neutrino masses, while maintaining three different charged-lepton masses and suppressing neutrinoless double beta decay. The resulting radiative splitting of the two neutrinos is shown to be suitable for solar neutrino vacuum oscillations.

PACS number(s): 14.60.Pq, 11.30.Hv, 12.15.Ff

## I. INTRODUCTION

There are three known charged leptons,  $e$ ,  $\mu$ ,  $\tau$ , with very different masses:

$$\begin{aligned} m_e &= 0.511 \text{ MeV}, & m_\mu &= 105.66 \text{ MeV}, \\ m_\tau &= 1777 \text{ MeV}. \end{aligned} \quad (1)$$

Their accompanying neutrinos,  $\nu_e$ ,  $\nu_\mu$ ,  $\nu_\tau$ , are not necessarily mass eigenstates. In general,

$$\nu_\alpha = \sum_{i=1}^3 U_{\alpha i} \nu_i, \quad (2)$$

where  $\alpha = e, \mu, \tau$  and  $\nu_i$  are mass eigenstates and

$$U = \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta / \sqrt{2} & \cos \theta / \sqrt{2} & -1/\sqrt{2} \\ -\sin \theta / \sqrt{2} & \cos \theta / \sqrt{2} & 1/\sqrt{2} \end{bmatrix} \quad (3)$$

is a typical mixing matrix which allows one to understand recent atmospheric neutrino data [1] and the long-standing solar neutrino deficit [2] in terms of neutrino oscillations. The form of  $U$  in Eq. (3) has been advocated by many authors [3]. It has the virtue of maximal mixing between  $\nu_\mu$  and  $\nu_\tau$  which agrees well with the atmospheric data, and it allows the solar data to be interpreted with either the small-angle or large-angle matter-enhanced neutrino-oscillation solution [4] or the necessarily large-angle vacuum solution.

The masses  $m_{1,2,3}$  are now subject to the conditions that

$$\Delta m_{13}^2 \simeq \Delta m_{23}^2 \sim 10^{-3} \text{ eV}^2 \quad (4)$$

for atmospheric neutrino oscillations and

$$\Delta m_{12}^2 \sim 10^{-5} \text{ or } 10^{-10} \text{ eV}^2 \quad (5)$$

for solar matter-enhanced or vacuum neutrino oscillations, where  $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$ . This allows for the intriguing possibility that neutrino masses are degenerate [5] with very small splittings. However, since the charged-lepton masses break this degeneracy, there must be radiative corrections [6] which may or may not be consistent with the actual phenomenological solutions desired, especially so if a value of  $10^{-10} \text{ eV}^2$  for  $\Delta m_{12}^2$  is to be maintained.

In this paper, a detailed model of lepton mass matrices is presented, based on permutation symmetry  $S_3$ . It is the outgrowth of a previous proposal [7] which shows how  $\Delta m_{12}^2$  of order  $10^{-10} \text{ eV}^2$  for solar neutrino vacuum oscillations can be obtained as a second-order perturbation of a twofold degenerate neutrino mass matrix, resulting in a successful formula relating atmospheric and solar neutrino oscillations. In Sec. II the model is described and it is shown how the breaking of  $S_3$  together with electroweak gauge symmetry allows the charged-lepton masses to be all different while maintaining a twofold degeneracy in the neutrino mass matrix  $\mathcal{M}_\nu$  at the tree level. In Sec. III radiative corrections to  $\mathcal{M}_\nu$  are derived in terms of the parameters of the charged-lepton mass matrix  $\mathcal{M}_l$ . The latter is chosen such that neutrinoless double beta decay [8] is absent at the tree level. In Sec. IV the resulting phenomenon of lepton flavor nonconservation beyond that of neutrino oscillations is discussed in the context of other processes involving charged leptons. Finally in Sec. V there are some concluding remarks.

II. LEPTON MASS MATRICES UNDER  $S_3$ 

Let the three families of leptons be denoted by  $(\nu_i, l_i)_L$  and  $l_i^c$ ,  $i = 1, 2, 3$ . In this convention,  $l_i l_j^c$  is a Dirac mass term for the charged leptons (instead of the usual  $\bar{l}_i l_j$ ) and  $\nu_i \nu_j$  is a Majorana mass term for the neutrinos. Consider the discrete permutation symmetry  $S_3$ . Its irreducible representations are  $\underline{2}$ ,  $\underline{1}$ , and  $\underline{1}'$ , with the following multiplication rules:  $\underline{2} \times \underline{2} = \underline{2} + \underline{1} + \underline{1}'$  and  $\underline{1}' \times \underline{1}' = \underline{1}$ . Under  $S_3$ , let

$$\left[ \begin{pmatrix} \nu_1 \\ l_1 \end{pmatrix}_L, \begin{pmatrix} \nu_2 \\ l_2 \end{pmatrix}_L \right] \sim \underline{2}, \quad \begin{pmatrix} \nu_3 \\ l_3 \end{pmatrix}_L \sim \underline{1}, \quad [l_1^c, l_2^c] \sim \underline{2}, \quad l_3^c \sim \underline{1}. \quad (6)$$

The Higgs sector of this model consists of three doublets  $\Phi_i = (\phi_i^0, \phi_i^-)$ ,  $i = 1, 2, 3$ , and one triplet  $\xi = (\xi^{++}, \xi^+, \xi^0)$ . Under  $S_3$ , let

$$(\Phi_1, \Phi_2) \sim \underline{2}, \quad \Phi_3 \sim \underline{1}, \quad \xi \sim \underline{1}. \quad (7)$$

Neutrinos and charged leptons couple to  $\xi$  according to

$$f_{ij} [\xi^0 \nu_i \nu_j + \xi^+ (\nu_i l_j + l_i \nu_j) / \sqrt{2} + \xi^{++} l_i l_j] + \text{H.c.}, \quad (8)$$

where  $f_{ij}$  is restricted by  $S_3$  to have the form

$$f = \begin{pmatrix} 0 & f_0 & 0 \\ f_0 & 0 & 0 \\ 0 & 0 & f_3 \end{pmatrix}. \quad (9)$$

As shown recently [9,10], this is an equally natural way to obtain small Majorana neutrino masses as the canonical seesaw mechanism [11], because the vacuum expectation value  $\langle \xi^0 \rangle = u$  is inversely proportional to  $m_\xi^2$ . Let  $m_0 = 2f_0u$  and  $m_3 = 2f_3u$ ; then the Majorana neutrino mass matrix spanning  $\nu_{1,2,3}$  is given by

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_0 & 0 \\ m_0 & 0 & 0 \\ 0 & 0 & m_3 \end{pmatrix}. \quad (10)$$

The eigenvalues of  $\mathcal{M}_\nu$  are  $-m_0$ ,  $m_0$ , and  $m_3$ . (A negative mass here means that the corresponding Majorana neutrino is odd under  $CP$  after a  $\gamma_5$  rotation to remove the minus sign.) Hence there is an effective twofold degeneracy in the  $\nu_1$ - $\nu_2$  sector, and it corresponds to an additional global symmetry, i.e.  $L_1$ - $L_2$ , if the charged-lepton mass matrix  $\mathcal{M}_l$  is diagonal in the same basis.

There are five Yukawa interaction terms of the charged leptons with the Higgs doublets which are invariant under  $S_3$ , i.e.

$$\begin{aligned} h_1 & [l_1 l_1^c \phi_1^0 + l_2 l_2^c \phi_2^0 - \nu_1 l_1^c \phi_1^- - \nu_2 l_2^c \phi_2^-] \\ & + h_2 [(l_1 l_2^c + l_2 l_1^c) \phi_3^0 - (\nu_1 l_2^c + \nu_2 l_1^c) \phi_3^-] \\ & + h_3 [(l_1 \phi_2^0 + l_2 \phi_1^0) l_3^c - (\nu_1 \phi_2^- + \nu_2 \phi_1^-) l_3^c] \\ & + h_4 [l_3 (l_1^c \phi_2^0 + l_2^c \phi_1^0) - \nu_3 (l_1^c \phi_2^- + l_2^c \phi_1^-)] \\ & + h_5 [l_3 l_3^c \phi_3^0 - \nu_3 l_3^c \phi_3^-] + \text{H.c.} \end{aligned} \quad (11)$$

As  $\phi_{1,2,3}^0$  acquire vacuum expectation values  $v_{1,2,3}$ , the  $3 \times 3$  mass matrix linking  $l_{1,2,3}$  to  $l_{1,2,3}^c$  is given by

$$\mathcal{M}_l = \begin{bmatrix} h_1 v_1 & h_2 v_3 & h_3 v_2 \\ h_2 v_3 & h_1 v_2 & h_3 v_1 \\ h_3 v_2 & h_4 v_1 & h_5 v_3 \end{bmatrix}. \quad (12)$$

The scalar potential of this model is assumed to respect  $S_3$  only in its dimension-4 terms; i.e.,  $S_3$  is broken *softly* by its dimension-2 and dimension-3 terms:

$$\begin{aligned} & \sum_{i,j=1}^3 m_{ij}^2 (\bar{\phi}_i^0 \phi_j^0 + \phi_i^+ \phi_j^-) + m_\xi^2 (\xi^{--} \xi^{++} + \xi^- \xi^+ + \bar{\xi}^0 \xi^0) \\ & + \sum_{i,j=1}^3 \mu_{ij} \left[ \xi^{++} \phi_i^- \phi_j^- + \frac{1}{\sqrt{2}} \xi^+ (\phi_i^0 \phi_j^- + \phi_i^- \phi_j^0) \right. \\ & \left. + \xi^0 \phi^0 \phi^0 \right] + \text{H.c.} \end{aligned} \quad (13)$$

This allows  $\mathcal{M}_l$  to break  $S_3$  with  $v_2 \neq v_1$ . In fact, the limits  $h_2 = 0$  and  $v_2 = 0$  (which are of course not required by the

symmetry but are crucial for the subsequent discussion) are assumed here so that  $\mathcal{M}_l$  becomes of the form

$$\mathcal{M}_l = \begin{pmatrix} m_e & 0 & 0 \\ 0 & 0 & a \\ 0 & b & d \end{pmatrix}, \quad (14)$$

with  $l_1$  identified as the electron, and

$$m_{\tau,\mu}^2 = \frac{1}{2} (d^2 + a^2 + b^2) \pm \frac{1}{2} \sqrt{d^2 + (a+b)^2} \sqrt{d^2 + (a-b)^2}. \quad (15)$$

From Eqs. (10) and (14), the eigenstates of  $\mathcal{M}_\nu$  are easily read off:

$$\begin{aligned} \nu_1 &= \frac{1}{\sqrt{2}} (\nu_e - c \nu_\mu - s \nu_\tau), & \nu_2 &= \frac{1}{\sqrt{2}} (\nu_e + c \nu_\mu + s \nu_\tau), \\ \nu_3 &= -s \nu_\mu + c \nu_\tau, \end{aligned} \quad (16)$$

where  $c = \cos \theta_L$  and  $s = \sin \theta_L$  are determined by the  $\mu_L$ - $-\tau_L$  sector of Eq. (14). This means that  $\nu_e$  mixes maximally with  $c \nu_\mu + s \nu_\tau$ , i.e.  $\theta = \pi/4$  in Eq. (3). If  $c = s = 1/\sqrt{2}$  is also assumed in the above (corresponding to  $a^2 = b^2 + d^2$ ), the so-called bimaximal form of neutrino oscillations is obtained. In that case,

$$a^2 = \frac{1}{2} (m_\tau^2 + m_\mu^2), \quad b^2 = \frac{2m_\tau^2 m_\mu^2}{m_\tau^2 + m_\mu^2}, \quad d^2 = \frac{(m_\tau^2 - m_\mu^2)^2}{2(m_\tau^2 + m_\mu^2)}. \quad (17)$$

### III. RADIATIVE CORRECTIONS TO NEUTRINO MASS DEGENERACY

To discuss radiative corrections to  $\mathcal{M}_\nu$ , consider first the case of keeping only one Higgs doublet  $\Phi$  and one Higgs triplet  $\xi$ . In this scenario,  $S_3$  is explicitly broken by the Yukawa interactions of the charged leptons. Consequently, there is an arbitrariness in choosing the mass scale at which  $S_3$  is assumed to be exact. The most natural choice in the present context is of course  $m_\xi$ ; hence there are two contributions to the radiatively corrected  $\mathcal{M}_\nu$ . One is a finite correction to the mass matrix, as shown in Fig. 1. The other is a renormalization of the coupling matrix from the shift in mass scale from  $m_\xi$  to  $m_W$ . This was the specific case presented in Ref. [7].

Here the situation is different in two ways. First,  $S_3$  is a good symmetry as far as the Yukawa couplings are concerned. Second, it is softly broken only at the electroweak energy scale. Hence there is no  $S_3$  breaking contribution from the renormalization of the coupling matrix. As for the finite correction to the mass matrix, because of the approximations  $h_2 = 0$  and  $v_2 = 0$ , there are now only two contributions  $\nu_1 \nu_3 \bar{\phi}_1^0 \phi_3^0$  and  $\nu_3 \nu_3 \bar{\phi}_3^0 \phi_3^0$ , as shown in Fig. 1. Assuming that  $\mu_{33}$  dominates among all the  $\mu_{ij}$ 's in Eq. (13) so that




FIG. 1. One-loop radiative breaking of neutrino mass degeneracy.

[9]  $u \simeq -\mu_{33}v_3^2/m_\xi^2$ , the mass matrix  $\mathcal{M}_\nu$  is now corrected to read

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_0 & adIm_0 \\ m_0 & 0 & 0 \\ adIm_0 & 0 & (1+2d^2I)m_3 \end{pmatrix}. \quad (18)$$

The integral  $I$  is given by [7]

$$I = \frac{G_F}{4\pi^2\sqrt{2}\sin^2\beta} \ln \frac{m_\xi^2}{m_W^2}, \quad (19)$$

where  $\sin^2\beta = v_3^2/(v_1^2 + v_3^2)$ . The twofold degeneracy of the  $\nu_1$ - $\nu_2$  sector is then lifted, with the following mass eigenvalues:

$$-m_0 - \frac{a^2d^2I^2m_0^2}{2(m_0+m_3)}, \quad m_0 + \frac{a^2d^2I^2m_0^2}{2(m_0-m_3)}, \quad (20)$$

where  $adI \ll (m_0 - m_3)/(m_0 + m_3)$  has been used, being justified numerically. Hence their mass-squared difference is

$$\Delta m^2 \simeq a^2d^2I^2m_0^3 \left[ \frac{1}{m_0 - m_3} - \frac{1}{m_0 + m_3} \right] \simeq \frac{2a^2d^2I^2m_\nu^4}{m_0^2 - m_3^2}, \quad (21)$$

where  $m_\nu \simeq m_0 \simeq m_3$  has been used. Identifying this with solar neutrino vacuum oscillations then yields

$$\frac{(\Delta m^2)_{sol}(\Delta m^2)_{atm}}{m_\nu^4} = 2a^2d^2I^2 = \frac{2.16 \times 10^{-13}}{\sin^4\beta} \left( \ln \frac{m_\xi^2}{m_W^2} \right)^2, \quad (22)$$

where  $a = 1259$  MeV and  $d = 1250$  MeV from Eq. (17) have been used. In the above, bimaximal mixing (i.e.

$\sin^2 2\theta_{sol} = \sin^2 2\theta_{atm} = 1$ ) has been assumed. For  $(\Delta m^2)_{sol} \sim 4 \times 10^{-10}$  eV<sup>2</sup> in the case of vacuum oscillations and  $(\Delta m^2)_{atm} \sim 4 \times 10^{-3}$  eV<sup>2</sup>, this would require

$$\frac{m_\nu^4}{\sin^4\beta} \left( \ln \frac{m_\xi^2}{m_W^2} \right)^2 \sim 7.4 \text{ eV}^4, \quad (23)$$

which gives the bound  $m_\nu < 0.6$  eV for  $m_\xi > 1$  TeV and  $\sin^2\beta < 0.7$ . It is interesting to note that this same numerical limit in the case of three nearly mass-degenerate neutrinos was recently obtained [12] from the consideration of cosmic structure formation in the light of the latest astronomical observations.

The choice of Eq. (14) in conjunction with Eq. (10) means that neutrinoless double beta decay [8] is absent to lowest order. It also eliminates any one-loop correction to the diagonal entries of  $\mathcal{M}_\nu$  in the  $\nu_1$ - $\nu_2$  sector. This allows the mass splitting to be quadratic (as opposed to linear) in  $I$  as shown in Eq. (20), which is crucial for obtaining the very small phenomenological value of  $(\Delta m^2)_{sol}$  for vacuum oscillations.

#### IV. LEPTON FLAVOR NONCONSERVATION

Both lepton number and lepton flavor are not conserved in this model. Whereas lepton number nonconservation originates from the heavy Higgs triplet  $\xi$  and manifests itself at low energy only through the very small Majorana neutrino masses, lepton flavor nonconservation originates from the much less heavy Higgs doublets which are presumably in the 100 GeV mass range. On the other hand, the  $h_i$ 's of Eq. (11) are suppressed relative to the gauge couplings because they are related to  $\mathcal{M}_l$  as shown in Eqs. (12) and (14).

Using Eqs. (14) and (17), it is easily shown that

$$\begin{pmatrix} l_1 \\ l_2 \\ l_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/\sqrt{2} & -1/\sqrt{2} \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} e \\ \mu \\ \tau \end{pmatrix} \quad (24)$$

and

$$\begin{pmatrix} l_1^c \\ l_2^c \\ l_3^c \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta_R & \sin\theta_R \\ 0 & -\sin\theta_R & \cos\theta_R \end{pmatrix} \begin{pmatrix} e^c \\ \mu^c \\ \tau^c \end{pmatrix}, \quad (25)$$

where

$$\tan\theta_R = \frac{m_\mu}{m_\tau} \left( \frac{m_\tau^2 - m_\mu^2}{m_\tau^2 + m_\mu^2} \right). \quad (26)$$

The interactions of  $\phi_{1,2,3}^0$  are then given by

$$\begin{aligned}
& \phi_1^0 \left[ h_1 e e^c + \frac{-h_3 s_R + h_4 c_R}{\sqrt{2}} \mu \mu^c + \frac{-h_3 c_R + h_4 s_R}{\sqrt{2}} \tau \tau^c + \frac{h_3 c_R + h_4 s_R}{\sqrt{2}} \mu \tau^c + \frac{h_3 s_R + h_4 c_R}{\sqrt{2}} \tau \mu^c \right] \\
& + \phi_2^0 \left[ \frac{h_1 c_R}{\sqrt{2}} \mu \mu^c - \frac{h_1 s_R}{\sqrt{2}} \tau \tau^c - h_3 s_R e \mu^c + \frac{h_4}{\sqrt{2}} \mu e^c + h_3 c_R e \tau^c + \frac{h_4}{\sqrt{2}} \tau e^c + \frac{h_1 s_R}{\sqrt{2}} \mu \tau^c - \frac{h_1 c_R}{\sqrt{2}} \tau \mu^c \right] \\
& + \phi_3^0 \left[ -\frac{h_5 s_R}{\sqrt{2}} \mu \mu^c + \frac{h_5 c_R}{\sqrt{2}} \tau \tau^c + h_1 c_R e \mu^c + \frac{h_1}{\sqrt{2}} \mu e^c + h_1 s_R e \tau^c - \frac{h_1}{\sqrt{2}} \tau e^c + \frac{h_5 c_R}{\sqrt{2}} \mu \tau^c - \frac{h_5 s_R}{\sqrt{2}} \tau \mu^c \right], \quad (27)
\end{aligned}$$

where  $s_R = \sin \theta_R$  and  $c_R = \cos \theta_R$ . In the above,

$$\begin{aligned}
h_1 &= \frac{m_e}{v_1}, \quad h_3 \simeq \frac{m_\tau}{v_1 \sqrt{2}}, \quad h_4 \simeq \frac{\sqrt{2} m_\mu}{v_1}, \\
h_5 &\simeq \frac{m_\tau}{v_3 \sqrt{2}}, \quad s_R \simeq \frac{m_\mu}{m_\tau}, \quad c_R \simeq 1. \quad (28)
\end{aligned}$$

Consequently, the most prominent rare decays are  $\tau^- \rightarrow e^- e^- \mu^+$ ,  $\tau^- \rightarrow e^- \mu^- \mu^+$ , and  $\tau^- \rightarrow \mu^- \mu^- \mu^+$ , with branching fractions

$$B(\tau^- \rightarrow e^- e^- \mu^+) \simeq 2B(\tau^- \rightarrow e^- \mu^- \mu^+) \simeq \frac{m_\mu^2 m_\tau^2}{8 \cos^4 \beta m_{\phi_2^0}^4} \quad (29)$$

and

$$B(\tau^- \rightarrow \mu^- \mu^- \mu^+) \simeq \frac{m_\mu^2 m_\tau^2}{16} \left( \frac{1}{\cos^2 \beta m_{\phi_1^0}^2} - \frac{1}{\sin^2 \beta m_{\phi_3^0}^2} \right)^2. \quad (30)$$

With scalar masses of order 100 GeV, these branching fractions are of order  $10^{-10}$ , much below the present experimental upper limits [13] which are of order  $10^{-6}$ . Note that  $\mu^- \rightarrow e e e$  is suppressed even more strongly in this model because its amplitude is proportional to  $m_e m_\mu$ .

Whereas low-energy tests of this model are limited to neutrino masses and oscillations, dramatic effects are predicted at high energies. The production of  $\phi_{1,2,3}^0$  at future colliders would yield very clear signals from decays such as  $\phi_2^0 \rightarrow \tau^- e^+$  and  $\phi_{1,3}^0 \rightarrow \tau^- \mu^+$ , provided of course that their masses are such that other would-be dominant channels such as  $WW$  and  $ZZ$  do not obscure these signals.

## V. CONCLUDING REMARKS

To understand the present experimental data on atmospheric and solar neutrinos, a model of neutrino and charged-lepton mass matrices based on permutation symmetry  $S_3$  has been proposed. It has a twofold degeneracy in the neutrino mass matrix which is broken radiatively, and allows for a very small mass splitting, suitable for solar neutrino vacuum oscillations, as given by Eq. (22). The  $S_3$  symmetry is maintained in the scalar sector by three Higgs doublets which determine the charged-lepton mass matrix, whereas the neutrinos obtain naturally small Majorana masses from their couplings to a heavy Higgs triplet. Lepton flavor nonconservation at low energy is suppressed by the small charged-lepton masses. This model may be tested at high energy with the production and decay of its scalar doublets.

## ACKNOWLEDGMENTS

This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG03-94ER40837.

---

[1] Y. Fukuda *et al.*, Phys. Lett. B **433**, 9 (1998); **436**, 33 (1998); Phys. Rev. Lett. **81**, 1562 (1998); **82**, 2644 (1999); **82**, 5194 (1999); Phys. Lett. B **467**, 185 (1999).

[2] R. Davis, Prog. Part. Nucl. Phys. **32**, 13 (1994); P. Anselmann *et al.*, Phys. Lett. B **357**, 237 (1995); **361**, 235 (1996); J. N. Abdurashitov *et al.*, *ibid.* **328**, 234 (1994); Phys. Rev. Lett. **83**, 4686 (1999); Y. Fukuda *et al.*, *ibid.* **77**, 1683 (1996); **81**, 1158 (1998); **82**, 1810 (1999); **82**, 2430 (1999).

[3] C. H. Albright, K. S. Babu, and S. M. Barr, Phys. Rev. Lett. **81**, 1167 (1998); V. Barger, S. Pakvasa, T. J. Weiler, and K. Whisnant, Phys. Lett. B **437**, 107 (1998); S. F. King, *ibid.* **439**, 350 (1998); hep-ph/9904210; B. C. Allanach, Phys. Lett. B **450**, 182 (1999); J. K. Elwood, N. Irges, and P. Ramond, Phys. Rev. Lett. **81**, 5064 (1998); R. Barbieri, L. J. Hall, D. Smith, A. Strumia, and N. Weiner, J. High Energy Phys. **12**, 017 (1998); A. J. Baltz, A. S. Goldhaber, and M. Goldhaber, Phys. Rev. Lett. **81**, 5730 (1998); G. Altarelli and F. Feruglio, Phys. Lett. B **439**, 112 (1998); J. High Energy Phys. **811**, 021 (1998); Phys. Lett. B **451**, 388 (1999); M. Jezabek and Y. Sumino, *ibid.* **440**, 327 (1998); **457**, 139 (1999); H. Fritzsch and Z. Xing, *ibid.* **440**, 313 (1998); R. N. Mohapatra and S. Nussinov, *ibid.* **441**, 299 (1998); E. Ma, *ibid.* **442**, 238 (1998); Y. Nomura and T. Yanagida, Phys. Rev. D **59**, 017303 (1999); M. Tanimoto, *ibid.* **59**, 017304 (1999); Phys. Lett. B **456**, 220 (1999); N. Haba, Phys. Rev. D **59**, 035011 (1999); K. Oda, E. Takasugi, E. Tanaka, and M. Yoshimura, *ibid.* **59**, 055001 (1999); J. Ellis, G. K. Leontaris, S. Lola, and D. V. Nanopoulos, Eur. Phys. J. C **9**, 389 (1999); E. Ma, D. P. Roy, and U.

Sarkar, Phys. Lett. B **444**, 391 (1998); S. Davidson and S. F. King, *ibid.* **445**, 191 (1998); R. Barbieri, L. J. Hall, and A. Strumia, *ibid.* **445**, 407 (1999); Y. Grossman, Y. Nir, and Y. Shadmi, J. High Energy Phys. **10**, 007 (1998); S. L. Glashow, P. J. Kernan, and L. M. Krauss, Phys. Lett. B **445**, 412 (1999); A. S. Joshipura and S. D. Rindani, hep-ph/9811252; Phys. Lett. B **463**, 303 (1999); C. Jarlskog, M. Matsuda, S. Skadhauge, and M. Tanimoto, *ibid.* **449**, 240 (1999); E. Malkawi, Phys. Rev. D **61**, 013006 (2000); E. Ma and D. P. Roy, Phys. Rev. D **59**, 097702 (1999); M. Fukugita, M. Tanimoto, and T. Yanagida, *ibid.* **59**, 113016 (1999); L. J. Hall and N. Weiner, *ibid.* **60**, 033005 (1999); Z. Berezhiani and A. Rossi, J. High Energy Phys. **03**, 002 (1999); K. S. Babu, J. C. Pati, and F. Wilczek, hep-ph/9812538; S. Lola and G. G. Ross, Nucl. Phys. **B553**, 81 (1999); Q. Shafi and Z. Tavartkiladze, Phys. Lett. B **451**, 129 (1999); Y. Nomura and T. Sugimoto, hep-ph/9903334; T. Blazek, S. Raby, and K. Tobe, Phys. Rev. D **60**, 113001 (1999); R. Barbieri, P. Creminelli, and A. Romanino, hep-ph/9903460; K. S. Babu, B. Dutta, and R. N. Mohapatra, Phys. Lett. B **458**, 93 (1999); C.-K. Chua, X.-G. He, and W.-Y. P. Hwang, hep-ph/9905340; B. Stech, Phys. Lett. B **465**, 219 (1999); H. B. Benaoum and S. Nasri, Phys. Rev. D **60**, 113003 (1999); C. H. Albright and S. M. Barr, Phys. Lett. B **461**, 218 (1999); P. H. Frampton and S. L. Glashow, *ibid.* **461**, 95 (1999); J. C. Romao, M. A. Diaz, M. Hirsch, W. Porod, and J. W. F. Valle, hep-ph/9907499; G. Altarelli, F. Feruglio, and I. Masina, hep-ph/9907532; A. Abada and M. Losada, hep-ph/9908352.

[4] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978); S. P. Mikheyev and A. Yu. Smirnov, Yad. Fiz. **42**, 1441 (1985) [Sov. J. Nucl. Phys. **42**, 913 (1985)].

[5] D. Caldwell and R. N. Mohapatra, Phys. Rev. D **48**, 3259 (1993); A. S. Joshipura, Z. Phys. C **64**, 31 (1994); Phys. Rev. D **51**, 1321 (1995); P. Bamert and C. P. Burgess, Phys. Lett. B **329**, 289 (1994); D.-G. Lee and R. N. Mohapatra, *ibid.* **329**, 463 (1994); A. Ioannissian and J. W. F. Valle, *ibid.* **332**, 93 (1994); A. Ghosal, *ibid.* **398**, 315 (1997); A. K. Ray and S. Sarkar, Phys. Rev. D **58**, 055010 (1998); C. D. Carone and M. Sher, Phys. Lett. B **420**, 83 (1998); F. Vissani, hep-ph/9708483; H. Georgi and S. L. Glashow, Phys. Rev. D (to be published), hep-ph/9808293; U. Sarkar, Phys. Rev. D **59**, 037302 (1999); R. N. Mohapatra and S. Nussinov, *ibid.* **60**, 013002 (1999); G. C. Branco, M. N. Rebelo, and J. I. Silva-Marcos, Phys. Rev. Lett. **82**, 683 (1999); hep-ph/9906368; C. Wetterich, Phys. Lett. B **451**, 397 (1999); Y. L. Wu, Phys. Rev. D **60**, 073010 (1999); Eur. Phys. J. C **10**, 491 (1999); Int. J. Mod. Phys. A **14**, 4313 (1999); hep-ph/9905222; hep-ph/9906435; R. Barbieri, L. J. Hall, G. L. Kane, and G. G. Ross, hep-ph/9901228; M. Tanimoto, T. Watari, and T. Yanagida, Phys. Lett. B **461**, 345 (1999); A. K. Ray and S. Sarkar, Phys. Rev. D (to be published), hep-ph/9908294.

[6] E. Ma, Phys. Lett. B **456**, 48 (1999); **456**, 201 (1999); J. Phys. G **25**, L97 (1999), hep-ph/9907503; J. Ellis and S. Lola, Phys. Lett. B **458**, 310 (1999); J. A. Casas, J. R. Espinosa, A. Ibarra, and I. Navarro, Nucl. Phys. **B556**, 3 (1999); hep-ph/9905381; J. High Phys. **09**, 015 (1999); R. Barbieri, G. G. Ross, and A. Strumia, *ibid.* **10**, 020 (1999); N. Haba and N. Okamura, hep-ph/9906481; N. Haba, Y. Matsui, N. Okamura, and M. Sugiura, hep-ph/9908429.

[7] E. Ma, Phys. Rev. Lett. **83**, 2514 (1999).

[8] L. Brandis *et al.*, Phys. Rev. Lett. **83**, 41 (1999); H. V. Klein-grothaus, hep-ex/9901021.

[9] E. Ma and U. Sarkar, Phys. Rev. Lett. **80**, 5716 (1998).

[10] E. Ma, Phys. Rev. Lett. **81**, 1171 (1998).

[11] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p. 315; T. Yanagida, in “Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe,” edited by O. Sawada and A. Sugamoto, KEK Report No. 79-18, Tsukuba, Japan, 1979, p. 95; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).

[12] M. Fukugita, G.-C. Liu, and N. Sugiyama, hep-ph/9908450.

[13] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. C **3**, 1 (1998).