

Nacional de Córdoba, Córdoba, Argentina). *Uses.* Medicinal.² *Previous work.* Phytochemical screening.³ Sister species.^{4,5}

Present work. The air-dried above ground plant (without capsules) was extracted exhaustively with 95% EtOH. The extract was concentrated to a syrup *in vacuo*†, dissolved in CH₂Cl₂, extracted with 5% HCl, alkalinized with NH₄OH and extracted with CH₂Cl₂, which upon concentration yielded berberine crystallized as the chloride. After most of the berberine was removed, the alkaloid mixture was then separated by conventional procedures, including fractional crystallization from EtOH (protopine and allocryptopine), and preparative TLC on silica gel P F₂₅₄ using xylene-MeCOEt-MeOH-Et₂NH (20:20:3:1). Bands at *R*_f 0.75 and 0.82 were extracted with CHCl₃-EtOH (3:1) and yielded chelerythrine and sanguinarine respectively. In addition TLC analysis also revealed the presence of traces of at least three other unidentified bases. Yield of crude mixed alkaloids: 0.40% of the dried plant weight. The approximate alkaloid composition (% of alkaloid from the total mixture) was as follows: protopine 41%; allocryptopine 28%; berberine 9%; sanguinarine 5%; chelerythrine 4%.

All the alkaloids were identified by comparing the UV, IR and NMR spectra with those of authentic samples and by m.p., m.m.p., co-chromatography (TLC 3 solvents) and GC.

Acknowledgements—The authors are indebted to Dr. R. Juliani and Ing. A. Hunziker (Universidad Nacional de Córdoba) for the supply of plant material and botanical advice respectively; to Drs. R. H. F. Manske, J. Slavik and D. Giacopello for authentic samples. Thanks also to Dr. M. Vernengo (Instituto Nacional de Farmacología y Bromatología) for the award of a research fellowship to one of us (ALB) and for financial assistance.

² A. L. CABRERA, *Flora de la Provincia de Buenos Aires*, Vol. IV, Part III, p. 266, Colección Científica del I.N.T.A., Buenos Aires (1967).

³ R. V. D. RONDINA, M. E. MENDIONDO and J. D. COUSSIO, *Rev. Invest. Agropec.* Serie 2, **VII**, 271 (1970).

⁴ F. R. STERMITZ, S. M. WORKMAN and W. M. KLEIN, *Phytochem.* **10**, 675 (1971).

⁵ M. H. BENN and R. E. MITCHELL, *Phytochem.* **11**, 461 (1972).

Phytochemistry, 1972, Vol. 11, pp. 3548 to 3549. Pergamon Press. Printed in England.

SOLANACEAE

SOLASODINE GLYCOSIDES AND DIOSGENIN FROM *SOLANUM PINNATUM*

A. URZÚA and B. K. CASSELS

Departamento de Química, Universidad Técnica del Estado, Santiago, Chile

(Received 5 June 1972. Accepted 14 August 1972)

Key Word Index—*Solanum pinnatum*; Solanaceae; solasonine; solamargine.

A sample of a plant growing near Longotoma, Aconcagua, identified as *Solanum pinnatum* Cav. (Voucher specimen No. BC10 deposited in the herbarium of the Museo Nacional de Historia Natural, Santiago), was examined for glycoalkaloids and saponins. The crude basic glycoside mixture isolated from stems and leaves afforded solasonine, solamargine, and a glucose- and rhamnose-containing glycoside of solasodine, probably identical

with a previously described product obtained by partial hydrolysis of solamargine.¹ Hydrolysis of the remaining glycosides furnished two neutral compounds: diosgenin and (25*R*)-spirosta-3,5-diene.²

EXPERIMENTAL

M.p.s of the glycoalkaloids are uncorrected. TLC of the glycosides and aglycones was carried out on Merck silica gel G using *n*-BuOH-(H₂O-satd)-Et₂NH-MeOH (85:10:2) (System I), or C₆H₆-CHCl₃ (9:1) (System II) respectively, spraying with 20% SbCl₃ in CHCl₃. The sugars were examined by TLC on Merck Kieselguhr G prepared with 0.02 M NaOAc, using AcOEt-*i*-PrOH-H₂O (65:23:12) (System III), spraying with anisaldehyde-H₂SO₄.³

Isolation of the alkaloids. Dried, powdered leaves and stems of *S. pinnatum* collected in December 1970 (1.2 kg) were extracted with 5% AcOH, and the basic glycosides were precipitated with NH₄OH. The dried, ground precipitate was extracted with EtOH (Soxhlet), yielding 40 g crude alkaloids (3%). 8 g of the alkaloid mixture were chromatographed on Al₂O₃, eluting with H₂O-satd *n*-BuOH.

SP-1. 0.85 g, *R*_f 0.7 (System I), m.p. 235-240°, [α]_D²³ -90 ± 3° (c 0.62, EtOH). Hydrolysis with 0.1 N HCl yielded solasodine hydrochloride, m.p. 295-300°, *R*_f (System II) and IR spectrum identical with those of an authentic sample. The acid solution contained glucose and rhamnose, identified by co-chromatography (System III). (Found: C, 63.44; H, 8.89; N, 1.92. C₃₉H₆₃NO₁₁·H₂O requires: C, 63.30; H, 8.85; N, 1.89%).

Solamargine. 1.1 g, *R*_f 0.44 (System I), m.p. 274-276°, [α]_D²³ -111 ± 3° (c 0.48, MeOH). Identical with a reference sample (m.m.p., TLC, IR, hydrolysis products).

Solasonine. 1.46 g, *R*_f 0.3 (System I), m.p. 266-268°, [α]_D²³ -90 ± 4° (c 0.55, pyridine). Identical with a reference sample (m.m.p., TLC, IR, hydrolysis products, picrate).

Isolation of the sapogenins. The basic mother liquors of the crude alkaloid precipitate were acidified to 2.5 N with HCl and refluxed (10 hr) to yield a mixture of crude aglycones. These, dried and ground, were extracted with C₆H₆ (Soxhlet), and the soluble material was fractionated on Al₂O₃, eluting with C₆H₆ and C₆H₆-CHCl₃ mixtures.

(25R)-spirosta-3,5-diene. 214 mg, *R*_f 0.72 (System II), m.p. 142-150°, [α]_D²³ -164 ± 5° (c 1.4, CHCl₃), λ_{max} 228, 233, 242 nm (log ε 4.15, 4.18, 4.00). Identical with a reference sample (TLC, IR).

Diosgenin. 550 mg, *R*_f 0.44 (System II), m.p. 206.5-210°, [α]_D²³ -119 ± 3° (c 1.1, CHCl₃). Identical with an authentic sample (m.m.p., TLC, IR, acetate).

Acknowledgements—The authors are indebted to Mrs. M. Muñoz for the botanical determination, Drs. M. Silva, E. Guerreiro and J. Kavka for reference samples, Mrs. A. Bau and Mr. R. Clavijo for technical assistance, and FORGE, New York, for financial help.

¹ R. KUHN and I. LÖW, *Chem. Ber.* **88**, 289 (1955).

² P. BITE and G. MAGYAR, *Acta Chem. Acad. Sci. Hung.* **48**, 255 (1966).

³ E. STAHL and U. KALTENBACH, *J. Chromatog.* **5**, 351 (1961).