



REVISED STRUCTURE FOR HORTENSIN, A FLAVONOID FROM *MILLINGTONIA HORTENSIS*

TAKESHI HASE, KAZUHIRO OHTANI, RYOJI KASAI, KAZUO YAMASAKI* and CHAYAN PICHEANSOONTHON†

Institute of Pharmaceutical Sciences, Hiroshima University, School of Medicine, 1-2-3 Kasumi, Minami-ku, Hiroshima 734, Japan;

†Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand

(Received in revised form 31 January 1995)

Key Word Index—*Millingtonia hortensis*; Bignoniaceae; flavonoid; hortensin; cirsimaritin; pectolinarigenin.

Abstract—The original structure of hortensin, from *Millingtonia hortensis*, and analysed as 4'-hydroxy-6,7-dimethoxyflavonol, has been changed once to 5,4'-dihydroxy-6,7-dimethoxyflavone (cirsimaritin). A further reinterpretation of the NMR data of newly isolated 'hortensin' showed that it is 5,7-dihydroxy-6,4'-dimethoxyflavone (pectolinarigenin). Several related flavonoids from the same plant were also identified.

INTRODUCTION

Hortensin [1], a methoxylated flavone possessing anticancer properties, was isolated from the flower of *Millingtonia hortensis* and given the structure 4'-hydroxy-6,7-dimethoxyflavonol (1). However, re-examination of the published data and comparisons of hortensin with isolated and authentic samples suggested that it was identical with 5,4'-dihydroxy-6,7-dimethoxyflavone (cirsimaritin, 2). In addition, hortensin was different from a sample of synthetic 4'-hydroxy-6,7-dimethoxyflavonol. Thus, the structure of hortensin was revised to cirsimaritin (2) [2].

In the course of our study on the constituents of the flower of *M. hortensis*, we isolated a compound 3, whose melting point and NMR data were the same as those of hortensin [1]. The NMR spectral data of 3, including NOE evidence, revealed that its structure was identical with neither 4'-hydroxy-6,7-dimethoxyflavonol (1) nor 5,4'-dihydroxy-6,7-dimethoxyflavone (cirsimaritin) (2). Instead, 3 was identical with 5,7-dihydroxy-6,4'-dimethoxyflavone (pectolinarigenin, 3) [3, 4]. In this paper, the correction of the structure of 'hortensin' and the identification of four other known flavonoids (4–7) from this plant are presented.

RESULTS AND DISCUSSION

From the hexane-soluble portion of the hot methanolic extract of the dried flowers of *Millingtonia hortensis* (Thai origin), a flavonoid (3) was isolated. The melting point (210–211°, lit. [1]: 212–213°) and the ¹H and ¹³CNMR data were essentially the same (with an addi-


tional ¹H NMR signal at δ13.00) as the reported data of hortensin, which was isolated from the same part of the same plant collected in the same country, and characterized as 4'-hydroxy-6,7-dimethoxyflavonol (1) through its spectroscopic properties, including CSCM 1D and selective INEPT experiments [1] (Table 1).

However, the ¹H and ¹³CNMR data of 3 (Tables 1 and 2) could not satisfy the proposed structure for (1), but were identical with those of 5,7-dihydroxy-6,4'-dimethoxyflavone (pectolinarigenin, 3) [3, 4]. The structure was confirmed with NOE experiments. When the signal for H-2' and H-6' (δ8.03, *d*, *J* = 9 Hz) was irradiated, NOE was observed at H-3 (δ6.87, *s*; 8%) and H-3' and H-5' (δ7.10, *d*, *J* = 9 Hz; 9%). With irradiation on 4'-OMe (δ3.75, *s*) NOE was observed at H-3' and H-5' (5%), while the irradiation of 6-OMe (δ3.86, *s*) resulted in no NOE.

Therefore, the structure of our compound 3, which was previously named hortensin [1], is 5,7-dihydroxy-6,4'-dimethoxyflavone (pectolinarigenin, 3), and the name 'hortensin' should be deleted.

Prior to our correction of the structure of 'hortensin', Nair and Sivakumar [2] claimed the non-identity of hortensin with synthetic 4'-hydroxy-6,7-dimethoxyflavonol. Instead, they identified 'hortensin' as 5,4'-dihydroxy-6,7-dimethoxyflavone (cirsimaritin, 2) based on comparisons of hortensin with their sample, which they isolated from the same plant, and with authentic samples of hortensin from Cordell and co-workers [1] and authentic cirsimaritin from *Flourensia ceruna*. However, the ¹H NMR data of their cirsimaritin [2] were neither identical with those of original hortensin [1], nor with those of cirsimaritin [5] recorded in the same solvent (Table 1). In addition, the melting point of their cirsimaritin was 258–260°, significantly different from that of hortensin

*Author to whom correspondence should be addressed.

Table 1. ¹H NMR data of 'hortensin' and related compounds (in DMSO-*d*₆)

C	Original hortensin [1]*	This work 3	Revised 'hortensin' [2]†	Cirsimaritin 2 [5]	Pectolinarigenin 3 [3]
3		6.87 s	6.54 s	6.87 s	6.86 s
5	6.88 s				
8	6.62 s	6.62 s	7.34 s	6.75 s	6.63 s
2'/6'	8.04 d, 9.0	8.03 d, 9.0	7.82 d, 9.0	7.98 d, 9.0	8.03 d
3'/5'	7.11 d, 9.0	7.10 d, 9.0	7.01 d, 9.0	6.96 d, 9.0	7.12 d
6-OMe	3.86 s	3.86 s	3.90 s	3.79 s	3.86 s
7-OMe	3.76 s		3.95 s	3.96 s	
5-OH		13.00 s	12.95 s	12.9 s	13.00 s
4'-OH			9.95 s		
4'-OMe		3.75 s			3.77 s

*The structure (1) was proposed.

†The structure (2) was proposed.

(212–213°). They did not measure the ¹³C NMR of their sample, but reassigned the reported data of hortensin [1] for cirsimaritin (Table 2).

Apart from 3, four structurally related flavonoids (4–7) were isolated from the same plant, along with 21 phenylethanoids and related compounds [6]. Compounds 4–7 were identified as hispidulin [7, 8], apigenin 7-*O*-glucuronide [9], hispidulin 7-*O*-glucoside [10] and hispidulin 7-*O*-glucuronide methyl ester [11], respectively, by means of their NMR data. The NMR data of 4 also confirmed the structure of 3.

EXPERIMENTAL

Mps: uncorr. ¹H and ¹³C NMR (TMS as int. standard): 400 and 100 MHz, respectively, in DMSO-*d*₆.

Plant material. *Millingtonia hortensis* was collected at the suburb of Khon Kaen City, Thailand. A voucher specimen is deposited at the Herbarium of Khon Kaen University, Thailand.

Extraction and isolation. Dried flower (300 g) was extracted with MeOH at room temp. to give 149 g of extract, a part (51 g) of which was suspended in H₂O and

Table 2. ^{13}C NMR data of 'hortensin', in descending delta values and their assignments (in DMSO- d_6)

Original hortensin [1]*	C	Revised hortensin [2]†	This work 3	C	Pectolinarigenin‡ 3[4]	C
182.1	4	4	182.1	4	181.3	4
163.2	6	2	163.3	2	163.0	2
162.2	7	4'	162.3	4'	162.0	4'
157.2	4'	7	157.4	7	157.0	7
152.7	2	5	152.7	5	153.4	5
152.4	9	9	152.4	9	152.1	9
131.3	3	6	131.4	6	131.2	6
128.2	2'/6'	2'/6'	128.3	2'/6'	127.9	2'/6'
122.8	1'	1'	122.9	1'	122.7	1'
114.5	3'/5'	3'/5'	114.5	3'/5'	114.3	3'/5'
104.1	10	10	104.1	10	103.9	10
103.0	5	3	103.0	3	102.8	3
94.2	8	8	94.3	8	94.0	8
59.9	6-OMe	6-OMe	59.9	6-OMe	59.6	6-OMe
55.5	7-OMe	7-OMe	55.5	4'-OMe	55.3	4'-OMe

*The structure (1) was proposed.

†Quoted from the data of lit. [1], structure 2 was proposed.

‡Measured at 50°.

extracted with hexane and EtOAc, successively. The hexane-soluble portion was concd and chromatographed on a column of silica gel eluted with CH_2Cl_2 -MeOH, and on Sephadex LH-20 eluting with MeOH to afford 3 (33 mg) and 4 (120 mg). The first MeOH extract was chromatographed on Diaion HP-20 eluting with H_2O , 40% MeOH, 80% MeOH, MeOH and Me_2CO . From the 40% MeOH fr., compound 5 (360 mg) was isolated with silica gel and MPLC column (ODS). The 80% eluate was mixed with H_2O and extracted with EtOAc. The EtOAc extract was chromatographed on silica gel (solvent: CHCl_3 -MeOH- H_2O) followed by HPLC (Amide 80: 21.5 mm ϕ $\times$ 300 mm) using MeOH- H_2O and/or MeCN- H_2O system at a flow rate of 6 ml min^{-1} to afford 6 (36 mg) and 7 (21 mg).

Pectolinarigenin (3). Crystals from MeOH, mp 210–211°, ^{13}C and ^1H NMR: see Table 1.

Hispidulin (4). Powder, ^{13}C NMR [7]: δ (from C-2 to C-10): 163.8, 102.4, 182.1, 152.8, 131.4, 157.3, 94.2, 152.4, 104.0; (from C-1' to C-4'): 121.2, 128.4, 115.9, 161.2, 6-OMe: 59.9; ^1H NMR [8]: δ 13.05 (1H, s, 5-OH), 7.90 (2H, d, J = 8.8 Hz, H-2', H-6'), 6.90 (2H, d, J = 8.8 Hz, H-3', H-5'), 6.75 (1H, s, H-3), 6.57 (1H, s, H-8), 3.73 (3H, s, 6-OMe).

Apigenin 7-O-glucoronide (5) [9]. Oil, $[\alpha]_D^{17}$ – 30° (MeOH; c 0.47). ^{13}C NMR: δ (from C-2 to C-10): 164.3, 103.2, 182.0, 161.4, 99.4, 162.5, 94.7, 156.9, 105.4; (from C-1' to C-4'): 121.0, 128.6, 116.0, 161.2, (from GlcA-1 to GlcA-6): 99.2, 72.7, 75.4, 71.2, 75.6, 170.0; ^1H NMR: δ 12.98 (1H, br s, 5-OH), 10.42 (1H, br s, 6''-COOH), 7.96 (2H, d, J = 8.8 Hz, H-2' H-6'), 6.94 (2H, d, J = 8.8 Hz, H-3' H-5'), 6.87 (1H, s, H-3), 6.86 (1H, d, J = 2.2 Hz, H-8), 6.47 (1H, d, J = 2.2 Hz, H-6), 5.27 (1H, d, J = 7.1 Hz, GlcA-1), 4.05 (1H, d, J = 9.5 Hz, GlcA-5).

Hispidulin 7-O-glucoside (6) [10]. Oil, $[\alpha]_D^{17}$ – 66° (MeOH; c 0.67). ^{13}C NMR: δ (from C-2 to C-10): 164.2, 102.5, 182.2, 152.3, 132.4, 156.3, 94.3, 152.0, 105.6; (from C-1' to C-4'): 120.9, 128.5, 115.9, 161.3; 6-OMe: 60.2; (from Glc-1 to Glc-6): 100.1, 73.1, 77.1, 69.5, 76.6, 60.5; ^1H NMR: δ 12.95 (1H, br s, 5-OH), 7.96 (2H, d, J = 8.9 Hz, H-2', H-6'), 7.02 (1H, s, H-8), 6.95 (1H, d, J = 8.9 Hz, H-3' H-5'), 6.85 (1H, s, H-3), 5.12 (1H, d, J = 7.4 Hz, Glc-1), 3.78 (3H, s, 6-OMe).

Hispidulin 7-O-glucuronide methyl ester (7) [11]. Powder, $[\alpha]_D^{17}$ – 53° (MeOH; c 0.67). ^{13}C NMR: δ (from C-2 to C-10): 164.3, 102.7, 182.2, 152.6, 132.6, 155.8, 93.9, 152.0, 105.9; (from C-1' to C-4'): 121.0, 128.5, 116.0, 161.3, 6-OMe: 60.2; (from GlcA-1 to GlcA-6): 99.5, 72.8, 75.6, 71.2, 75.3, 169.0; COOMe: 51.9; ^1H NMR: δ 7.95 (2H, d, J = 8.8 Hz, H-2', H-6'), 7.06 (1H, s, H-8), 6.95 (1H, d, J = 8.8 Hz, H-3', H-5'), 6.85 (1H, s, H-3), 5.36 (1H, d, J = 7.4 Hz, GlcA-1), 3.77 (3H, s, 6-OMe), 3.67 (3H, s, COOMe).

Acknowledgements—We thank the Japanese Society of the Promotion of Science (JSPS) and the National Research Council of Thailand (NRCT) for the promotion of international collaborative research. Our thanks are also due to the Research Centre for Molecular Medicine (RCMM), Hiroshima University School of Medicine, for the use of NMR equipment.

REFERENCES

- Bunyapraphatsara, N., Blasko, G. and Cordell, G. A. (1989) *Phytochemistry* **28**, 1555.
- Nair, A. G. R. and Sivakumar, R. (1992) *Phytochemistry* **31**, 671.

3. Imre, S., Oztung, A. and Wagner, H. (1977) *Phytochemistry* **16**, 799.
4. Markham, K. R., Chari, V. M. and Mabry, T. J. (1982) in *The Flavonoids: Advances in Research* (Harbone, J. B. and Mabry, T. J., eds). pp. 52. Chapman and Hall, London.
5. Mesquita, A. A. L., Corea, D. D. D., Padua, A. P. D., Guedes, M. L. O. and Gottlieb, O. R. (1986) *Phytochemistry* **25**, 1255.
6. Hase, T., Ohtani, K., Kasai, R., Yamasaki, K. and Picheansoonth, C. (1995) *Phytochemistry* **39**, 235
7. Collado, I. G., Macias, F. A., Massanet, G. M. and Luis, F. R. (1985) *J. Nat. Prod.* **48**, 819.
8. Annlakanapakorn, K., Bunyarpraphatsara, N. and Satayavivad, J. (1987) *J. Sci. Soc. Thailand* **13**, 71.
9. Ogawa, K. (1990) Master Thesis submitted to Hiroshima University School of Medicine.
10. Fernandez, I., Garcia, B., Pedro, J. R. and Varea, A. (1991) *Phytochemistry* **30**, 1030.
11. Subramanian, S. S. and Nair, A. G. R. (1972) *Phytochemistry* **11**, 464.