



Erratum

Erratum to “Distribution of ginkgolides and terpenoid biosynthetic activity in *Ginkgo biloba*”
[*Phytochemistry*, 48 (1998) 89–92][☆]

D.J. Carrier^{a,*}, T. A. van Beek^b, R. van der Heijden^c, R. Verpoorte^c

^a Department of Agricultural and Bioresource Engineering, University of Saskatchewan, 57 Campus Drive, Saskatoon, SK, Canada S7N 5A9

^b Phytochemical Section, Laboratory of Organic Chemistry, Agricultural University, Dreijenplein 8, 6703 HB Wageningen, The Netherlands

^c Division of Pharmacognosy, Leiden/Amsterdam Center for Drug Research, Leiden University, P.O. Box 9502, 2300 RA Leiden, The Netherlands

The correct version of Table 1 is presented.

Table 1

Bilobalide, G-A, G-B, G-C and G-J and total terpene content with standard deviation of various parts obtained from 3-year-old *G. biloba* plants. Contents are expressed as mg terpene 100 mg⁻¹ dry wt

Plant part	Bilobalide	G-A	G-B	G-C	G-J	Total
Terminal	0.22 ± 0.04	0.18 ± 0.01	0.09 ± 0.02	0.09 ± 0.02	0.03 ± 0.00	0.61 ± 0.09
Side branch	0.16 ± 0.06	0.13 ± 0.01	0.06 ± 0.00	0.07 ± 0.01	0.01 ± 0.01	0.43 ± 0.09
Rosette	0.27 ± 0.12	0.19 ± 0.04	0.08 ± 0.01	0.08 ± 0.00	0.02 ± 0.01	0.64 ± 0.18
Root meristem	0.00 ± 0.00	0.13 ± 0.01	0.05 ± 0.01	0.06 ± 0.01	0.03 ± 0.01	0.27 ± 0.04
Root	0.01 ± 0.02	0.13 ± 0.07	0.05 ± 0.02	0.05 ± 0.02	0.02 ± 0.01	0.26 ± 0.14
Root bark	0.00 ± 0.00	0.02 ± 0.02	0.01 ± 0.01	0.02 ± 0.00	0.01 ± 0.01	0.06 ± 0.04
Stem bark	0.01 ± 0.00	0.01 ± 0.01	0.00 ± 0.00	0.02 ± 0.00	0.01 ± 0.01	0.05 ± 0.02

[☆] PII of original article: S0031-9422(97)00450-0

* Corresponding author.