

Phenolic compounds from the flowers of *Garcinia dulcis*

S. Deachathai ^a, W. Mahabusarakam ^{a,*}, S. Phongpaichit ^b, W.C. Taylor ^c,
Y.-J. Zhang ^d, C.-R. Yang ^d

^a Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand

^b Department of Microbiology, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand

^c School of Chemistry, University of Sydney, NSW 2006, Australia

^d Laboratory of Phytochemistry, Kunming Institute of Botany, Chinese Academy of Sciences, China

Received 7 July 2005; received in revised form 14 September 2005

Available online 1 December 2005

Abstract

Dulcisxanthones C–F (**1**–**4**) and dulcinone (**5**) together with 22 known compounds were isolated from the flowers of *Garcinia dulcis*. Their structures were determined by spectroscopic methods. The abilities of some of these compounds to act as radical scavengers and antibacterial agents were investigated.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: *Garcinia dulcis*; Guttiferae; Xanthones; Chromones; Radical scavenging; Antibacterial

1. Introduction

The sub-woody plant *Garcinia dulcis* Kurz. (Guttiferae), local Thai name Ma-Phut, grows mainly in southeast Asia. Its leaves and seeds have been used in traditional medicine against lymphatitis, parotitis, struma and other disease conditions (Kasahara and Henmi, 1986). In our previous work (Deachathai et al., 2005), we have investigated the chemical constituents from its fruit and their biological activities. In our continuing work, we have examined the chemical constituents from the flowers. This investigation had led to the isolation and structural determination of five new and 22 known compounds.

2. Results and discussion


The flowers of *G. dulcis* were sequentially extracted with acetone. Purification of the extract, produced four new

xanthones named dulcisxanthones C–F (**1**–**4**), one new chromone named dulcinone (**5**), along with 22 known compounds: volkensiflavone (**6**), morelloflavone (**7**) (Ansari et al., 1976), 1-hydroxy-3,4,5-trimethoxyxanthone (**8**) (Stout et al., 1973), rhamnazin (**9**) (Subhadhirasakul et al., 2003), quercetin 3-*O*- β -galactopyranoside (**10**) (Kartnig et al., 1985), podocarpusflavone A (**11**) (Harrison et al., 1994), xanthochymusside (**12**), fukugeside (**13**) (Kono-shima et al., 1970), cowaxanthone (**14**) (Deachathai et al., 2005), GB-2a (**15**) (Ansari et al., 1976), xanthochymol (**16**) (Blount and Williams, 1976), BR-xanthone A (**17**), 1,3,6-trihydroxy-7-methoxy-2,5-bis(3-methyl-2-butenyl)-xanthone (**18**) (Deachathai et al., 2005), guttiferone E (**19**) (Gustafson et al., 1992), rheediaxanthone A (**20**) (Delle Monache et al., 1981), α -mangostin (**21**), 1,7-dihydroxy-3-methoxy-2-(3-methyl-2-butenyl)xanthone (**22**) (Deachathai et al., 2005), 3-isomangostin (**23**) (Mahabusarakam et al., 1987), kaempferol 3-*O*-glucopyranosyl-7-*O*-rhamnopyranoside (**24**), garcinone B (**25**), morusignin J (**26**) (Deachathai et al., 2005) and β -mangostin (**27**) (Mahabusarakam et al., 1987).

Dulcisxanthone C (**1**) is 1-hydroxy-2,3,4,6-tetramethoxyxanthone, and a yellow solid, m.p. 125–128 °C. Its

* Corresponding author. Tel.: +66 7428 8432; fax: +66 7421 2918.

E-mail addresses: mwilawan@ratree.psu.ac.th, wilawan.m@psu.ac.th (W. Mahabusarakam).



molecular formula of $C_{17}H_{16}O_7$ was established on the basis of its mass spectrum ($[M]^+$ m/z 332). The UV spectrum showed absorption maxima at 376, 309, 273, 241 nm. The IR spectrum exhibited O–H stretching at 3400 cm^{-1} and C=O stretching at 1646 cm^{-1} . The ^1H NMR spectrum showed a singlet signal of a chelated hydroxy proton (1-OH) at δ 12.60. The signals in the aromatic region, δ 7.84 (*dd*), 7.32 (*d*) and 7.26 (*d*) that appeared as an ABX type were proposed for the signals of H-7, H-8 and H-5, respectively. These assignment were supported by the correlations of H-7 to C-5, C-6, C-9; H-8 to C-6, C-8a, C-10a and H-5 to C-6, C-7, C-10a from an HMBC experiment. Four singlet signals at δ 4.04, 4.16, 3.97 and 4.05 were assigned for the methoxy protons of 2-OCH₃, 3-OCH₃, 4-OCH₃ and 6-OCH₃, respectively, and confirmed by the 3J coupling of the methoxy protons to C-2, C-3, C-4 and C-6, respectively, in HMBC. The assigned structure was further confirmed by analysis of HMBC correlations (Table 1).

Dulcisxanthone D (2) is 1,6-dihydroxy-5-(3-methyl-2-butenyl)-2',2'-dimethylchromeno(5',6':2,3)-2''',2'''-dimethylchromeno(5''',6''':8,7)xanthone. It is an orange solid, m.p. 218–220 °C, with the molecular formula $C_{28}H_{28}O_6$. The UV spectrum showed absorption maxima at 335, 301, 289, 221 nm. Absorption bands of O–H stretching and C=O stretching were shown in the IR spectrum at 3424 cm^{-1} and at 1615 cm^{-1} . The ^1H NMR spectrum showed the signals of a chelated hydroxy proton at δ 13.70 (1-OH), a non-chelated hydroxy proton at δ 6.34 (6-OH) and an isolated aromatic proton at δ 6.30 (H-4). The characteristic signals of a prenyl group were present at δ 3.57 (H-1''), 5.28 (H-2''), 1.69 (H-4'') and 1.88 (H-5'')

Table 1

HMBC correlation data of compounds 1 and 5

H-position	1	5
3		C-2, C-4, C-4a, 2-Me
5	C-6, C-7, C-10a	C-4, C-4a, C-6, C-7, C-8a
7	C-5, C-6, C-9	
8	C-6, C-8a, C-10a	
2-OMe	C-2	
3-OMe	C-3	
4-OMe	C-4	
6-OMe	C-6	
2-Me		C-2, C-3
7-Me		C-6, C-7, C-8

and it was located at C-5 according to the HMBC correlation of H-1'' to C-6, and C-10a. The signals of four methyl groups at δ 1.49 (6H, *s*, 2'''-Me) and 1.47 (6H, *s*, 2'-Me) and vicinal olefinic protons at δ 5.77 (*d*, H-3'''), 7.98 (*d*, H-4'''), 5.57 (*d*, H-3') and 6.73 (*d*, H-4') associated with a chromene rings were present. The 3J correlations of H-4''' to C-7 and H-4' to C-1 suggested that the chromene rings were connected to the parent structure at C-7, C-8 and C-2, C-3. The complete HMBC data (Table 2) confirmed this structure.

Dulcisxanthone E (3) is 1,3,6,7-tetrahydroxy-2-(3,7-dimethyl-2,6-octadienyl)-5-(3-methyl-2-butenyl)xanthone. It is a yellow solid with the molecular formula $C_{28}H_{32}O_6$. The UV spectrum showed absorption maxima at 369, 322, 258, 242 nm. The IR spectrum showed absorption bands of O–H stretching at 3450 cm^{-1} and C=O stretching at 1690 cm^{-1} . The ^1H NMR spectrum exhibited resonances of a chelated hydroxy proton 1-OH at δ 13.19 and aromatic protons H-4 at δ 6.42 and H-8 at δ 7.33. The characteristic

Table 2
HMBC correlation data of compounds **2–4**

H-position	2	3	4
4	C-2, C-3, C-4a, C-9, C-4'	C-2, C-3, C-4a, C-9a	C-2, C-3, C-4a, C-9a
5			C-6, C-7, C-8a, C-10a
8		C-6, C-7, C-8a, C-9, C-10a	
1'		C-1, C-2, C-3, C-2', C-3'	C-1, C-2, C-3, C-2', C-3'
2'		C-1', C-4'	C-2, C-4', C-5'
3'	C-2, C-2', 2'-Me		
4'	C-1, C-2'	C-3', C-5', C-6'	C-2', C-3', C-5'
5'		C-3', C-4', C-6'	C-2', C-3', C-4'
6'		C-5', C-8', C-10'	
8'		C-6', C-7', C-10'	
9'		C-2', C-3', C-4', C-5'	
10'		C-6', C-7', C-8'	
1''	C-6, C-8a, C-10a, C-2'', C-3''	C-5, C-6, C-10a, C-2'', C-3''	
2''	C-5, C-4'', C-5''	C-1'', C-4'', C-5''	
3''			C-8, C-4''
4''	C-2'', C-3'', C-5''	C-2'', C-3'', C-5''	C-2''
5''	C-2'', C-3'', C-4''	C-2'', C-3'', C-4''	
3'''	C-8, C-2'''		
4'''	C-7, C-2'''		
1-OH	C-1, C-2, C-9a		C-9a, C-1, C-2
3-OMe			C-3
2'-Me	C-2', C-3'		
2''-Me			C-3''
2'''-Me	C-2''', C-3'''		

signals of a prenyl unit were displayed at δ 6.30 (*d*, H-1''), 5.28 (*br t*, H-2''), 1.69 (*s*, H-4'') and 1.88 (*s*, H-5''). The prenyl unit was linked at C-5 according to the HMBC correlation of H-1'' to C-5, C-6 and C-10a. It also exhibited the typical signal of a geranyl group: three singlets of three methyl groups at δ 1.57 (H-8'), 1.80 (H-9') and 1.52 (H-10'), a doublet of methylene protons at δ 3.37 (H-1'), two multiplets of methylene protons at δ 1.99 (H-4') and 2.03 (H-5') and two broad triplets of two olefinic methine protons at δ 5.28 (H-2') and 5.02 (H-6'). The correlation of H-1' to C-1, C-2 and C-3 in the HMBC indicated that the geranyl side chain was at C-2. The complete HMBC (Table 2) supported this structure.

Dulcisxanthone **F** (**4**) is 1,6-dihydroxy-2-(3-methyl-2-butenyl)-3-methoxy-2'',2''-dimethylchromeno(5'',6'':8,7)-xanthone. It is a yellow solid with the molecular formula $C_{24}H_{24}O_6$. The UV spectrum showed absorption maxima at 331, 322, 265, 244 nm. The IR spectrum exhibited absorption bands of O–H stretching at 3479 cm^{-1} and C=O stretching at 1675 cm^{-1} . As was found for compound **2**, the ^1H NMR spectrum indicated the presence of a chelated hydroxy group 1-OH at δ 13.35 (*s*), aromatic protons H-4 at δ 6.36 (*s*), H-5 at δ 6.83 (*s*), methoxy protons 3-OCH₃ at δ 3.91 (*s*) and proton resonances corresponded to a prenyl group at δ 1.80 (H-5', *s*), 1.68 (H-4', *s*), 3.36 (H-1', *d*) and 5.23 (H-2', *br t*). The characteristic signals of two methyl groups (H-5'', H-6'') and vicinal olefinic protons (H-3'' and H-4'') associated with a chromene ring were shown at δ 1.50, 5.83 and 8.04, respectively. The deshielded effect on the resonance of H-4'' indicated that the chromene ring was attached to the xanthone nucleus close to its carbonyl group. The HMBC correlation of H-1' to C-1, C-2

and C-3 indicated that the prenyl unit was at C-2. The methoxy group was assigned at C-3 according to the HMBC correlations of H-4 to C-3 and methoxy protons to C-3. Irradiation of the methoxy protons at δ 3.91 in a NOE experiment, enhanced the signal of H-4 and this also confirmed the placement of the methoxy group. The complete HMBC (Table 2) supported the assigned structure.

Dulcinone (**5**) is 6,8-dihydroxy-2,7-dimethyl-4*H*-chromen-4-one. It is a yellow solid with molecular formula $C_{11}H_{10}O_4$. The IR spectrum showed absorption bands of O–H stretching at 3343 cm^{-1} and C=O stretching at 1657 cm^{-1} . The ^{13}C NMR spectrum also showed the resonance of a carbonyl carbon at δ 181.5. The ^1H NMR spectrum showed a singlet signal of an olefinic proton H-3 at δ 5.99, a singlet signal of an aromatic proton H-5 at δ 6.33, a singlet signal of methyl protons 2-Me at δ 2.39 and a singlet signal of methyl protons 7-Me at δ 2.13. These assignment were supported by the HMBC correlations: H-3 to C-2, C-4, C-4a; H-5 to C-4a, C-6, C-7, C-8a; 2-Me to C-2, C-3 and 7-Me to C-6, C-7, C-8. The assigned structure was further confirmed by HMBC correlations (Table 1).

Compounds **1–5** are new natural products. Compounds **6**, **7**, **11**, **14**, **15**, **17**, **18**, **21**, **22**, **24**, **25**, and **26** were previously isolated from *G. dulcis* (Ansari et al., 1976; Harrison et al., 1994; Deachathai et al., 2005). This is the first report of the known compounds **8**, **9**, **10**, **12**, **13**, **16**, **19**, **20**, **23**, **27** in *G. dulcis*. Evaluation of the radical scavenging activities of some of the compounds at the concentration of $10\text{ }\mu\text{M}$ (Table 3) revealed that none of the newly isolated compounds showed any antioxidant activity. However **10**, **13**, **16** and **19** had potent antioxidant properties produced 57%, 56%, 60% and 59% scavenging properties, respec-

Table 3
Radical scavenging activity of compounds from the flower of *G. dulcis* (10 μ M)

Compounds	% Scavenging of DPPH	IC ₅₀ (μ M)
1	2	—
2	16	—
3	15	—
4	2	—
5	3	—
6	5	—
8	2	—
9	8	—
10	57	10.50
11	5	—
12	36	—
13	56	11.40
15	33	—
16	61	8.50
19	59	10.10
20	8	—
23	3	—
27	2	—
BHT	43	19.00

tively. These values corresponded to IC₅₀ values of 10.50, 11.40, 8.50 and 10.10 μ M, respectively. Compound **9** and **16** were tested for the antibacterial activity. Compound **16** was found to inhibit the growth of both the penicillin-sensitive strain, ATCC 25923, and the methicillin-resistant strain, MRSA SK1, of *Staphylococcus aureus* with MIC values of 8 μ g/mL whereas **9** showed only weak activity with MIC values greater than 128 μ g/mL. The results of radical scavenging and antibacterial activities of compounds **7**, **14**, **17**, **18**, **21**, **22**, **24**, **25**, **26** have been reported in the previous work (Deachathai et al., 2005).

3. Experimental

3.1. General methods

Melting points were measured on a digital Electrothermal 9100 Melting Point Apparatus and are uncorrected. Infrared spectra were recorded on an FTS 165 FT-IR spectrometer. Ultraviolet absorption spectra were recorded using a UV-160A spectrometer (SHIMADZU). ¹H and ¹³C NMR spectra were performed on a Varian UNITY INOVA 500 spectrometer and on a 300 MHz FTNMR, Brüker spectrometer.

The high resolution mass spectra were recorded on an MS25RFA spectrometer. Pre-coated TLC sheets of silica gel 60 PF₂₅₄ were used. Quick column chromatography (QCC) was performed with silica gel 60H. Column chromatography (CC) was performed with silica gel 100 and sephadex LH-20.

3.2. Plant material

The flowers of *G. dulcis* were collected from Songkhla province in the southern part of Thailand. The voucher

specimen (Coll. No. 02, Herbarium No. 0012652) has been deposited at Prince of Songkla University Herbarium, Biology Department, Faculty of Science, Prince of Songkla University, Thailand.

3.3. Extraction and isolation

The flowers of *G. dulcis* (1.2 kg) were extracted at room temperature sequentially with acetone (5 and 7 days). Removal of the solvents from the extracts yielded the acetone extracts A (148.2 g) and B (38.4 g). An aliquot of the acetone extract A (92.0 g) was further separated by QCC and eluted with CH₂Cl₂, CH₂Cl₂–Me₂CO, Me₂CO, Me₂CO–MeOH gradient solvent system. The eluted fractions were combined into 17 fractions (A1–A17) on the basis of their TLC behaviour. Fraction A6 (7.7 g) was subjected to CC on sephadex LH-20 and eluted with a gradient of H₂O–MeOH to give compounds **1** (3.0 mg), **8** (9.0 mg), **9** (62.0 mg), **10** (27.0 mg) and **11** (10.0 mg). Fractions A5 (2.2 g), A7 (4.8 g) and A13 (1.0 g) were further purified producing **6** (29.1 mg), **7** (1.5 g), **12** (148.0 mg) and **13** (51.0 mg). The acetone extract A (55.8 g) was also fractionated by dissolving in CH₂Cl₂ to give a soluble (15.2 g) and insoluble (40.2 g) fraction. This CH₂Cl₂ soluble fraction was subjected to silica gel CC and eluted sequentially with CH₂Cl₂, and a CH₂Cl₂–Me₂CO gradient to afford **2** (10.0 mg), **14** (21.5 mg), **15** (38.2 mg) and **16** (2.3 g). The CH₂Cl₂ insoluble fraction after CC produced **17** (15.2 mg) and **18** (5.4 mg). Extraction of the acetone extract B (38.4 g) with CH₂Cl₂, EtOAc and then Me₂CO gave CH₂Cl₂ soluble – (8.7 g), EtOAc soluble – (8.5 g) and Me₂CO soluble (3.5 g) fractions. After CC of CH₂Cl₂ soluble fraction and eluting with CH₂Cl₂–Me₂CO in a polarity gradient manner, compounds **19** (10.2 mg), **20** (5.5 mg), **3** (14.3 mg), **21** (31.7 mg), **22** (20.2 mg), **23** (3.2 mg) and **24** (12.3 mg) were obtained. The EtOAc soluble fraction was subjected to silica gel CC eluted with CH₂Cl₂–Me₂CO in a polarity gradient manner to afford **4** (23.8 mg) and **5** (5.2 mg), whereas the Me₂CO soluble fraction subjected to silica gel CC and eluted with CH₂Cl₂–MeOH in a polarity gradient manner gave **25** (2.6 mg), **26** (2.3 mg) and **27** (15.2 mg).

3.3.1. Dulcisxanthone C (**1**)

Yellow solid, m.p. 125–128 °C. HRESIMS *m/z* 332.0892 [M]⁺ (calcd. for C₁₇H₁₆O₇, 332.0896). UV (CH₃OH) $\lambda_{\max}$ (nm) (log ϵ): 376 (3.37), 309 (3.78), 273 (3.87), 241 (3.91). IR (neat), ν (cm^{−1}): 3400, 1646. ¹H NMR (CDCl₃) (δ ppm): 12.60 (1H, *s*, 1-OH), 7.84 (1H, *dd*, *J* = 7.8, 1.5 Hz, H-7), 7.32 (1H, *d*, *J* = 7.8 Hz, H-8), 7.26 (1H, *d*, *J* = 1.5 Hz, H-5), 4.16 (3H, *s*, 3-OMe), 4.05 (3H, *s*, 6-OMe), 4.04 (3H, *s*, 2-OMe), 3.97 (3H, *s*, 4-OMe). EIMS *m/z* (% rel. int): ([M]⁺ 332, 100), 317 (98), 302 (21), 287 (15), 259 (17), 203 (10), 175 (13). ¹³C NMR (CDCl₃) (δ ppm): 181.7 (C-9), 154.1 (C-3), 150.6 (C-1), 148.7 (C-6), 146.4 (C-10a), 145.7 (C-4a), 135.4 (C-4), 132.7 (C-2), 123.7 (C-8), 120.9 (C-8a), 116.5 (C-7), 115.9 (C-5), 104.5

(C-9a), 61.9 (2-OMe), 61.7 (3-OMe), 61.2 (4-OMe), 56.5 (6-OMe).

3.3.2. *Dulcisxanthone D* (2)

Yellow solid, m.p. 218–220 °C. HRFABMS m/z 460.1929 $[M]^+$ (calcd. for $C_{28}H_{28}O_6$ 460.1886). UV (CH₃OH) λ_{max} (nm) (log ϵ): 335 (4.10), 301 (4.24), 289 (4.30), 221 (4.21), 205 (4.25). IR (neat) ν (cm⁻¹): 3424, 1615, 1440 cm⁻¹. ¹H NMR (CDCl₃) (δ ppm): 13.70 (1H, s, 1-OH), 7.98 (1H, d, J = 10.2 Hz, H-4''), 6.73 (1H, d, J = 10.2 Hz, H-4'), 6.34 (1H, s, 6-OH), 6.30 (1H, s, H-4), 5.77 (1H, d, J = 10.2 Hz, H-3'''), 5.57 (1H, d, J = 10.2 Hz, H-3'), 5.28 (1H, br t, J = 5.7 Hz, H-2''), 3.57 (2H, d, J = 7.5 Hz, H-1''), 1.88 (3H, s, H-5''), 1.69 (3H, s, H-4''), 1.49 (6H, s, 2'''-Me), 1.47 (6H, s, 2'-Me). FABMS m/z (% rel. int.): ($[M]^+$ 460, 100), 444 (50), 61 (42). ¹³C NMR (CDCl₃) (δ ppm): 182.8 (C-9), 159.8 (C-1), 157.7 (C-3), 156.5 (C-10a), 150.9 (C-4a), 148.6 (C-6), 136.5 (C-7), 132.6 (C-3''), 131.3 (C-3'''), 127.0 (C-3'), 120.9 (C-4'''), 120.9 (C-2''), 117.1 (C-8), 115.7 (C-4'), 115.3 (C-8a), 108.3 (C-5), 104.2 (C-9a), 103.8 (C-2), 94.2 (C-4), 77.9 (C-2'), 76.8 (C-2'''), 28.3 (2'-Me), 27.4 (2'''-Me), 25.8 (C-4''), 22.5 (C-1''), 18.0 (C-5'').

3.3.3. *Dulcisxanthone E* (3)

Yellow solid, m.p. 185–187 °C. HRESIMS m/z 464.2199 $[M]^+$ (calcd. for $C_{28}H_{32}O_6$ 464.2199). UV (CH₃OH) λ_{max} (nm) (log ϵ): 369 (4.06), 322 (4.28), 258 (4.46), 242 (4.50). IR (neat) ν (cm⁻¹): 3450, 2923, 1690, 1456. ¹H NMR (CDCl₃ + CD₃OD one drop) (δ ppm): 13.19 (1H, s, 1-OH), 7.33 (1H, s, H-8), 6.42 (1H, s, H-4), 6.30 (2H, d, J = 6.0 Hz, H-1''), 5.28 (2H, br t, H-2', H-2''), 5.02 (1H, br t, H-6'), 3.37 (2H, d, J = 6.0 Hz, H-1'), 2.03 (2H, m, H-5'), 1.99 (2H, m, H-4'), 1.88 (3H, s, H-5''), 1.80 (3H, s, H-9'), 1.69 (3H, s, H-4''), 1.57 (3H, s, H-8'), 1.52 (3H, s, H-10'). EIMS m/z (% rel. int.): ($[M]^+$ 464, 80), 421 (50), 409 (100), 339 (40), 297 (55), 285 (54), 69 (26). ¹³C NMR (CDCl₃ + CD₃OD one drop) (δ ppm): 180.6 (C-9), 162.5 (C-3), 159.8 (C-1), 156.1 (C-4a), 150.7 (C-6), 150.2 (C-10a), 141.9 (C-7), 135.9 (C-3''), 132.2 (C-3'), 131.5 (C-7'), 124.2 (C-6'), 122.4 (C-2'), 121.3 (C-2''), 115.5 (C-8a), 112.7 (C-5), 110.4 (C-2), 105.1 (C-8), 102.5 (C-9a), 93.4 (C-4), 39.8 (C-4'), 25.8 (C-4''), 25.5 (C-8'), 22.4 (C-1''), 21.7 (C-5'), 21.4 (C-1'), 17.8 (C-9'), 17.6 (C-10'), 16.2 (C-5'').

3.3.4. *Dulcisxanthone F* (4)

Yellow solid, m.p. 213–215 °C. HRESIMS m/z 408.1573 $[M]^+$ (calcd. for $C_{24}H_{24}O_6$ 408.1573). UV (CH₃OH) λ_{max} (nm) (log ϵ): 331 (4.44), 322 (4.43), 265 (4.53), 244 (4.54). IR (neat) ν (cm⁻¹): 3479, 1675, 1587, 1447. ¹H NMR (CDCl₃ + DMSO-*d*₆ one drop) (δ ppm): 13.35 (1H, s, 1-OH), 8.04 (1H, d, J = 9.7 Hz, H-4''), 6.83 (1H, s, H-5), 6.36 (1H, s, H-4), 5.83 (1H, d, J = 9.7 Hz, H-3''), 5.23 (1H, br t, J = 7.0 Hz, H-2'), 3.91 (3H, s, 3-OMe), 3.36 (2H, d, J = 6.7 Hz, H-1'), 1.80 (3H, s, H-5'), 1.68 (3H, s, H-4'), 1.50 (6H, s, 2''-Me). EIMS m/z (% rel. int.): ($[M]^+$

408, 60), 393 (55), 365 (70), 353 (100), 169 (15). ¹³C NMR (CDCl₃ + DMSO-*d*₆ one drop) (δ ppm): 182.4 (C-9), 163.6 (C-3), 159.6 (C-1), 155.5 (C-4a), 153.0 (C-10a), 150.7 (C-6), 136.7 (C-7), 132.2 (C-3''), 131.7 (C-3'), 122.2 (C-2'), 121.0 (C-4''), 119.7 (C-8), 111.4 (C-2), 108.7 (C-8a), 102.2 (C-9a, C-5), 88.9 (C-4), 71.0 (C-2''), 55.8 (3-OCH₃), 29.6 (2''-Me), 27.3 (2''-Me), 25.8 (C-4'), 21.3 (C-1'), 17.7 (C-5').

3.3.5. *Dulcinone* (5)

Yellow solid, m.p. 227–229 °C. HRESIMS m/z 206.0570 $[M]^+$ (calcd. for $C_{11}H_{10}O_4$ 206.0579). UV (CH₃OH) λ_{max} (nm) (log ϵ): 327 (3.51), 300 (3.62), 258 (4.25), 226 (3.96). IR (neat) ν (cm⁻¹): 3343, 1657, 1598, 1417. ¹H NMR (CDCl₃ + DMSO-*d*₆ one drop) (δ ppm): 6.33 (1H, s, H-5), 5.99 (1H, s, H-3), 2.39 (3H, s, H-2), 2.13 (3H, s, H-7). EIMS m/z (% rel. int.): ($[M]^+$ 206, 100), 205 (95), 177 (11), 165 (8). ¹³C NMR (CDCl₃ + DMSO-*d*₆ one drop) (δ ppm): 181.5 (C-4), 165.6 (C-2), 160.9 (C-6), 158.2 (C-8a), 154.6 (C-8), 106.8 (C-3), 102.9 (C-4a), 101.2 (C-7), 97.5 (C-5), 19.3 (2-CH₃), 6.4 (7-CH₃).

3.4. Radical scavenging activity

This was carried out according to the procedure of Deachathai et al. (2005).

3.5. Antibacterial activity

This was carried out according to the procedure of Mahabusarakam et al. (2004).

Acknowledgements

This research was supported by a scholarship to S.D. from the Postgraduate Education and Research Program in Chemistry (PERCH), funded by The Royal Thai Government and the Graduate School, Prince of Songkla University.

References

- Ansari, W.H., Rahman, W., Barraclough, D., Maynard, M.R., Scheinmann, F., 1976. Biflavonoids and a flavanone-chromone from the leaves of *Garcinia dulcis* (Roxb.). *Kurz. J. Chem. Soc. Perkin Trans I*, 1458–1463.
- Blount, J.F., Williams, T.H., 1976. Revised structure of xanthochymol. *Tetrahedron Lett.* 34, 2921–2924.
- Deachathai, S., Mahabusarakam, W., Towatana, N., Phongpaichit, S., Taylor, W.C., 2005. Phenolic compounds from the fruit of *Garcinia dulcis*. *Phytochemistry* 66, 2368–2375.
- Delle Monache, F., Botta, B., Nicoletti, M., de Barros Coelho, J.S., de Andrade Lyra, F.D., 1981. Three new xanthenes and macluraxanthone from *Rheedia benthamiana* Pl. Triana (Guttiferae). *J. Chem. Soc. Perkin Trans. I*, 484–488.
- Gustafson, K.R., Blunt, J.W., Munro, M.H.G., Fuller, R.W., McKee, T.C., Cardellina II, J.H., McMahon, J.B., Cragg, G.M., Boyd, M.R., 1992. The guttiferones, HIV-inhibitory benzophenones from *Sympho-*

- nia globulifera*, *Garcinia livingstonei*, *Garcinia ovalifolia* and *Chusia rosea*. Tetrahedron 48, 10093–10102.
- Harrison, L.J., Leong, L.-S., Leong, Y.-W., Sia, G.-L., Sim, K.-Y., Tan, H.T.-W., 1994. Xanthone and flavonoid constituents of *Garcinia dulcis* (Guttiferae). Nat. Prod. Lett. 5, 111–116.
- Kartnig, Th., Gruber, A., Stachel, J., 1985. Zur kenntnis des flavonoidmusters von *Asparagus officinalis*. Planta Med. 3, 288.
- Kasahara, S., Henmi, S., 1986. Medicine Herb Index in Indonesia. Eisai Indonesia, Jakarta, p. 92.
- Konoshima, M., Ikeshiro, Y., Miyahara, S., Yen, K.-Y., 1970. The constitution of biflavonoids from *Garcinia plants*. Tetrahedron Lett. 48, 4203–4206.
- Mahabusarakam, W., Wiriyaichitra, P., Taylor, W.C., 1987. Chemical constituents of *Garcinia mangostana*. J. Nat. Prod. 50, 474–478.
- Mahabusarakam, W., Deachathai, S., Phongpaichit, C., Jansakul, C., Taylor, W.C., 2004. A benzil and isoflavone derivatives from *Derris scandens* Benth. Phytochemistry 65, 1185–1191.
- Stout, G.H., Christensen, E.N., Balkenhol, W.J., Stevens, K.L., 1973. Xanthones of the Gentianaceae-II *Fraseria albicaulis* Dougl. EX Griesb. Tetrahedron 25, 1961–1973.
- Subhadhirasakul, S., Jankeaw, B., Malinee, A., 2003. Chemical constituents and antioxidative activity of the extracts from *Dyera costulata* leaves. Songklanakarin J. Sci. Technol. 25, 351–353.