

LETTERS TO THE EDITOR

Condensation of Acetamide in Aqueous Solutions in the Presence of Sulfur(IV) Dioxide

R. E. Khoma^{a,b}, A. V. Mazepa^c, A. A. Shestaka^b, A. A. Ennann^b, and V. O. Gelmboldt^b

^a Mechnikov Odessa National University,
ul. Dvoryanskaya 2, Odessa 65082, Ukraine
e-mail: rek@onu.edu.ua, r_khoma@farlep.net

^b Physico-Chemical Institute of Environment and Human Protection,
Ministry of Education and Sciences of Ukraine and National Academy of Sciences of Ukraine, Odessa, Ukraine


^c Bogatskii Physico-Chemical Institute, National Academy of Sciences of Ukraine, Odessa, Ukraine

Received December 10, 2008

DOI: 10.1134/S107036320906036X

The reactions leading to formation of cyclic structures are not typical for amides, in distinct to their thio analogs that under the conditions of acid catalyzed hydrolysis are prone to cyclization processes with formation of N,S-containing heterocycles [1–3].

We found that reaction of acetamide proceeding in SO₂–H₃CC(O)NH₂–H₂O solution affords cyclic *o*-quinoid compound **I** that formally can be regarded as a product of acetamide condensation along the scheme:



In the process of the synthesis of **I** also has been registered formation of acetic acid that can be a result of acetamide hydrolysis in the presence of SO₂. Not excluded that the ability of the observed transformation in the presence of SO₂ is provided due to weak CH-acid properties of H₃CC(O)NH₂ [1].

4,5-Bis(aminomethyl)-3,6-bis(1-aminovinyl)cyclohexa-3,5-diene-1,2-dione (I). A mixture of acetamide (0.2 mol) and 3.6 ml of water (0.2 mol) was placed to a reactor, cooled and kept at the temperature 0°C for 20 min. Then through the mixture was passed gaseous

SO₂ with the rate 10 to 15 ml min^{−1} until homogenous mixture has formed. The viscous oily yellow solution as placed to a freezing chamber (≈ −15°C). After a few weeks the SO₂ excess was neutralized at −5 to −10°C by adding Na₂CO₃ in small portions until gas evolution stopped. The solution was decanted and again placed into freezing chamber (≈ −15°C). After ~14 days the solution obtained was placed to a desiccator over calcium chloride at room temperature and kept until a precipitate has formed. We isolated 5.2 g (62 %) of product **I**, white, easily sublimating at room temperature. The IR spectrum, ν , cm^{−1}: 3420 m [$\nu_{\text{as}}(\text{NH})$]; 3342 s [$\nu_{\text{s}}(\text{NH})$]; 3080 sh [$\nu_{\text{as}}(\text{CH}_2)(=\text{CH}_2)$]; 2930 s [$\nu_{\text{s}}(\text{CH}_2)(=\text{CH}_2)$]; 2857 s [$\nu_{\text{s}}(\text{CH}_2)$]; 1637 m [$\nu(\text{CO})$, $\nu(\text{C}=\text{C})$]; 1562 s [$\delta(\text{NH}_2)$, $\nu(\text{C}=\text{C})_{\text{ring}}$]; 1453 s [$\delta(\text{CH}_2)$, $\delta(\text{CNH})$]; 1397 m [$\delta(\text{CH}_2)_{\text{planar}}$]; 1361 m [$\delta(\text{CH}_2)$, $\nu(\text{NH}_2)$]; 1071 w [$\nu_{\text{as}}(\text{CN})(\text{C}-\text{NH}_2)$]; 1061 m [$\delta(\text{CCC})_{\text{planar, ring}}$]; 969 w [ν_{ring}]; 823 m, br [$\nu(\text{NH}_2)$]; 797 m [$\delta(\text{CH}_2)$]; 733 m [$\delta(\text{CCC})_{\text{out-of-plane, ring}}$]; 711 m [$\nu(\text{CH}_2)$, $\nu(\text{NH}_2)$]; 611 m [$\delta(\text{CCC})_{\text{out-of-plane, ring}}$]; 497 m [$\delta(\text{OCC})_{\text{ring}}$]. The ¹H NMR spectrum (CD₃OD), δ , ppm (*J*, Hz): 1.33–1.37 m (4H, C–NH₂), 1.44–1.53 m (4H, CH₂–NH₂) 2.63 t (4H, CH₂–NH₂, *J*_{HH} 7.2), 4.87 s (4H, C=CH₂). The mass spectrum (FAB, solution of **I** in 3-nitrobenzyl alcohol), *m/z* (*I*_{rel}, %): 250 (100) [*M* + 2H]⁺; (FAB, solution of **I** in 3-nitrobenzyl alcohol with NaI adding) *m/z* (*I*_{rel}, %): 271 (65) [*M* + Na]⁺, 250 (53) [*M* + 2H]⁺. In the FAB mass-spectrum occurs a peak of [*M* + 2H]⁺, characteristic of *o*-quinones that are reduced to the corresponding catechols, that is not inherent in *p*-quinones [5]. Found, %: C 57.43; H 6.41;

N 23.21. $C_{12}H_{16}N_4O_2$. Calculated, %: C 58.05; H 6.50; N 22.57. M 248.3.

The carbon and hydrogen contents were analyzed on a CHN-analyzer. The IR spectra were recorded on a Spectrum BX II FT-IR System instrument (Perkin-Elmer) in the range 4000–350 cm^{-1} , the samples were prepared as pellets with KBr. The 1H , ^{13}C NMR spectra were registered on a Bruker WM 400 spectrometer. The FAB mass spectra were registered on a VG 7070 instrument, desorption of ions from a liquid matrix was carried out by the action of the beam of argon atoms with the energy 8 keV.

REFERENCES

1. *Comprehensive Organic Chemistry*, Barton, D. and Ollis, W.D., Eds., vol. 4, Liverpool: Pergamon Press, 1979. Translated under the title *Obshchaya organicheskaya khimiya*, Moscow: Khimiya, 1983.
2. *Ibid.*, vol. 5, p. 656.
3. *Ibid.*, vol. 9, p. 548.
4. Gel'mbol'dt, V.O., Ilyukhin, A.B., Ganin, E.V., and Ennan, A.A., *Koord. Khimiya*, 2001, vol. 27, p. 573.
5. Vul'fson, N.S., Zaikin, V.G., and Mikaya, A.I., *Mass-spektrometriya organicheskikh soedinenii* (Mass-Spectrometry of Organic Compounds), Moscow: Khimiya, 1986.