

LETTERS TO THE EDITOR

Synthesis, Structure and Hydrazinolysis of 4-(4-Bromo-3,5-dimethyl-1*H*-pyrazol-1-yl)benzaldehyde

S. D. Fazylov^a, O. A. Nurkenov^a, A. E. Arinova^a, Zh. B. Satpaeva^a,
T. M. Seilkhanov^b, A. N. Zhakupova^a, and I. S. Tolepbek^a

^a Institute of Organic Synthesis and Coal Chemistry of the Republic of Kazakhstan,
ul. Alikhanova 1, Karaganda, 100008 Kazakhstan
e-mail: iosu8990@mail.ru

^b Ualikhanov Kokshetau State University, Kokshetau, Kazakhstan

Received July 21, 2014

Keywords: 4-bromo-3,5-dimethylpyrazole, 4-fluorobenzaldehyde, microwave activation

DOI: 10.1134/S1070363214120263


Pyrazole derivatives are used as anti-inflammatory and analgesic drugs, so the synthesis of new representatives of this class of compounds is one of the most promising directions in the search for new biologically active compounds [1, 2]. Since aldehydes are important reagents in organic synthesis of various compounds including heterocyclic, it was interesting to obtain an aromatic aldehyde based on 4-bromo-3,5-dimethylpyrazole and 4-fluoro-benzaldehyde.

However, the nucleophilic substitution reaction involving fluorobenzaldehyde requires prolonged heating [3],


due to the nature of the C–F bond. We found that the synthesis of 4-(4-bromo-3,5-dimethyl-1*H*-pyrazol-1-yl)benzaldehyde **I** can be successfully carried out in DMF under the microwave activation and the catalysis with a specially prepared catalyst supported on silica gel Silpearl activated with potassium carbonate (Scheme 1).

In addition, we performed this reaction under classical conditions (K₂CO₃, DMF, reflux, 18–20 h). The synthesized 4-(4-bromo-3,5-dimethyl-1*H*-pyrazol-1-yl)benzaldehyde **I** is a reactive synthon which is of

Scheme 1.



Scheme 2.



R¹ = OH, R² = H (**II**); R¹ = H, R² = OH (**III**).

interest for a variety of chemical transformations. Thus, the reaction of aldehyde **I** with hydrazides of *ortho*- and *para*-hydroxybenzoic acids in alcoholic medium resulted in the formation of hydrazones **II** and **III** in 85 and 44% yields respectively (Scheme 2).

Composition and structure of the synthesized compounds **I–III** were confirmed by elemental analysis, ^1H NMR and IR spectra.

4-(4-Bromo-3,5-dimethyl-1H-pyrazol-1-yl)benzaldehyde (I). *a. Under microwave irradiation.* To a solution of 0.62 g (0.005 mol) of 4-fluorobenzaldehyde in 10 mL of DMF was added 1 g (0.0056 mol) of 4-bromo-3,5-dimethylpyrazole and 1.54 g of catalyst. The reaction mixture was subjected to microwave irradiation in a Monowave 300 Anton Paar (Austria) apparatus at 140°C for 5 h. After the reaction completion the mixture was filtered and evaporated. The resulting precipitate was recrystallized from 2-propanol. Yield 0.92 g (66%), mp 109–110°C. IR spectrum, ν , cm^{-1} : 1685 (C=O). ^1H NMR spectrum, δ , ppm: 2.12 s (3H, CH_3), 2.39 s (3H, CH_3), 7.79 d (1H, CH_{arom} , J_{HH} 7.8 Hz), 8.05 d (1H, CH_{arom} , J_{HH} 7.9 Hz), 10.07 s [1H, C(O)H]. Found, %: C 52.83; H 4.17; N 10.19. $\text{C}_{12}\text{H}_{11}\text{N}_2\text{OBr}$. Calculated, %: C 51.63; H 3.97; N 10.04.

b. Under classical conditions. To a solution of 1.24 g (0.01 mol) of 4-fluorobenzaldehyde in 20 mL of DMF was added 2 g (0.0113 mol) of 4-bromo-3,5-dimethylpyrazole and 2.07 g (0.015 mol) of potassium carbonate. The reaction mixture was heated on a glycerol bath at 140–150°C for 20 h. The reaction progress was monitored by TLC. After cooling potassium carbonate was filtered off, and the solvent was evaporated. The residue was recrystallized from 2-propanol. Yield 1.68 g (60%).

2-Hydroxy-*N*-[4-(4-bromo-3,5-dimethyl-1H-pyrazol-1-yl)benzylidene]benzoylhydrazide (II). To a solution of 0.23 g (0.00083 mol) of benzaldehyde **I** in 5 mL of ethanol was added 0.13 g (0.00083 mol) of *o*-hydroxybenzoic acid hydrazide. The reaction mixture was heated at 50°C for 15 min. The resulting precipitate was filtered off, washed with ethanol, and

recrystallized from 2-propanol. Yield 0.15 g (44%), yellow powder, mp 169–170°C. IR spectrum, ν , cm^{-1} : 1665 (C=O). ^1H NMR spectrum, δ , ppm: 2.21 s (3H, CH_3), 2.34 s (3H, CH_3), 7.59 d (1H, CH_{arom} , J_{HH} 7.4 Hz), 7.87 d (1H, CH_{arom} , J_{HH} 7.8 Hz), 6.71 d (1H, $\text{CH}_{\text{arom}}^1$, J 8.7 Hz), 7.21 t (1H, $\text{CH}_{\text{arom}}^2$, J 7.8 Hz), 6.59 t (1H, $\text{CH}_{\text{arom}}^3$, J 7.4 Hz), 7.82 d (1H, $\text{CH}_{\text{arom}}^4$, J 8.7 Hz), 8.39 s (1H, CH=N), 14.02 br.s (1H, OH), 14.08 br.m (1H, NH). Found, %: C 55.72; H 4.65; N 14.06. $\text{C}_{19}\text{H}_{17}\text{BrN}_4\text{O}_2$. Calculated, %: C 55.22; H 4.15; N 13.56.

4-Hydroxy-*N*-[4-(4-bromo-3,5-dimethyl-1H-pyrazol-1-yl)benzylidene]benzoylhydrazide (III) was prepared similarly from benzaldehyde **I** and *p*-hydroxybenzoic acid hydrazide. Yield 85%, mp 248–249°C (2-propanol). IR spectrum, ν , cm^{-1} : 1668 (C=O). ^1H NMR spectrum, δ , ppm: 2.19 s (3H, CH_3), 2.32 s (3H, CH_3), 7.59 d (1H, CH_{arom} , J_{HH} 7.4 Hz), 7.87 d (1H, CH_{arom} , J_{HH} 7.8 Hz), 6.68 d (1H, CH_{arom} , J_{HH} 8.7 Hz), 7.75 d (1H, CH_{arom} , J_{HH} 8.7 Hz), 8.4 s (1H, CH=N), 13.95 br.s (1H, OH), 14.05 br.m (1H, NH). Found, %: C 55.57; H 4.49; N 13.81. $\text{C}_{19}\text{H}_{17}\text{BrN}_4\text{O}_2$. Calculated, %: C 55.22; H 4.15; N 13.56.

IR spectra of the compounds **I–III** were recorded on a Nicolet Avatar-320 spectrometer from KBr pellets. ^1H NMR spectra were registered on a Bruker DRX500 spectrometer (500 MHz) in $\text{DMSO}-d_6$ solution, internal reference TMS. Melting points were determined on a Boetius instrument. TLC analysis was performed using Sorbfil plates, eluting with 2-propanol–benzene–ammonia mixture (10 : 5 : 2) and detecting with iodine vapor.

REFERENCES

1. Ivanskii, I.V., *Khimiya geterotsiklicheskikh soedinenii* (Chemistry of Heterocyclic Compounds), Moscow: Vysshaya Shkola, 1978.
2. Mashkovskii, M.D., *Lekarstvennye sredstva* (Drugs), Moscow: RIA "Novaya Volna," 2012.
3. Dryachenko, E.V., Glukhareva, T.V., Nikolaenko, E.F., Tkachev, A.V., and Morzherina Yu.Yu., *Russ. Chem. Bull. Int. Ed.*, 2004, vol. 53, no. 6, p. 1240. DOI: 10.1023/B:RUCB.0000042280.71728.d0.