

Synthesis and Study of New Nitrogen-Containing Heterocycles Based on Glycoluril Derivatives

L. K. Sal'keeva^a, G.-V. Röschenthaler^b, A. A. Bakibaev^c, P. Voitiček^d, A. K. Shibaeva^a,
E. K. Taishibekova^a, A. A. Zhortarova^a, and L. M. Sugralina^a

^a Buketov Karaganda State University, ul. Universitetskaya 28, Karaganda, 100028 Kazakhstan
e-mail: LSalkeeva@mail.ru

^b Jacobs University Bremen, Bremen, Germany

^c National Research Tomsk Polytechnic University, Tomsk, Russia

^d Charles University, Prague, Czech Republic

Received July 28, 2014

Abstract—Functionalization of some cyclic bisureas (glycolurils) and their methylol derivatives was investigated with the goal to obtain new azaheterocycles. The modification of tetra-*N*-methylglycoluril gave unknown before tetra-*N*-methylbisimidazolium tetrachloride and provided a possibility of the synthesis therefrom of glycoluril tetrafluoro derivatives.


Keywords: glycolurils (bisureas), tetra-*N*-methylglycoluril (mebicar), tetra-*N*-methylbisimidazolium tetrachloride, antioxidant activity


DOI: 10.1134/S1070363215010156

Glycoluril (2,4,6,8-tetraazabicyclo[3.3.0]octane-3,7-dione) and its derivatives are applied in various fields of industry. They are used at water purification as disinfectors, as burning inhibitors, at paper manufacturing [1]. The resins containing glycoluril are used at dyes and coatings preparation. Glycoluril derivatives are widely used in the production of hardening polymer compositions, psychotropic agents [2, 3], explosives, and as prolonged fertilizers [4].

Glycoluril is a bicyclic substrate extremely attractive for further functionalization with the goal of synthesis and investigation of new azaheterocycles [5]. A high interest to glycoluril and its derivatives is due to a wide potential of chemical modification of the compounds [6, 7]. In extension of the research in this direction [8, 9] in the present work new possibilities of chemical modification of some glycoluril derivatives were studied, in particular, of 2,4,6,8-tetramethyl-

Scheme 1.





2,4,6,8-tetraazabicyclo[3.3.0]octane-3,7-dione (1,3,4,6-tetramethylglycoluril, mebicar) **I** [10].

Thus, the reaction of 1,3,4,6-tetramethylglycoluril **I** with phosgene in methylene chloride led to formation of before unknown glycoluril tetra-*N*-methylbisimidazolium tetrachloride **II**.

It is known that the introduction of a halogen atom, chlorine or fluorine, into the molecule of potential biologically active compound instead of, e.g., the carbonyl oxygen may cause a change or even strengthening of the biological activity [11]. To obtain the fluorine-containing derivatives of glycoluril, the reaction of compound **II** with potassium fluoride was carried out. The reaction proceeded in anhydrous acetonitrile and led to formation of tetra-*N*-methyl-tetrafluoroglycoluril **III**. The alternative method for preparation of tetra-*N*-methyltetrafluoroglycoluril **III** is the reaction of glycoluril **II** with COF_2 .

The reaction progress was monitored by ^{19}F NMR spectroscopy. In the ^{19}F NMR spectrum of compound **III** there was a signal at 70.9 ppm corresponding to four fluorine atoms.

Aiming to obtain new potential biologically active compounds, functionalization of glycoluril tetra-*N*-methylbisimidazolium tetrachloride **II** was performed with amines of various structures. Thus, tetramethylguanidine, dimethyl- and diethylamines were applied. The reaction was performed at room temperature in anhydrous methylene chloride. The obtained compounds **IV–VI** were crystalline.

The synthesized compounds **I–V** were examined for antioxidant activity. The results are presented in the table. The data illustrate that the activity of compounds

II, IV, V is higher than the activity of trolox used as a reference.

As a result of the performed research new derivatives of glycoluril were prepared; some of them show high antioxidant activity.

EXPERIMENTAL

^1H NMR spectra were recorded on a Bruker DRX-300 and Jeol ECX-400 spectrometers, internal reference TMS. ^{19}F NMR spectra (376 MHz) were registered on a Jeol ECX-400 instrument, internal reference CFCl_3 . Mass spectra were obtained on a Varian Saturn 2000K instrument. Elemental analysis was done on a Eurovector-EA3000 apparatus (Italy). Melting points were determined on a Boetius and MP50 Melting Point System.

2,4,6,8-Tetramethyl-2,4,6,8-tetraazabicyclo[3.3.0]-octane-3,7-dione (I) was prepared by the known procedure [10].

2,5-Dichloro-1,3,4,6-tetramethyl-3,3a,6,6a-tetrahydroimidazo[4,5-*d*]imidazol-1,4-diylum dichloride (II). To a solution of tetramethylglycoluril **I** (1.0 g, 0.005 mol) in CH_2Cl_2 was added with stirring 2.5 g, (0.025 mol) of COCl_2 . The reaction mixture was heated for 2 h at 50–60°C. The solvent was distilled off; the residue was washed with diethyl ether (3–4 times) and dried in argon. Yield 0.8 g (67%), mp 235–237°C. ^1H NMR spectrum (CDCl_3), δ , ppm: 2.95 s (12H, CH_3), 4.91 s (2H, CH). Mass spectrum: m/z 308. Found, %: C 31.36; H 4.32; N 18.47; Cl 45.91. $\text{C}_8\text{H}_{14}\text{N}_4\text{Cl}_4$. Calculated, %: C 31.17; H 4.55; N 18.18; Cl 46.10.

Coefficients of antioxidant activity of glycoluril and its derivatives

Compound	C_w , mol/L	K , $\mu\text{mol L}^{-1} \text{min}^{-1}$	K_{av}
Glycoluril	4.5×10^{-5}	55.9 66.1 59.7 78.2 60.1	64.0
I	4.5×10^{-5}	77.28 81.56 75.69 120.74 80.25	87.104
II	4.5×10^{-5}	130.0 139.64 126.8 138.9 159.6	138.99
IV	4.5×10^{-5}	156.6 170.2 166.38 154.26 160.5	161.59

2,2,5,5-Tetrafluoro-1,3,4,6-tetramethyloxyhydroimidazo[4,5-*d*]imidazole (III). *a.* A mixture of compound **II** (99.74 g, 0.590 mol) and potassium fluoride (102.84 g, 1.770 mol) in anhydrous acetonitrile was heated at 85°C under nitrogen for 6 h. After cooling inorganic salts were separated from the reaction mixture, the residue was distilled. Yield 88.5 g (62%), mp 47°C (37 mmHg). ^1H NMR spectrum (CDCl_3), δ , ppm: 2.52 s (12H, CH_3), 3.05 s (2H, CH). ^{19}F NMR spectrum (CDCl_3): δ_F 70.9 ppm. Mass spectrum: m/z 242. Found, %: C 39.43; H 5.98; N 23.37; F 31.22. $\text{C}_8\text{H}_{14}\text{N}_4\text{F}_4$. Calculated, %: C 39.67; H 5.79; N 23.14; F 31.40.

b. COF_2 (16.5 g, 0.25 mol) was passed through a solution of compound **II** (9.9 g, 0.05 mol) in anhydrous acetonitrile under argon at continuous stirring at 25°C; the reaction progress was monitored by ^{19}F NMR. After distilling off the solvent, the residue was distilled in a vacuum. Yield 8.2 g (68%), mp 47°C (37 mmHg).

***N,N'*-{1,3,4,6-Tetramethyltetrahydroimidazo[4,5-*d*]imidazole-2,5(1*H*,3*H*)-diylidene}bis(*N*-ethyl-ethanoaminium) dichloride (IV).** Diethylamine (0.9 g, 0.006 mol) was added dropwise to compound **II** (0.7 g, 0.003 mol) at room temperature in anhydrous methylene chloride for 3 h. The solvent was distilled off; the formed crystals were washed with anhydrous diethyl ether and dried. Yield 1.8 g (67%), mp 198°C. ^1H NMR spectrum (CDCl_3), δ , ppm: 2.83 s (12H, CH_3), 5.12 s (2H, CH). Mass spectrum: m/z 381. Found, %: C 50.48; H 8.66; N 22.37; Cl 18.41. $\text{C}_{16}\text{H}_{34}\text{N}_6\text{Cl}_2$. Calculated, %: C 50.39; H 8.92; N 22.05; Cl 18.64.

***N,N'*-{1,3,4,6-Tetramethyltetrahydroimidazo[4,5-*d*]imidazole-2,5(1*H*,3*H*)-diylidene}bis(*N*-methyl-methanoaminium) dichloride (V)** was prepared similarly from 0.6 g (0.003 mol) of **II** and 0.7 g (0.006 mol) of dimethylamine. Yield 0.8 g (67%), mp 235°C. ^1H NMR spectrum (CDCl_3), δ , ppm: 2.95 s (12H, CH_3), 4.91 s (2H, CH). Mass spectrum: m/z 325. Found, %: C 44.52; H 8.21; N 25.63; Cl 21.59. $\text{C}_{12}\text{H}_{26}\text{N}_6\text{Cl}_2$. Calculated, %: C 44.31; H 8.00; N 25.85; Cl 21.85.

2,5-Bis{[bis(dimethylamino)methylidene]amino}-1,3,4,6-tetramethyl-3,3a,6,6a-tetrahydroimidazo[4,5-*d*]imidazole-1,4-diylum dichloride (VI) was prepared similarly from 0.8 g (0.003 mol) of **II** and 0.9 g (0.006 mol) of tetramethylguanidine. Yield 1.8 g (57%), mp 204.9°C. ^1H NMR (CDCl_3), δ , ppm: 2.92 s (12H, CH_3), 5.08 s (2H, CH). Mass spectrum: m/z 465. Found, %: C 46.65; H 8.39; N 30.38; Cl 15.02. $\text{C}_{18}\text{H}_{38}\text{N}_{10}\text{Cl}_2$. Calculated, %: C 46.45; H 8.17; N 30.11; Cl 15.27.

Antioxidant activity was determined by the method of cathodic voltammetry (process of oxygen electro-reduction). Trolox was used as a reference.

Experimental procedure included obtaining the voltammogram of cathodic electro-reduction of oxygen with the help of an analyzer connected to PC. The electrochemical cell consisted of a glass vessel filled with a solution of background electrolyte with an indicator glass-carbon electrode immersed in it, a silver chloride electrode of comparison, and a silver chloride auxiliary electrode. 0.1 M solution of sodium perchlorate in ethanol was used as a background solution.

Antioxidant activity of the investigated sample was estimated by the kinetic criterion of antioxidant activity K ($\mu\text{mol L}^{-1} \text{min}^{-1}$), which reflects the amount

of the oxygen forms reacted with the sample in time and is determined by the following equation:

$$K = c(\text{O}_2)/t(1 - I_i/I_0),$$

where $c(\text{O}_2)$ is the concentration of oxygen in the starting solution without the investigated sample, $\mu\text{mol L}^{-1}$; I_i is the value of limiting current of oxygen electro-reduction, μA ; I_0 is the value of limiting current of oxygen electro-reduction in the absence of the investigated sample in a solution, μA ; t is the time of the process, min.

REFERENCES

1. Gazieva, G.A., Nelyubina, Yu.V., Kravchenko, A.N., Lysenko, K.A., Kolotyrkina, N.G., Belyakov, P.A., Korlyukov, A.A., Lebedev, O.V., and Makhova, N.N., *Russ. J. Org. Chem.*, 2009, vol. 45, no. 2, p. 260. DOI: 10.1134/S107042800902016X.
2. Gazieva, G.A., Kravchenko, A.N., Lysenko, K.A., Gaziev, R.G., Lebedev, O.V., and Makhova N.N., *Russ. Chem. Bull. Int. Ed.*, 2008, vol. 57, no. 8, p. 1744. DOI: 10665285/08/57081744.
3. Gazieva, G.A., Kravchenko, A.N., Trunova, N.S., and Makhova, N.N., *Russ. Chem. Bull. Int. Ed.*, 2007, vol. 56, no. 11, p. 2272. DOI: 10665285/07/56112272.
4. Lagona, J., Wagner, B.D., and Isaacs, L., *J. Org. Chem.*, 2006, vol. 71, p. 1181. DOI: 10.1021/jo052294i.
5. Lucas, D., Minami, T., Iannuzzi, G., Cao, L., Wittenberg, J.B., Anzenbacher, P., and Isaacs, L., *J. Am. Chem. Soc.*, 2011, vol. 133, p. 17966. DOI:10.1021/ja208229d.
6. Kravchenko, A.N., Maksareva, E.Yu., Belyakov, P.A., Sigachev, A.S., Chegaev, K.Yu., Lysenko, K.A., Lebedev, O.V., and Makhova, N.N., *Russ. Chem. Bull. Int. Ed.*, 2003, vol. 52, no. 1, p. 192. DOI: 10665285/03/5201192.
7. Kravchenko, A.N., Sigachev, A.S., Maksareva, E.Yu., Gazieva, G.A., Trunova, N.S., Lozhkin, B.V., Pivina, T.S., Ilyin, M.M., Lysenko, K.A., Nelyubina, Yu.V., Davankov, V.A., Lebedev, O.V., Makhova, N.N., and Tartakovsky V.A., *Russ. Chem. Bull. Int. Ed.*, 2005, vol. 54, no. 3, p. 691. DOI: 10665285/05/54030691.
8. Petersen, H., *Synthesis*, 1973, vol. 5, p. 257.
9. Salkeeva, L.K., Shibaeva, A.K., Bakibaev, A.A., Tayshibekova, E.K., Minaeva, E.V., Zhortarova, A.A., and Salkeeva, A.K., *Russ. J. Gen. Chem.*, 2014, vol. 84, no. 2, p. 338. DOI: 10.1134/S1070363214020339.
10. Kravchenko, A.N., Gazieva, G.A., Kolotyrkina, N.G., and Makhova N.N., *Russ. Chem. Bull. Int. Ed.*, vol. 57, no. 8, p. 1760. DOI: 10665285/08/57081760.
11. Dekaprilevich, M.O., Kravchenko, A.N., Khmelnytsky, L.I., Lebedev, O.V., Struchkov, Yu.T., and Strelenko, Yu.A., *Russ. Chem. Bull.*, 1995, vol. 44, no. 1, p. 131. DOI: 10665285/95/44010131.