

REVIEW

*Dedicated to 125th Anniversary of outstanding German chemist,
Nobel Prize Winner H. Fischer*

Modern Aspects of the Chemistry of Protoporphyrin IX

V. Yu. Pavlov

*Faculty of Chemistry, Moscow State University, Vorob'evy gory 1, Moscow, 119992 Russia
e-mail: vyupavlov@rambler.ru*

Lomonosov Moscow State Academy of Fine Chemical Technology, pr. Vernadskogo 86, Moscow, 119571 Russia

Received February 27, 2006

Abstract—Numerous recently published data on the methods for modification of protoporphyrin IX and its derivatives have been summarized and analyzed. Special attention has been given to those transformations which are important from the practical viewpoint for the preparation of compounds possessing valuable medical, biological, and physical properties, as well as to specific transformations inherent to the tetrapyrrole macroring.

DOI: 10.1134/S10704280007010010

1. Introduction	2
2. Nomenclature and Trivial Names of Basic Protoporphyrin IX Derivatives	2
3. Modification of Peripheral Substituents	2
3.1. Modification of Vinyl Groups	3
3.1.1. Reduction	3
3.1.2. Oxidation	3
3.1.3. Addition to Vinyl Groups	3
3.1.4. Electrocyclic Reactions Involving Vinyl Groups	9
3.1.5. Substitution at Vinyl Groups	11
3.1.6. Elimination of Vinyl Groups	11
3.1.7. Olefin Metathesis	13
3.2. Modification of Formyl Groups	15
3.2.1. Reduction	15
3.2.2. Oxidation	16
3.2.3. Nucleophilic Addition to Formyl Groups	16
3.2.3.1. Synthesis of Imino Derivatives	16
3.2.3.2. Synthesis of Vinyl Derivatives	17
3.2.3.3. Nucleophilic Addition to Formyl Groups without Formation of Multiple Bonds	20
3.3. Modification of 13- and 17-(3-Alkoxy-3-oxopropyl) Groups	20
3.3.1. Reduction of Ester Groups	20
3.3.2. Hydrolysis of Ester Groups	20
3.3.3. Different Versions of Esterification of 13- and 17-(2-Carboxyethyl) Groups in Porphyrins	22
3.3.4. Amide Derivatives Based on 13,17-Bis(2-carboxyethyl)porphyrins	22
3.4. Modification of Methyl Groups	23



Vsevolod Pavlov was born in 1978 in Moscow. In 2000, he graduated from the Faculty of Chemistry, Moscow State University, and in 2001, from the English Language Department, Higher Courses of Foreign Languages at the Ministry of Foreign Economic Relations of the Russian Federation. Candidate of Chemical Sciences since 2006. V. Pavlov is scientific worker at the Organic Chemistry Department, Faculty of Chemistry, Moscow State University, and at the Faculty of Biotechnology and Organic Synthesis, Lomonosov Moscow State Academy of Fine Chemical Technology.

Fields of scientific interest: organic and medicinal chemistry, synthesis and reactivity of low-molecular heterocyclic compounds, and chemistry and biological activity of derivatives of natural macrocyclic compounds.

4. Modification of Macroring	23
4.1. Modifications at <i>meso</i> Positions	23
4.2. Modifications Involving 2–3, 7–8, 12–13, and 17–18 Double Bonds	27
4.3. Metal Complexes and Their Demetalation	28
4.4. Modifications Involving Rearrangement Processes	28
5. Conclusion	30

1. INTRODUCTION

Protoporphyrin IX occupies a specific place among numerous families of tetrapyrroles. It is a parent structure of a large group of natural tetrapyrroles responsible for vital functions of aerobic organisms [1, 2]. In particular, protoporphyrin IX acts as ligand toward prosthetic group of proteins which ensure transport (hemoglobin) and deposition of oxygen (myoglobin) [2–4], as well as toward enzymes responsible for key bioregulatory functions such as detoxication and synthesis of sex hormones and antiinflammatory and anti-hypertensive compounds (cytochrome P450 family) [5–7]. Strong interest in the chemistry of protoporphyrin IX originates from unique combination of valuable physical, chemical, and biological properties [8] of its purposefully modified derivatives and their synthetic accessibility [9]. Effective catalysts of organic reactions [10–12], components of sensor complexes [13–15], and convenient models for studying photosynthesis [16, 17] and oxygen transport *in vivo* [18] were found among protoporphyrin IX derivatives.

An exceptional combination of profitable photophysical properties, low systemic toxicity, and affinity for hyperproliferating tissues makes protoporphyrin IX derivatives very promising for use in the diagnostics and therapy of cancer and eye diseases, as well as in cardiology, cosmetology, and other fields of medicine [8, 19–28].


The chemistry of protoporphyrin IX has started to extensively develop since mid XXth century due to H. Fischer's studies; since 1934 till 1940 he published a collection of fundamental works on the synthesis and chemical transformations of tetrapyrrole compounds [29–31]. Voluminous data on the chemistry of protoporphyrin IX derivatives were covered by books [32–34] and several reviews [9, 20–22, 35–37], as well as by the 20-volume series *The Porphyrin Handbook* published in 2000–2003 [38, 39].

The present review summarizes and analyzes data published in the past decade on modifications of protoporphyrin IX derivatives with account taken of publications not covered by the above reviews and mono-

graphs. The data are discussed with respect to the substitution pattern in protoporphyrin IX and the possibility for modification of the tetrapyrrole macroring.

2. NOMENCLATURE AND TRIVIAL NAMES OF BASIC PROTOPORPHYRIN IX DERIVATIVES

The structures of protoporphyrin IX (**1**) and its some most important derivatives **2–13** are given below. Atoms in the porphyrin macroring are numbered according to Fischer or IUPAC recommendations [36]. This review uses the IUPAC atom numbering, except for trivial names proposed by Fischer.



1, M = 2H, R¹ = R² = CH₂=CH (protoporphyrin IX); **2**, M = 2H, R¹ = CHO, R² = CH₂=CH (chlorocruoroporphyrin); **3**, M = 2H, R¹ = R² = H (deuteroporphyrin); **4**, M = 2H, R¹ = R² = HOCOCH₂CH₂ (coproporphyrin III); **5**, M = 2H, R¹ = CH₂=CH, R² = HOCOCH₂CH₂ (harderoporphyrin); **6**, M = 2H, R¹ = CHO, R² = CH=CHR³ (R³ = C₁₂H₂₅ to C₁₅H₃₁) (cryptoporphyrin *a*); **7**, M = 2H, R¹ = R² = MeCH(OH) (hematoporphyrin IX); **8**, M = 2H, R¹ = CH₂=CH, R² = H (pentoporphyrin); **9**, M = 2H, R¹ = HOCOCH=CH, R² = HOCOCH₂CH₂ (porphyrin S-411); **10**, M = Fe²⁺, R¹ = R² = CH₂=CH (heme); **11**, M = [FeCl]²⁺, R¹ = CHO, R² = CH₂=CH (chlorocruorohemin); **12**, M = [FeCl]²⁺, R¹ = R² = CH₂=CH (protohemin IX); **13**, M = [FeOH]²⁺, R¹ = R² = CH₂=CH (hematin).

3. MODIFICATION OF PERIPHERAL SUBSTITUENTS

Analysis of the structure of protoporphyrin IX (**1**) shows that the main peripheral substituents capable of being involved in various chemical transformations are vinyl, 13- and 17-(2-carboxyethyl), and methyl groups. Data on possible modifications of these substituents are discussed below.


3.1. Modification of Vinyl Groups

A combination of high reactivity of the vinyl groups in protoporphyrin IX derivatives and considerable stability of the porphyrin macroring is responsible for their easy transformations such as reduction, oxidation, addition, substitution, elimination, electrocyclic reactions, and olefin metathesis.

3.1.1. Reduction. Specific features of the reduction of peripheral vinyl groups in porphyrins to ethyl substituents were considered in the classical treatise [33]. A detailed discussion of practical large-scale methods for hydrogenation of the vinyl groups in protohemin IX (**12**) was given in patent application [40].

The porphyrin vinyl groups are converted into branched 1-formylethyl substituents by hydroformylation catalyzed by rhodium complexes with organophosphorus ligands [41, 42]. The regioselectivity of the process strongly depends on the central metal ion. Zinc complex **14** gives rise to 100% of aldehyde **15**, while the yield of formylethyl derivative **17** from nickel complex **16** is 75% (Scheme 1) [42].

Scheme 1.



14, 15, M = Zn(II); **16, 17**, M = Ni(II); hereinafter, "p" stands for $\text{CH}_2\text{CH}_2\text{CO}_2$, i.e., $\text{pMe} = \text{CH}_2\text{CH}_2\text{CO}_2\text{Me}$.


3.1.2. Oxidation. The vinyl groups in divinylporphyrin **18** can be oxidized to formyl substituents with the aid of a large number of reagents [9, 33, 43]. The general disadvantage of direct oxidation methods is that they lead to the formation of mixtures of monoformyl derivatives **19** and **20** which are difficult to separate (Scheme 2). The use of the OsO_4 -4-methylmorpholine *N*-oxide system ensures complete oxidation of both vinyl groups. In this case, porphyrin **18** is converted into bis-diol **21** in almost quantitative yield, and treatment of the latter with HIO_4 gives dialdehyde **22** in a high yield [44]. Compound **21** is the key intermediate product in the synthesis of BOPP, an efficient agent for photodynamic therapy of cancer [44]. Divinylporphyrin **18** is oxidized in high yield to bis-

acetal **23** by the action of 3 equiv of $\text{Ti}(\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$ in $\text{MeOH}-\text{CH}_2\text{Cl}_2$; acid hydrolysis of **23** gives bis-aldehyde **24**, and reduction of the latter with NaBH_4 leads to diol **25** which may formally be regarded as the anti-Markownikow vinyl group hydration product [33, 44]. The direct anti-Markownikow hydration (by treatment with B_2H_6) of vinyl groups in porphyrins is characterized by a poor yield [9]. Nucleophilic replacement of the hydroxy groups in diol **25** by chlorine atoms may be achieved by treatment with SOCl_2 in CHCl_3 -DMF in the presence of K_2CO_3 . Dichloro derivative **26** is transformed in a high yield into initial divinylporphyrin **18** by the action of *t*-BuOK over a period of 3 days (Scheme 3) [33]. Under analogous conditions, 5-methylporphyrin **27** is not converted into divinylporphyrin **28**, but closure of six-membered ring occurs to give compound **29** (Scheme 4) [45]. An effective and universal method for regeneration of vinyl group from 2-hydroxyethyl substituent consists of treatment of diols **25** and **30** with the system $o\text{-NO}_2\text{C}_6\text{H}_4\text{SeCNO}-\text{Bu}_3\text{P}$, followed by oxidation of the corresponding organoselenium porphyrin derivatives **31** and **32** with hydrogen peroxide [45]. This procedure makes it possible to shorten the reaction time to 5 h, avoid undesirable cyclization to compound **29**, and obtain 5-methyl-substituted analog of protoporphyrin IX **28** with high regioselectivity [45]. Transformations of the hydroxyethyl groups in **25** were used in the design of efficient photosensitizers for photodynamic therapy of cancer [22, 46] and in some classical syntheses [9, 33].


Treatment of divinylporphyrin **18** with 2 equiv of $\text{Ti}(\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$ in $\text{MeOH}-\text{CH}_2\text{Cl}_2$ resulted in oxidation of only one vinyl group; the subsequent hydrolysis, reduction with NaBH_4 , and chromatographic separation gave individual monohydroxy isomers **33** and **34** that are important intermediate products in a number of classical syntheses of key natural porphyrins [33]. The oxidation of **18** under severe conditions (with a hot solution of KMnO_4 in Me_2CO) leads to the formation of dicarboxylic acid **35** [33].

3.1.3. Addition to vinyl groups. Electrophilic addition of aliphatic amines to the vinyl groups in protoporphyrin IX derivatives occurs on prolonged heating. Heating of porphyrin **18** in boiling ethylenediamine was accompanied by amidation of the ester groups to give compound **36**. Brunner et al. [47] proposed to use the diamine fragments in **36** as bidentate chelating ligands toward Pt(II). The reaction of **36** with K_2PtCl_4 gave complex **37** containing a photodynamically active

Scheme 2.




Scheme 3.



31, R = *o*-O₂NC₆H₄Se.

Scheme 4.




27, R = Cl; **30**, R = OH; **31**, **32**, R = *o*-O₂NC₆H₄Se.


porphyrin chromophore and a cytotoxic Pt(II) coordination entity (Scheme 5). Pilloud et al. [48, 49] reported on effective addition of thiols immobilized on the surface of a quartz or gold electrode to the por-

phyrin vinyl groups. Sulfanyl derivatives **38** were obtained via autooxidation of a non-porphyrin macrocyclic ring to porphyrin, by reaction of cysteine and other S-nucleophiles with protoporphyrinogen **39** under

Scheme 5.



Scheme 6.



38, R = HOCOCH(NH₂) or CH-peptide.

acidic conditions. Compound **39** was synthesized from porphyrinogen **40** disodium salt which was generated from protoporphyrin IX disodium salt (**41**) by the action of Na/Hg under nitrogen (Scheme 6) [50].


A widely used procedure for functionalization of the vinyl groups in protoporphyrin IX is based on hydrobromination with the system HBr–AcOH, followed by replacement of the labile bromine atoms by various nucleophiles [20, 21, 51, 52]. The efficiency of this approach was demonstrated by the synthesis of promising photosensitizers for photodynamic therapy of cancer [20, 21, 53, 54].

Addition of alcohols to the vinyl groups in divinylporphyrin **18** was the initial step in the synthesis of effective diagnostic agents [22, 55]. Successive treatment of **18** with HBr–AcOH and ethylene glycol gave a mixture of isomeric ethers **42** and **43** (Scheme 7).

Metalation of **43**, hydrolysis of the ester groups, and esterification of the hydroxy group in the 8-[1-(2-hydroxyethoxy)]ethyl substituent with diethylenetriaminepentaacetic acid led to the formation of compound **44** (ATN-10). Due to the presence in its molecule of a diethylenetriaminepentaacetic acid residue which effectively coordinates various metals, ATN-10 was used to chelate radioactive metal isotopes. Intravenous administration of a solution obtained by mixing ATN-10 with sodium pertechnetate (as source of [⁹⁹Tc]) in saline at a dose of 0.1 mg/kg gave a distinct scintigraphic pattern of bone marrow in 5 min after injection [55].


Tian et al. [56] reported on acid hydrolysis of the ether bond in the 3(8)-(1-methoxyethyl) group in porphyrins. The subsequent transformation of the 3(8)-(1-hydroxyethyl) group into vinyl was effected by heating in a mixture of benzoyl chloride with DMF or in

Scheme 7.




44, M = Mn(III).

Scheme 8.



45, R¹ = R² = (*R,S*)-CF₃CH(OH), R³ = R⁴ = Me; 49, R¹ = (*R*)-CF₃CH(OH), R² = (*S*)-CH₃CH(OH), R³ = R⁴ = Me; 50, R¹ = (*S*)-CF₃CH(OH), R² = (*R*)-CF₃CH(OH), R³ = R⁴ = Me; 51, R¹ = R² = (*R*)-CF₃CH(OH), R³ = R⁴ = Me; 52, R¹ = R² = (*S*)-CF₃CH(OH), R³ = R⁴ = Me; 61, R¹ = (*S*)-CF₃CH(OH), R² = (*R*)-CF₃CH(OH), R³ = R⁴ = Na.

Scheme 9.



1,2-dichlorobenzene in the presence of *p*-toluenesulfonic acid; protoporphyrin IX dialkyl esters were thus obtained [9, 21, 57].

Hematoporphyrin IX (**7**) possesses two chiral centers and is therefore a mixture of four diastereoisomers [58]. With the goal of elucidating the effect of steric structure on biological activity of porphyrins, Ando et al. [58, 59] tried to perform optical resolution of trifluoromethyl analogs **45–47** of hematoporphyrin IX with the aid of (*S*)-camphoroyl chloride (**48**). The authors failed to separate bis(2,2,2-trifluoro-1-hydroxyethyl) derivative **45** into individual stereoisomers **49–52**; the latter were obtained only via total syntheses from chiral monopyrrole compounds [58]. Monosubstituted derivatives **46** and **47** were separated into *R* and *S* isomers **53/54** and **55/56** by column chromatography at the stage of formation of the corresponding esters which were treated with sodium hexamethyldisilazide in MeOH–THF to obtain disodium salts **57–60** (Scheme 8). Experiments on a stomach cancer model *in vitro* showed that (3*S*)-isomer **59** is taken up by tumor cells 15 times more selectively than is (3*R*)-isomer **57** and that the selectivity of (8*R*)-isomer **58** for tumor cells is slightly lower as compared to (8*S*)-isomer **60** [58, 59]. The most active among disubstituted derivatives is (3*S*,8*R*)-isomer **61** [58].

The 3- and 8-[1-hydroxy(alkoxy)ethyl] groups in porphyrins are capable of reacting with various nucleo-

philes to give the corresponding replacement products due to influence of the tetrapyrrole macroring which activates pseudobenzylic positions in the peripheral substituents. Various products can be obtained, depending on the activation method and nucleophile nature [20, 21, 36, 60–62]. For example, treatment of 8-(1-hydroxyethyl)benzoporphyrin **62** with trifluoromethanesulfonic acid gives a mixture of *cis*- and *trans*-bis-benzoporphyrins **63** and **64** in more than 40% yield [57] (Scheme 9). An alternative method of activation of 3- and 8-(1-hydroxyethyl) groups makes use of a Zn(OAc)₂·2H₂O–nucleophile system [60, 61, 63]. Bis-acetylacetonate derivative **65** was obtained in a high yield by heating hematoporphyrin IX (**7**) with acetylacetone in the presence of Zn(OAc)₂·2H₂O, followed by esterification with H₂SO₄–MeOH [61]. Further transformations involving the acetylacetonate moieties in **65** were described in [60, 63–65]. Condensation of **65** with substituted phenylhydrazines **69–71** gave pyrazole derivatives **66–68** (Scheme 10), thus demonstrating the possibility of using acetylacetonate porphyrin derivatives as synthons [64, 65]. Some products of modification of the acetylacetonate moiety showed a high photodynamic activity [66–69].


1-Hydroxyethyl groups in positions 3 and 8 of the porphyrin macroring are readily oxidized to acetyl; depending on the oxidizing system, partial or complete oxidation is achieved. The system Pr₄NRuO₄–4-meth-

ylmorpholine *N*-oxide oxidizes both hydroxy groups in porphyrin **72** in a high yield [70]; in reactions with less active MnO_2 , monoacetyl derivatives **73** and **74** were mainly formed (yield 67%), while the fraction of diacetylporphyrin **75** was insignificant [62] (Scheme 11). By treatment with NaBH_4 the acetyl groups can be reduced to hydroxyethyl in a nonstereoselective fashion


[9, 57, 62, 71]. Asymmetric reduction of diacetyl derivative **75** with $\text{BH}_3 \cdot \text{Me}_2\text{S}$ in the presence of chiral methyloxazaborolidines ensured enantioselective preparation of hematoporphyrin IX dimethyl ester stereoisomers **72** [72].

Classical nucleophilic addition reactions of acetyl groups in porphyrins were discussed in detail in


Scheme 10.



Scheme 11.




Scheme 12.



reviews [9, 33]. Kabachnik et al. recently reported on the reaction of diacetylporphyrin **75** with $(\text{EtO})_2\text{P}(\text{O})\text{H}-t\text{-BuNH}_2$ under conditions of microwave activation, which led to the formation of α -aminophosphonate **76** in a high yield [73, 74] (Scheme 12). Treatment of **75** with $\text{Ti}(\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}-\text{MeOH}$ in the presence of concentrated nitric acid involved oxidative rearrangement of the acetyl groups into methoxycarbonyl substituents to give tetraester **77** [75]. Diacetylporphyrin **75** reacted with POCl_3-DMF (Vilsmeier reaction), yielding dichloride **78**, and treatment of the latter with KOH in MeOH resulted in the formation of protoporphyrin IX analog **79** containing acetylenic groups instead of vinyl. Compound **79** was converted into hemin (**80**) by the action of FeCl_2 [76].

3.1.4. Electrocyclic reactions involving vinyl groups. Molecules of some protoporphyrin IX derivatives contain diene fragments formed by peripheral vinyl groups and double bonds in the macroring; such compounds are capable of reacting with C- [9, 57], N- [77], O- [58, 65, 78–83], and other dienophiles to give cycloaddition products. The ability of porphyrin **18** to effectively react with electron-deficient alkenes and alkynes is well known [9, 36, 84]. From the practical viewpoint, the most important is the reaction with dimethyl acetylenedicarboxylate; this reaction is the key step in the synthesis of Visudyne (**81, 82**) [20, 22–24, 85], an efficient drug for the photodynamic treatment of eye diseases. Compounds **84–87** containing heterocyclic substituents were obtained by reaction of porphyrin **18** with azo compounds **83** as dienophiles




81, 82 $\text{R}^1 = \text{Me}, \text{R}^2 = \text{H}; \text{82}, \text{R}^1 = \text{H}, \text{R}^2 = \text{Me}.$

(Scheme 13). The transformation involves concurrent [4+2]- and [2+2]-cycloaddition processes [77]. The reaction of divinylporphyrin **18** with singlet oxygen is a classical transformation in the chemistry of protoporphyrin IX [9]. Monovinylporphyrins also react with singlet oxygen [65, 78, 79, 81, 83]. Porphyrins **88–91** give rise to photoporphyrins **92–95** [65, 83] (Scheme 14). Presumably, the reaction involves intermediate formation of *endo*-peroxides **96–99** which instantaneously rearrange into stable products **92–95** (the photoporphyrin fragment is shown in Scheme 14 in boldface mode) [65, 83]. Grandadam et al. [78] showed that a chlorin-type photosensitizer is highly effective for the inactivation of human immunodeficiency viruses (HIV-1). Kygova et al. [86] demonstrated high immunosuppressive activity of photo-oxidation products of protoporphyrin IX dimethyl ester *in vivo*.


The reaction of divinylporphyrin **18** with diazomethane gives a mixture of pyrazolylporphyrins **100–**

Scheme 13.




83–87, $\text{R} = \text{Me}, \text{Et}, t\text{-Bu}, \text{Ph}, \text{NH}_2.$

Scheme 14.




88, 92, 96, R = OH; 89, 93, 97, R = OAc; 90, 94, 98, R = acac; 91, 95, 99, R = Ht.

Scheme 15.




Scheme 16.



107–109, R¹ = R² = R³; 112–115, R¹ = R² = R⁴; 110, Hlg = Br, I, R⁵ = H, CHO, AlkO.

Scheme 17.



102 in a good overall yield [87, 88], and irradiation of the latter with long-wave (λ 550–650 nm) light leads to cyclopropyl derivatives **103–105** [87] (Scheme 15). The use of pyrazolylporphyrins as photosensitizers for effective binding of substrates in selective photodynamic therapy was covered by patent [89]. The reaction of vinylporphyrins with ethyl diazoacetate to form cyclopropyl derivatives was used by Fischer and Medick [90] to prove the presence of vinyl groups.

3.1.5. Substitution at vinyl groups. Heck reaction of protoporphyrin IX derivatives allows preparation of functionalized olefin derivatives [36, 91–93]. The reaction of zinc complex **14** with modified nucleoside **106** in the presence of LiPdCl_3 , followed by demetalation, afforded a mixture of bis-*trans*, 3-*gem*-8-*trans*, and 8-*gem*-3-*trans* isomers **107–109** in an acceptable yield, while no bis-*gem* isomers were detected [92] (Scheme 16). Study of the reaction of zinc complex **14** with aryl halides **110** [93] showed that the use of palladium catalyst **111** improves the yield of arylvinylidene derivatives **112–115**; the reaction was nonstereoselective, and isomers **112–115** were difficult to separate. Stereoselective substitution at the vinyl groups of protoporphyrin IX and its metal complexes readily occurs by the action of Eschenmoser's salt ($\text{H}_2\text{C}=\text{NMe}_2^+ \Gamma^-$) [94]. Treatment of **14** with $\text{H}_2\text{C}=\text{NMe}_2^+ \Gamma^-$ gave bis-*trans*-dimethylaminomethyl derivative **116** with high stereo-selectivity; the subsequent reaction with methyl iodide led to water-soluble ammonium salt **117** in


100% yield (Scheme 17). Cationic porphyrin **117** was shown to effectively bind and cleave DNA in the presence of light [94].

Terminal protons in the vinyl groups of porphyrin metal complexes are replaced by formyl groups under the Vilsmeier reaction conditions (POCl_3 –DMF) to give acrolein derivatives [9]. Prolonged heating of protoporphyrin IX with *p*- $\text{MeC}_6\text{H}_4\text{SO}_3\text{D}$ and D_2O in 1,2-dichlorobenzene resulted in ~90% replacement of the terminal methylene protons by deuterium [9].

3.1.6. Elimination of vinyl groups. Vinylporphyrins readily lose vinyl groups on heating in fused resorcinol [9, 33]. Protohemin IX (**12**) is thus converted into deuterohemin IX (**118**) [9, 33]. The mechanism of this transformation was discussed in detail in [9]. Devinylation is an important reaction in the chemistry of protoporphyrin IX, for it makes the 3- and 8-positions in the macroring vacant, thus enabling electrophilic substitution reactions with various reagents [9, 95]. Additional synthetic approaches become possible via transformation of iron(III) complex **118** into the free base (metal-free porphyrin) **119** and then into a series of metal complexes, e.g., **120–123** (Scheme 18).


In some cases [9, 96], variation of the reaction conditions ensures controlled preparation of mono- and disubstituted products. For example, treatment of copper complex **120** with acetic anhydride– SnCl_4 for

Scheme 18.




120, M = Cu(II); **121**, M = Pt(II); **122**, M = Zn; **123**, M = Ni(II).

Scheme 19.




Scheme 20.



119, **129**, **130**, *M* = 2H; **120**, *M* = Cu(II); **121**, *M* = Pt(II); **122**, *M* = Zn; **123**, *M* = Ni(II); **127**, **128**, *M* = Pt(II); **131**, *M* = 2H, *R* = Me₂N; **132**, *M* = Zn, *R* = Me₂N; **133**, *M* = 2H, *R* = Me₃N⁺I⁻; **134**, *M* = Zn, *R* = Me₃N⁺I⁻; **138**, *Hlg* = Cl; **139**, *Hlg* = Br.

Scheme 21.



a short time (a few seconds), followed by HPLC separation, gives individual monoacetyl derivatives **124** and **125** in a high yield; considerable extension of the reaction time leads to predominant formation of diacetylporphyrin **126** [9] (Scheme 19). Monoacetylporphyrins **124** and **125** were intermediate products in classical syntheses of some important natural porphyrins [9].

Analysis of the classical data on the formylation of porphyrin metal complexes [97] shows that the reaction occurs at both *meso*- and 3- and/or 8-positions (provided that these positions are vacant). By contrast, Rumyantseva et al. [98] recently reported on the Vilsmeier formylation of Pt(II) complex **121** only at positions 3 and 8 with formation of monoformyl derivatives **127** and **128** (Scheme 20). High regio-

selectivity in the formylation of 3(8)-unsubstituted deuteroporphyrin IX copper complexes with a system HC(OMe)_3 –Lewis acid was demonstrated in [99, 100]. Thus, copper complex **120** was treated with the system HC(OMe)_3 – SnCl_4 , and subsequent demetalation and chromatographic separation gave pure porphyrins **129** and **130** in 46 and 47% yield, respectively [100].

Porphyrin **119** was converted into 3,8-bis-dimethylaminomethyl derivative **131** by reaction with dimethyl(methylidene)ammonium iodide. Analogous reaction with electron-donor Zn complex **122** gave the corresponding dimethylaminomethyl derivative **132** in a higher yield than in the reaction with free base **119**. Quaternization of dimethylaminomethyl derivatives **131** and **132** with methyl iodide leads to ammonium salts **133** and **134** which are capable of effectively binding DNA [94]. Ando and Kumadaki [58] used the system CF_3CHO – AlCl_3 to introduce $\text{CF}_3\text{CH(OH)}$ groups into positions 3 and 8 of deuteroporphyrin **119**; as a result, mono- and disubstituted products **135**–**137** were obtained in 27, 23, and 8% yield, respectively (Scheme 21).

Organomercury deuteroporphyrin IX derivatives **138** and **139** were synthesized in a high yield by heating zinc complex **122** with mercury(II) acetate in MeOH – THF , followed by treatment with a saturated solution of NaCl or NaBr ; compounds **138** and **139** are valuable synthons in the Heck reaction [9, 36, 101, 102]. 3,8-Diiodoporphyrin **140** [84, 102, 103] attracts strong interest as substrate for palladium-catalyzed reactions with protoporphyrin IX derivatives; it can be


prepared in more than 90% yield from deuteroporphyrin IX (**119**) by iodination with I_2 in the presence of K_2CO_3 [103]. The formation of mono- and dibromo derivatives in the reaction of deuteroporphyrin IX **119** with *N*-bromosuccinimide in an inert solvent may be controlled by varying the reactant ratio; the products are quantitatively converted into zinc complexes **141**–**143** [96] as substrates for Pd-catalyzed reactions [96, 101, 104] (Scheme 22).

An elegant synthesis of di- and trimeric porphyrins via Heck reaction using the system Pd(OAc)_2 – LiCl – K_2CO_3 – Bu_4NBr was proposed in [96]. The reaction of bromoporphyrin **142** with excess 1,4-divinylbenzene **144** gave 67% of *trans*-vinylstyryl derivative **145**, while in the presence of excess porphyrin **142** 73% of dimer **146** and 22% of **145** were formed (Scheme 22). An analogous approach may be used to synthesize trimeric porphyrins. For instance, ensemble **147** consisting of three porphyrin macrorings linked through 1,4-divinylbenzene moieties was obtained in 83% yield with high regioselectivity by reaction of **145** with dibromoporphyrin **143** at a ratio of 2.11. Finally, the reaction of 3 equiv of monobromo derivative **142** with 1,3,5-trivinylbenzene **148** gave 56% of trimer **149** [96] (Scheme 23).


A detailed analysis of prospects in using Pd-catalyzed reactions for functionalization of natural porphyrins was given in [58, 101].

3.1.7. Olefin metathesis. In 2004 Liu et al. [105] were the first to demonstrate the possibility of using protoporphyrin IX derivatives in metathesis of olefins

Scheme 22.




Scheme 23.



catalyzed by ruthenium complex **150** [105]. By reactions of benzoporphyrins **151** and **152** with alkenes **153** olefinic derivatives **154** and **155**, respectively, were obtained in high yields (Scheme 24). Electron-

donor zinc complex **152** reacted more readily than free base **151**. In all cases, the process was characterized by high *E*-stereoselectivity, presumably due to steric effect of the tetrapyrrole macroring [105].

Scheme 24.



3.2. Modification of Formyl Groups

Protoporphyrin IX (**1**) molecule has no formyl group, but such group is present in its closest analogs, e.g., chlorocruoroporphyrin (**2**), cryptoporphyrin *a* (**6**) and different hemes [1]. The data discussed in the preceding section indicate that the vinyl groups in protoporphyrin IX (**1**) can readily be transformed into formyl moieties as characteristic functionalities of fundamental natural porphyrins. Therefore, the present section considers methods for modification of formyl groups in protoporphyrin IX derivatives.


3.2.1. Reduction. The formyl groups in protoporphyrin IX derivatives are reduced to hydroxymethyl by the action of NaBH₄ [33, 79, 82, 83, 106]; they are converted into methyl substituents under the Wolff–Kishner reaction conditions [9]. Procedures for further modification of hydroxy groups in protoporphyrin IX derivatives were developed [21, 33, 47, 51, 62, 65, 79, 82, 83, 106] (see also Section 3.1.2).

From the practical viewpoint, introduction of carborane moieties (as a source of boron atoms for boron neutron capture therapy) via acylation of hydroxy


groups in protoporphyrin IX derivatives with carboranyl chlorides [22, 26, 62, 107] attracted interest. The reaction of bis-diol **21** with chloride **156** in the presence of 4-dimethylaminopyridine gave tetraester **157** containing 4 carborane fragments [22] (Scheme 25). Analogous reaction with bis-*p*-carboranyl acid chloride led to the formation of a porphyrin cluster containing eight carborane cages [107]. Tetraester **157** is a synthetic precursor of effective photosensitizer **158** (BOPP) which used in photodynamic therapy of cancer and is believed to be a potential agent for boron neutron capture therapy of brain tumors [22, 26, 108]. The ability of BOPP to accumulate in tumor cells was demonstrated by experiments *in vivo* (the ratio brain tumor-surrounding intact tissue exceeded 400:1) [26].

In some cases, modification of hydroxy groups leads to unexpected results. Attempts to obtain methanesulfonate derivative **159** by reaction of photoporphyrin **160** with MeSO₂Cl were unsuccessful; the product was 3-(1-chloro-2-oxoethyl)porphyrin **161** [79] (Scheme 26). The reduction of photoporphyrin **163** gave diol **162** whose treatment with H₅IO₆ promoted a series of rearrangements leading to formylporphyrin **19** [82].


Scheme 25.



Scheme 26.



Scheme 27.



3.2.2. Oxidation. The aldehyde moiety in protoporphyrin IX derivatives can be converted into carboxy group via oxidation with CrO_3 in Me_2CO [2, 33] or with atmospheric oxygen in the presence of HI [33]. The oxidation of diformylporphyrin **24** according to Jones, followed by esterification, was performed to introduce acetic acid residues into the 3- and 8-positions of protoporphyrin IX. The resulting tetraester **164** is the key intermediate in the synthesis of porphyrin *d*₁ tetramethyl ester (**165**) (Scheme 27). The corresponding tetracarboxylic acid is the ligand in heme *d*₁ (**166**) which plays an important role in the metabolism of sulfur and nitrogen in various organisms [2].

3.2.3. Nucleophilic addition to formyl groups. Formyl groups in protoporphyrin IX derivatives readily react with various nucleophiles. In this section, nucleophilic addition reactions leading to formation of multiple bonds and those not involving formation of multiple bonds are considered.


3.2.3.1. Synthesis of imino derivatives. Heating of formylporphyrins with hydroxylamine hydrochloride or hydroxylamine ethers give the corresponding oximes in high yields [9, 33, 106, 109, 110]. Isophotoporphyrin **167** is thus converted into oxime **168** [110] which is the key intermediate in the synthesis of ATX-S10 (**169**) used for sonodynamic therapy of cancer [111] (Scheme 28). Formylporphyrins also

undergo other transformations typical of common aldehydes [33, 82, 112]. By heating of formyl derivative **167** with *tert*-butylamine unstable Schiff base **170** was obtained [82].

Modification of formyl derivatives of protoporphyrin IX was studied with a view to integrate photodynamic capabilities of tetrapyrroles with pharmacophoric potential of an α -aminophosphonate fragment [74, 82]. Unlike common α -aminophosphonates, syntheses involving natural porphyrins and the model system $(\text{EtO})_2\text{P}(\text{O})\text{H}-t\text{-BuNH}_2$ may be effected only under conditions of microwave activation, especially in combination with catalysis by CdI_2 [82]. Formylporphyrin **167** was thus regioselectively converted into aminophosphonate **171** in 85% yield [82]. This transformation is the first example of microwave-assisted modification of porphyrin structures.


The well known radiosensitizing effect of nitroimidazoles stimulated studies aimed at combining nitroimidazole and porphyrin fragments in a single molecule [109]. Nitroimidazolyl-containing porphyrin manganese(III) complex **172** was successfully synthesized by condensation of oxime **173** with contrast agent **174** in MeOH in the presence of triethylamine [109] (Scheme 29). Dicarboxylic acid **175** was shown by experiments *in vivo* to be promising reagent for cancer diagnostics and therapy [22, 109].

Scheme 28.




169, Asp = $\text{HOCOCH}_2\text{CH}(\text{COOH})\text{NH}_2$.

Scheme 29.



172, 175, M = Mn(III).

Scheme 30.




3.2.3.2. Synthesis of vinyl derivatives. The Wittig reaction was used most frequently to convert formyl groups in protoporphyrin IX derivatives into ethylidene moiety [9, 43, 44, 58, 100, 113–115]. Some classical syntheses also involved the Knoevenagel reaction

[9, 33]. Protoporphyrins labeled with ^{13}C were synthesized by the Wittig reaction with $\text{Ph}_3\text{P}=\text{CH}_2^{13}\text{C}$ [43, 44]. The condensation of diformylporphyrin **176** with $\text{Ph}_3\text{P}=\text{CHCO}_2\text{CH}_3$ was the key stage in a practical synthesis of coproporphyrin III tetramethyl ester (**177**)

[44] (Scheme 30). Acrylate **178** obtained by the Wittig reaction in 96% yield was reduced with hydrogen over Pd/C and then treated with diazomethane; the yield of tetraester **177** was 80% [44]. A convenient procedure for the synthesis of fluoro-substituted protoporphyrin IX derivatives is based on reactions with fluorinated phosphorane generated *in situ* [58, 115, 116].


Fluorine-containing porphyrins **179–181** were synthesized in good yields by heating formylporphyrins **19**, **20**, and **22** with triphenylphosphine and sodium dichlorofluoroacetate in *N*-methylpyrrolidin-2-one [58] (Scheme 31). It was proposed to use porphyrine **181** as starting material in the preparation of difluorovinyl analog of protohemin IX (**12**) [117]. *In vitro* assays of

Scheme 31.




19, $R^1 = \text{CHO}$, $R^2 = \text{CH}_2=\text{CH}$; **20**, $R^1 = \text{CH}_2=\text{CH}$, $R^2 = \text{CHO}$; **22**, $R^1 = R^2 = \text{CHO}$; **179**, **182**, $R^1 = \text{CF}_2=\text{CH}$, $R^2 = \text{CH}_2=\text{CH}$; **180**, **183**, $R^1 = \text{CH}_2=\text{CH}$, $R^2 = \text{CF}_2=\text{CH}$; **181**, **184**, $R^1 = R^2 = \text{CF}_2=\text{CH}$.

Scheme 32.



Scheme 33.







disodium salts **182–184** derived from compounds **179–181** showed that 8-substituted derivative **183** accumulates mainly in stomach tumor cells, and 3,8-disubstituted compound **184**, in hepatoma cells, while 3-substituted analog **182** turned out to be weakly active [58].

A spirochlorin–chlorin dimer as potential model for the “special pair” of the photosynthetic reaction center was obtained via reaction of chlorin **167** with diphosphonium salt **185** [113, 114]. In the presence of DBU,

bis- and monochlorin derivatives **186** and **187** were formed in 60 and 20% yield; prolonged reaction (24 h) resulted in the transformation of bis-chlorin **186** into spirochlorin–chlorin dimer **188** (Scheme 32). The mechanism of formation of dimer **188** is likely to include intramolecular Diels–Alder reaction involving the double bond system between the chlorin fragments in initial dimer **186** [114]. Dimer **188** was converted into porphyrin–spirochlorin derivative **189** on storing

Scheme 34.



Scheme 36.**Scheme 37.****Scheme 38.**

211, L = Na⁺, K⁺; **212**, L = NH₄⁺.

COO–Nb bonds between the porphyrin and Nb₂O₅. Further modification of electrodes by the above system led to creation of an effective sensor for determination of dissolved oxygen [14]. An elegant procedure for modification of carboxy groups in porphyrins was proposed in [51, 84, 124]. Porphyrins **210** reacted with the platinum complex [Pt(NH₃)₂(H₂O)₂](OH)₂ in aqueous ethanol to produce coordination compounds **212** (Scheme 38) which showed high cyto- and photocytotoxicity [51].

3.3.3. Different versions of esterification of 13- and 17-(2-carboxyethyl) groups in porphyrins. Carboxy groups in porphyrins are converted into methoxycarbonyl by treatment with diazomethane or a 5% solution of H₂SO₄ in methanol [33]. Methods intrinsic to peptide chemistry are applied to obtain esters containing labile fragments. Variation of the alcohol component allows preparation of derivatives that may be promising for antitumor therapy [125–128], studying photosynthesis [129–131], modification of various proteins [132], and application in technics [13, 15, 123, 133].


Diester **214a** was synthesized in 75% yield by condensation of protoporphyrin IX (**1**) with alcohol **213** in the presence of *N,N'*-dicyclohexylcarbodiimide and 4-dimethylaminopyridine [125] with a view to combine anticarcinogenic activity of the anthraquinone derivative and photodynamic properties of tetrapyrroles. Murakami et al. [132] synthesized porphyrin–fullerene diads **214b** and **215** as unique compounds for modification of myoglobins and studied photophysical

properties of the modified proteins. The synthesis involved activation of the carboxy groups in protoporphyrin IX (**1**) with oxalyl chloride, condensation with fullerene derivatives **216** in the presence of 4-dimethylaminopyridine, and metalation (Scheme 39).

3.3.4. Amide derivatives based on 13,17-bis(2-carboxyethyl)porphyrins. Synthesis of amide derivatives of protoporphyrin IX is important from the viewpoint of design of compounds that may be used in biology [129, 134–142] and medicine [20, 21, 52, 80, 110–112, 126, 128, 130, 131, 135, 143–150]. Effective antitumor agent **216** was synthesized by successive treatment of protoporphyrin IX (**1**) with ethyl chloroformate, ethylenediamine, succinimidopoly(ethylene glycol) (*M* 5000), and zinc(II) acetate [147]. Evstigneeva et al. [135] reported on conjugation of protohemin IX (**12**) with peptides to obtain porphyrin-modified peptides **217** as convenient models for studying heme-containing proteins (Scheme 40).


Some amides were prepared from carboxy-containing porphyrins without involving peptide chemistry methods. Rogovina et al. [151] performed solid-phase amidation of carboxy groups in deuteroporphyrin derivatives using chitosan under high pressure (3 GPa). Liang et al. [138] described a successful synthesis of diamide derived from protoporphyrin IX zinc complex by reaction of the corresponding dimethyl ester with aluminum amide. An example of hydrazide modification of carboxy groups in protoporphyrin IX derivatives for conjugation with immunotoxins was also reported [152].

Scheme 39.



214b, *M* = Zn; **215**, *M* = [FeCl]²⁺.

Scheme 40.



$\mathbf{216}$, $\mathbf{R}$ = poly(ethylene glycol)-OC(O)(CH₂)₂C(O)NH(CH₂)₂.

3.4. Modification of Methyl Groups

Reactions of protoporphyrin IX and its derivatives accompanied by transformations of methyl groups are fairly rare. Electrophilic deuteration of methyl groups by heating protoporphyrin IX dimethyl ester (**18**) in solvents containing MeOD and MeO[−] was reported [9]. Mironov et al. [153] performed chlorination of methyl groups by prolonged reaction of porphyrin metal complexes with thionyl chloride.

4. MODIFICATION OF MACRORING

The structure of protoporphyrin IX (**1**) suggests that the macroring therein can be modified at the *meso* positions (5, 10, 15, 20), with participation of 2–3, 7–8, 12–13, and 17–18 double bonds, and via coordination with metals. Some examples of macroring identification implied rearrangements involving peripheral substituents. Data on such modifications of the macroring in porphyrins are briefly discussed below.


4.1. Modifications at *meso* Positions

Transformations involving *meso* positions in the macroring of protoporphyrin IX derivatives are classical reactions in the porphyrin chemistry; they were considered in detail in some reviews [60, 95]. In the recent years, modifications at the *meso* positions were performed to obtain derivatives interesting from the viewpoint of studying metabolism of heme [45, 154–157]. Formylation of copper complexes of mesoporphyrin IX (**218**) [154] and protected protoporphyrin IX (**219**) [45] with POCl₃–DMF gave mixtures of the corresponding *meso*-formylporphyrin complexes **220–223** and **224–227** [45, 154] (Scheme 41). The subsequent demetalation of copper complexes **224–227** with 5% H₂SO₄ in CF₃COOH and deprotection of vinyl groups with 30% H₂O₂ in THF afforded a mixture of

meso-formyl-substituted protoporphyrin IX analogs **228–231** which were separated by HPLC [45]. Successive treatment of copper complexes **220–223** with Bu₄NBH₄ and H₂SO₄–CF₃COOH (1:1) resulted in the formation of isomeric *meso*-hydroxymethyl derivatives **232–235** which were separated by column chromatography [154]. The exact position of the *meso*-substituent in individual isomers **228–235** was determined using NOE technique [45, 154]. In early studies, the structure of mesoporphyrin IX formylation products was determined by their transformation into the corresponding *meso*-dimethylaminomethylporphyrins **236–243** and borane complexes **244–251** [60] (in order to simplify interpretation of the ¹H NMR spectra of **236–251**, the corresponding analogs containing a deuterium label in the 13- and 17-CH₂CH₂COOCH₃ substituents were obtained). Reduction of *meso*-hydroxymethylporphyrins **232–235** gave a set of *meso*-methyl derivatives **252–255**, and the latter were subjected to hydrolysis followed by metalation to obtain four regioisomeric *meso*-methylmesohemins **256–259** [154]. Individual *meso*-formylmesohemins **260–263** were synthesized by oxidation of hydroxymethyl derivatives **232–235** with pyridinium chlorochromate in pyridine, and aldehydes **264–267** thus formed were hydrolyzed and metalated [157].


Apart from formylation, successful nitration of deuteroporphyrin IX dimethyl ester (**119**) with a mixture of nitric and sulfuric acids to obtain dinitro derivative **268** [51] and chlorination at the *meso* positions of protoporphyrin IX metal complexes with SOCl₂ [153] were reported. Vázquez et al. [158] synthesized tribenzoylporphyrin **269** by acylation of zinc complex **122** with benzoic anhydride in the presence of SnCl₄ (Scheme 42). Hemins **273–275** were treated with zinc amalgam to obtain hemes **270–272**, and reactions of the latter with hydrogen peroxide in pyridine in the absence of oxygen were studied to simulate heme

Scheme 41.




218, $R^1 = R^2 = \text{H}$; **220**, **228**, **260**, **264**, $R^1 = \text{CHO}$, $R^2 = R^3 = R^4 = \text{H}$; **221**, **229**, **261**, **265**, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{CHO}$; **222**, **230**, **262**, **266**, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{CHO}$; **223**, **231**, **263**, **267**, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{CHO}$; **224**, **228**, $R^1 = \text{CHO}$, $R^2 = R^3 = R^4 = \text{H}$; **225**, **229**, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{CHO}$; **226**, **230**, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{CHO}$; **227**, **231**, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{CHO}$; **232**, $R^1 = \text{HOCH}_2$, $R^2 = R^3 = R^4 = \text{H}$; **233**, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{HOCH}_2$; **234**, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{HOCH}_2$; **235**, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{HOCH}_2$; **236**, $M = \text{Ni(II)}$, $R^1 = \text{CH}_2\text{NMe}_2$, $R^2 = R^3 = R^4 = \text{H}$; **237**, $M = \text{Ni(II)}$, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{Me}_2\text{NCH}_2$; **238**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{Me}_2\text{NCH}_2$; **239**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{Me}_2\text{NCH}_2$; **240**, $M = 2\text{H}$, $R^1 = \text{Me}_2\text{NCH}_2$, $R^2 = R^3 = R^4 = \text{H}$; **241**, $M = 2\text{H}$, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{Me}_2\text{NCH}_2$; **242**, $M = 2\text{H}$, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{Me}_2\text{NCH}_2$; **243**, $M = 2\text{H}$, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{Me}_2\text{NCH}_2$; **244**, $M = \text{Ni(II)}$, $R^1 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$, $R^2 = R^3 = R^4 = \text{H}$; **245**, $M = \text{Ni(II)}$, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **246**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **247**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **248**, $M = 2\text{H}$, $R^1 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$, $R^2 = R^3 = R^4 = \text{H}$; **249**, $M = 2\text{H}$, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **250**, $M = 2\text{H}$, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **251**, $M = 2\text{H}$, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{BH}_3 \cdot \text{Me}_2\text{NCH}_2$; **252**, **256**, $M = \text{Ni(II)}$, $R^1 = \text{Me}$, $R^2 = R^3 = R^4 = \text{H}$; **253**, **257**, $M = \text{Ni(II)}$, $R^1 = R^3 = R^4 = \text{H}$, $R^2 = \text{Me}$; **254**, **258**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^4 = \text{H}$, $R^3 = \text{Me}$; **255**, **259**, $M = \text{Ni(II)}$, $R^1 = R^2 = R^3 = \text{H}$, $R^4 = \text{Me}$.


Scheme 44.



Scheme 45.



Scheme 46.



tate and then with acetic anhydride gave oxonium ion **308**. The final stage was amination of **308** with ammonia, followed by treatment with trimethylsilyl polyphosphate.


4.2. Modifications Involving 2–3, 7–8, 12–13, and 17–18 Double Bonds.


The porphyrin double bonds in positions 2–3, 7–8, 12–13, and 17–18 are oxidized with OsO_4 to form chlorin structures with vicinal diol fragments [36, 70, 163]; the oxidation of unsymmetrical octaalkylporphyrin systems leads to the formation of mixtures of isomeric vicinal diols [70, 163, 164]. Regioselective oxidation may be achieved with unsymmetrical tetrapyrroles containing a peripheral electron-withdrawing substituent. In this case, the reaction involves only one double bond in the pyrrole ring located opposite to the pyrrole ring containing electron-withdrawing substit-

uent [70]. For example, the oxidation of 3-acetyldeuteroporphyrin **309** gives 68% of 12,13-dihydroxychlorin **310** [70] (Scheme 47). Vicinal dihydroxychlorins are converted into oxochlorins under conditions of pinacol–pinacolone rearrangement (concentrated sulfuric acid). Chlorin **310** gives rise to a mixture of isomeric chlorins **311** and **312** [70]. Taking into account concomitant migration of peripheral substituents, the formation of oxochlorins is a fairly complex process (for details, see [70, 163]). Vicinal diols are capable of undergoing intramolecular lactonization. Heating of a solution of diol **310** in methanol in the presence of sodium acetate afforded 64% of *cis*-lactone **313** which underwent epimerization over silica gel to give *trans*-lactone **314** [70] (Scheme 47).

A combination of oxidation of the 2–3, 7–8, 12–13, and 17–18 double bonds in porphyrin with subsequent pinacol–pinacolone rearrangement, repetition of this

Scheme 47.





procedure, and some additional transformations are often performed to determine the structure of some biologically important highly saturated tetrapyrrole compounds [2].



An efficient procedure for regioselective transformation of a porphyrin system via oxidation of the 2–3 and 7–8 double bonds was proposed in [165]. It was found [166] that incubation of deuterioporphyrin IX (**3**) with horse radish peroxidase in the presence of glutathione gives isomeric chlorins **315** and **316** [165] (Scheme 48). Under analogous conditions protoporphyrin IX (**1**) was converted into an unidentified derivative [167]. Factor affecting the formation of chlorins **315** and **316** were discussed in detail in [168].

4.3. Metal Complexes and Their Demetalation

The ability of porphyrins to form coordination compounds with metals is a fundamental property of tetrapyrrole systems. Detailed analysis of the possibility for complex formation with porphyrins, stability of the complexes thus formed, and specificity of their demetalation was given in [32, 169, 170]. The recent data concerning these problems are briefly discussed below. A conventional procedure for the synthesis of porphyrin metal complexes is heating of free bases with metal salts in polar solvents [27, 51, 98, 106, 122, 144, 171, 172]. Reactions with radioactive metal isotopes open the way to radiopharmaceutical agents with a broad spectrum of applications in medicine [22, 28].

Platinum porphyrin complexes are characterized by a high luminescence quantum yield and are promising as luminescent probes. Platinum complex **317** possesses valuable photophysical properties and is capable of selectively accumulating in tumor cells [27]; it was patented as potential agent for luminescent diagnostics of cancer [19]. Complex **317** was synthesized by heating free base **177** with K_2PtCl_4 in benzonitrile, followed by alkaline hydrolysis of the ester groups [27] (Scheme 49). Treatment of dihydrochloride **318** with $SnCl_2$ in the presence of oxygen on heating gave

tin(IV) complex **319** (Stannosoporphin) in a good yield; the product is effective in the treatment of physiologic jaundice and psoriasis [40] (Scheme 50). Ishida et al. [173] recently demonstrated prospects in using template catalysis by apocytochrome b_{562} for metalation of porphyrins. Iron(III) complex **12** is widely used as starting compound in the chemistry of protoporphyrin IX. Complex **12** is isolated on a large scale by treatment of defibrinated bovine blood. Specificity of demetalation of protohemin IX (**12**) and other iron(III) porphyrin complexes was reviewed in detail in [9, 33].




318, $X = N \cdot HCl$; **319**, $M = [SnCl_2]^{2+}$.


4.4. Modifications Involving Rearrangement Processes

Variation of initial tetrapyrrole structures and conditions makes it possible to obtain tetrapyrrole systems with various modes of conjugation; transformations of partially saturated structures to porphyrins and vice versa are possible. Some aspects of the transformation of chlorins into thermodynamically more stable por-


Scheme 51.



Scheme 52.



Scheme 53.



pyrin derivatives were considered above (see Section 3.2). There are published data on double dehydration of vicinal diols obtained by oxidation of double bonds with OsO_4 ; here, vinylporphyrin fragment is formed as a result of successive treatment with hydrochloric acid and $\text{H}_2\text{SO}_4\text{-MeOH}$. The corresponding transformation of porphyrin **d**₁ tetramethyl ester **165** [2] (Scheme 51).

Examples of transformation of the chlorin system into dihydrobenzoporphyrin are known. Treatment of isophotoporphyrin **167** with phosphonium salt **321** in the presence of DBU gives the expected Wittig

reaction product **322**, and subsequent heating of compound **322** in 1,2-dichlorobenzene leads to elimination of water molecule and isomerization of intermediate chlorin **323** into stable dihydrobenzoporphyrin **324** [114] (Scheme 52). Transformation of the porphyrin system to chlorin is also possible. Syntheses of chlorin derivatives from vinylporphyrins via Diels–Alder reaction were discussed in Section 3.2. Diels–Alder reactions were also used to convert vinylporphyrins into benzoporphyrins [36] and isobacteriochlorins [9].

Montforts et al. [174, 175] described a new method for building up chlorin system with participation of

hydroxyethyl groups in porphyrins. The Claisen reaction of 8-(1-hydroxyethyl) derivative **325** with *N,N*-dimethyl-1,1-dimethoxyethanamine on heating afforded chlorin **326** through intermediate porphyrin **327** [174] (Scheme 53). Further modifications of chlorin **326** were also performed [174, 175]. For example, oxidation of **326** gave ketone **328** which was reduced with $\text{LiAlH}(\text{O}i\text{Bu})_3$ and then treated with a base to obtain lactone **329** [175].

5. CONCLUSION

Vast experimental data on modifications of protoporphyrin IX and its derivatives have been published over the past decade. The presence in the molecules of these compounds of various peripheral substituents (such as vinyl, formyl, and carbonyl groups) makes it possible to transform the porphyrin macroring into other tetrapyrrole systems, and the ability of tetrapyrroles to form stable complexes with many metals opens wide prospects in their modification using methods of organic and bioorganic chemistry.

The main recent trend in the chemistry of protoporphyrin IX is the application of flexible organic chemistry methods with a view to obtain derivatives possessing practically important properties. In particular, a combination of regioselective introduction of pharmacophoric fragments as peripheral substituents and variation of the central metal ion could give rise to effective agents for diagnostics and therapy of various diseases with the aid of powerful modern photomedical, radiological, X-ray, nuclear magnetic resonance, and radioisotope methods. Low systemic toxicity, valuable photophysical properties, and the ability to selectively accumulate in tumor and other affected cells make protoporphyrin IX derivatives especially promising.

The information collected in the present review on methods of modification of protoporphyrin IX and its derivatives demonstrates their huge synthetic potential and wide prospects in the development of synthetic approaches to compounds possessing a broad spectrum of useful properties, primarily valuable biological activity.

The author thanks Dr. M.S. Nechaev (Faculty of Chemistry, Moscow State University, Moscow, Russia), Dr. A.S. Lermontov (Topchiev Institute of Petrochemical Synthesis, Russian Academy of Sciences, Moscow, Russia), Dr. S.V. Shorunov (Zelinskii Institute of Organic Chemistry, Russian Academy of

Sciences, Moscow, Russia), I.O. Konstantinov (Research Institute of Chemical Diversity Ltd., Khimki, Russia), "First Print Yard" joint-stock company (Moscow, Russia), and personally O.I. Konstantinov for their assistance in the preparation of the present review. The review was prepared under financial support by the Russian Foundation for Basic Research (project no. 04-03-32710).

REFERENCES

1. Dawson, R.M.C., Elliott, D.C., Elliot, W.H., and Jones, K.M., *Data for Biochemical Research*, Oxford: Clarendon, 1986.
2. Montforts, F.-P., Gerlach, B., and Hoper, F., *Chem. Rev.*, 1994, vol. 94, p. 327.
3. Battersby, A.R., *Nat. Prod. Rep.*, 2000, vol. 17, p. 507.
4. Kundu, S., Premer, S.A., Hoy, J.A., Trent, J.T., III, and Hargrove, M.S., *Biophys. J.*, 2003, vol. 84, p. 3931.
5. Sono, M., Roach, M.P., Coulter, E.D., and Dawson, J.H., *Chem. Rev.*, 1996, vol. 96, p. 2841.
6. Shaik, S., de Visser, S.P., Ogliaro, F., Schwarz, H., and Schroeder, D., *Curr. Opin. Chem. Biol.*, 2002, vol. 6, p. 556.
7. Guallar, V., Baik, M.H., Lippard, S.J., and Friesner, R.A., *Proc. Natl. Acad. Sci. USA*, 2003, vol. 100, p. 6998.
8. *The Porphyrin Handbook*, Kadish, K., Smith, K.M., and Guillard, R., Boston: Academic, 2000, vol. 6.
9. Smith, K.M. and Cavaleiro, J.A.S., *Heterocycles*, 1987, vol. 26, p. 1947.
10. Woo, I., Hwang, M.W., and Cho, J.G., *Kayaku Gakkaishi*, 1996, vol. 57, p. 71; *Chem. Abstr.*, 1996, vol. 125, no. 122420.
11. Brecht, R., Buettner, F., Boehm, M., Seitz, G., Frenzen, G., Pilz, A., and Massa, W., *J. Org. Chem.*, 2001, vol. 66, p. 2911.
12. Smith, A.B., Kanoh, N., Ishiyama, H., Minakawa, N., Rainier, J.D., Hartz, R.A., Cho, Y.S., Cui, H., and Moser, W.H., *J. Am. Chem. Soc.*, 2003, vol. 125, p. 8228.
13. Cosnier, S., Gondran, C., Wessel, R., Montforts, F.-P., and Wedel, M., *J. Electroanal. Chem.*, 2000, vol. 488, p. 83.
14. Pessôa, C.A. and Gushikem, Y., *J. Porphyrins Phthalocyanines*, 2001, vol. 5, p. 537.
15. Cosnier, S., Gondran, C., Wessel, R., Montforts, F.-P., and Wedel, M., *Sensors*, 2003, vol. 3, p. 213.
16. Wedel, M. and Montforts, F.-P., *Tetrahedron Lett.*, 1999, vol. 40, p. 7071.
17. Montforts, F.-P., Vlasiouk, I., Smirnov, S., and Wedel, M., *J. Porphyrins Phthalocyanines*, 2003, vol. 7, p. 651.

18. Tsuchida, E., Komatsu, T., Nakagawa, A., and Ohmichi, N., US Patent Appl. Publ. no. 20040180837, 2004; *Chem. Abstr.*, 2004, vol. 141, no. 282880.
19. Sukhin, G.M., Chissov, V.I., Rumyantseva, V.D., Mironov, A.F., Pecherskikh, E.V., and Menenkov, V.D., Russian Patent no. 2074718, 1997; *Chem. Abstr.*, 1997, vol. 127, no. 158774.
20. Sternberg, E.D., Dolphin, D., and Bruckner, C., *Tetrahedron*, 1998, vol. 54, p. 4151.
21. Reshetnikov, A.V., Shvets, V.I., and Ponomarev, G.V., *Uspekhi khimii porfirinov* (Advances in Porphyrin Chemistry), St. Petersburg: Nauch.-Issled. Inst. Khimii Sankt-Peterb. Gos. Univ., 1999, vol. 2, p. 70.
22. Ali, H. and van Lier, J.E., *Chem. Rev.*, 1999, vol. 99, p. 2379.
23. Mody, T.D., *J. Porphyrins Phthalocyanines*, 2000, vol. 4, p. 362.
24. Pandey, R.K., *J. Porphyrins Phthalocyanines*, 2000, vol. 4, p. 368.
25. MacDonald, I. and Dougherty, T.J., *J. Porphyrins Phthalocyanines*, 2001, vol. 5, p. 105.
26. Bregadze, V.I., Sivaev, I.B., Gabel, D., and Wöhrle, D., *J. Porphyrins Phthalocyanines*, 2001, vol. 5, p. 767.
27. Bykhovskii, V.Ya., Zaitseva, N.I., Mironov, A.F., Osin, N.S., Pecherskikh, E.V., Rumyantseva, V.D., and Sukhin, G.M., *Prikl. Biokhim. Mikrobiol.*, 2001, vol. 37, p. 660.
28. Adair, E.L., US Patent no. 7067276, 2004; *Chem. Abstr.*, 2004, vol. 141, no. 327715.
29. Fischer, H. and Orth, H., *Die Chemie des Pyrrols*, Leipzig: Akademische Verlagsgesellschaft, 1934, vol. 1.
30. Fischer, H. and Orth, H., *Die Chemie des Pyrrols*, Leipzig: Akademische Verlagsgesellschaft, 1937, vol. 2, part 1.
31. Fischer, H. and Stern, A., *Die Chemie des Pyrrols*, Leipzig: Akademische Verlagsgesellschaft, 1940, vol. 2, part 2.
32. Falk, J.E., *Porphyrins and Metalloporphyrins*, Amsterdam: Elsevier, 1964.
33. *Porphyrins and Metalloporphyrins*, Smith, K.M., Ed., Amsterdam: Elsevier, 1975.
34. Dolphin, D., *The Porphyrins*, New York: Academic, 1978.
35. Shanmugathan, S., Edwards, C., and Boyle, R.W., *Tetrahedron*, 2000, vol. 56, p. 1025.
36. Vicente, M.G.H. and Smith, K.M., *Curr. Org. Chem.*, 2000, vol. 4, p. 139.
37. Smith, K.M. and Vicente, M.G.H., *Science of Synthesis*, Weinreb, S.M., Ed., Stuttgart: Thieme, 2003, vol. 17, p. 1081.
38. *The Porphyrin Handbook*, Kadish, K., Smith, K.M., and Guillard, R., Eds., Boston: Academic, 2000, vols. 1–10.
39. *The Porphyrin Handbook*, Kadish, K., Smith, K.M., and Guillard, R., Eds., San Diego: Academic, 2003, vols. 11–20.
40. Vukovich, R., Levinson, B., Drummond, G.S., Caroselli, R., Antczak, K.G., Boucher, C., Mortimer, R., Levin, D., and Cooke, K.A., US Patent Appl. Publ. no. 20040210048, 2004; *Chem. Abstr.*, 2004, vol. 141, no. 359890.
41. Peixoto, A., Pereira, M.M., Neves, M., Graca, P.M.S., Silva, A.M.S., and Cavaleiro, J.A.S., *Tetrahedron Lett.*, 2003, vol. 44, p. 5593.
42. Peixoto, A.F., Pereira, M.M., Sousa, A.F., Pais, A.A.C., Neves, M.G.P.M.S., Silva, A.M.S., and Cavaleiro, J.A.S., *J. Mol. Catal. A: Chem.*, 2005, vol. 235, p. 185.
43. Woo, P.W.K., Hayes, R., Beylin, V.G., Johnson, S., Lovdahl, M., Huang, Y., and Huang, C.C., *Proc. 6th Int. Symp.*, Philadelphia, PA, USA, 1998, p. 429.
44. Kahl, S.B., Schaeck, J.J., and Koo, M.-S., *J. Org. Chem.*, 1997, vol. 62, p. 1875.
45. Brantley, S.E., Gerlach, B., Olmstead, M.M., and Smith, K.M., *Tetrahedron Lett.*, 1997, vol. 38, p. 937.
46. Sylvain, I., Zerrouki, R., Granet, R., Huang, Y.M., Lagorce, J.-F., Guilloton, M., Blais, J.-C., and Krausz, P., *Bioorg. Med. Chem.*, 2002, vol. 10, p. 57.
47. Brunner, H., Obermeier, H., and Sziemies, R.-M., *Chem. Ber.*, 1995, vol. 128, p. 173.
48. Pilloud, D.L., Moser, C.C., Reddy, S., and Dutton, P.L., *Langmuir*, 1998, vol. 14, p. 4809.
49. Pilloud, D.L., Chen, X., Dutton, P.L., and Moser, C.C., *J. Phys. Chem. B*, 2000, vol. 104, p. 2868.
50. Razzaque, M.A., Lord, G.A., and Lim, C.K., *Rapid Commun. Mass Spectrom.*, 2002, vol. 16, p. 1675.
51. Brunner, H., Schellerer, K.-M., and Treitinger, B., *Inorg. Chim. Acta*, 1997, vol. 264, p. 67.
52. Karagianis, G. and Reiss, J.A., *Aust. J. Chem.*, 1995, vol. 48, p. 1693.
53. Ivanov, A.V., Gradyushko, A.T., Laptev, V.P., Panferova, N.G., Varlamov, V.P., Klyashchitsky, B.A., Reshetnikov, A.V., and Ponomarev, G.V., *Proc. SPIE, The International Society for Optical Engineering*, 1996, vol. 2728, p. 181; *Chem. Abstr.*, 1996, vol. 124, no. 219268.
54. Fickweiler, S., Szeimies, R.-M., Abels, C., Ponomarev, G.V., Hofstadter, F., Wolfbeis, O.S., and Landthaler, M., *Photoderm. Photoimmun. Photomed.*, 1998, vol. 14, p. 125.
55. Nakae, Y., Sakata, I., Nakajima, S., Hidege, N., and Aburano, T., JPN Patent Appl. no. 2002205959, 2002; *Chem. Abstr.*, 2002, vol. 137, no. 105844.
56. Tian, X.-L., Sun, H., and Wang, Z.-J., *Zhongguo Yiyao Gongye Zazhi*, 1998, vol. 29, p. 401; *Chem. Abstr.*, 1999, vol. 130, no. 52255.
57. Pandey, R.K., Dougherty, T.J., Smith, K.M., and Meunier, I., US Patent no. 5498710, 1996; *Chem. Abstr.*, 1996, vol. 125, no. 10486.

58. Ando, A. and Kumadaki, I., *J. Fluorine Chem.*, 1999, vol. 100, p. 135.
59. Omote, M., Matsumoto, T., Ando, A., Koyama, M., Takagi, T., and Kumadaki, I., *Heterocycles*, 1997, vol. 46, p. 259.
60. Ponomarev, G.V., *Khim. Geterotsikl. Soedin.*, 1997, p. 1299.
61. Ponomarev, G.V., Kirillova, G.V., and Yashunskii, D.V., *Khim. Geterotsikl. Soedin.*, 2000, p. 1197.
62. Isaac, M.F. and Kahl, S.B., *J. Organomet. Chem.*, 2003, vol. 680, p. 232.
63. Reshetnikov, A.V., Babushkina, T.A., Kirillova, G.V., and Ponomarev, G.V., *Khim. Geterotsikl. Soedin.*, 2001, p. 213.
64. Ponomarev, G.V., Pavlov, V.Yu., Konstantinov, I.O., Timofeev, V.P., and Kimel', B.G., *Russ. J. Org. Chem.*, 2003, vol. 39, p. 1683.
65. Pavlov, V.Yu., *Cand. Sci. (Chem.) Dissertation*, Moscow, 2006.
66. Uzdensky, A.B., Ivanov, A.V., Reshetnikov, A.V., Ponomarev, G.V., Dergacheva, O.Y., and Zhavoronkova, A.A., *Proc. SPIE, The International Society for Optical Engineering*, 2000, vol. 4059, p. 147.
67. Mansurova, G.V., Pogrebnaya, O.G., Ponomarev, G.V., Reshetnikov, A.V., Potapenko, A.Ya., Bezdetnaya, L.N., and Guillemin, F., *Internet Photochem. Photobiol.*, 2000; <http://www.photobiology.com/photobiology2000/mansurova/index>.
68. Uzdensky, A.B., Dergacheva, O.Yu., Zhavoronkova, A.A., Ivanov, A.V., Reshetnikov, A.V., and Ponomarev, G.V., *Biochem. Biophys. Res. Commun.*, 2001, vol. 281, p. 1194.
69. Mansurova, G.V., Pogrebnaya, O.G., Ponomarev, G.V., Reshetnikov, A.V., Potapenko, A.Ya., Bezdetnaya, L.N., and Gulya, F., *Biofizika*, 2003, vol. 48, p. 251.
70. Pandey, R.K., Isaac, M., MacDonald, I., Medforth, C.J., Senge, M.O., Dougherty, T.J., and Smith, K.M., *J. Org. Chem.*, 1997, vol. 62, p. 1463.
71. Shigeoka, T., Kuwahara, Y., Watanabe, K., Sato, K., Omote, M., Ando, A., and Kumadaki, I., *Chem. Pharm. Bull.*, 1999, vol. 47, p. 1326.
72. Kusch, D. and Montforts, F.-P., *Tetrahedron: Asymmetry*, 1995, vol. 6, p. 867.
73. Kabachnik, M.M., Zobnina, E.V., and Beletskaya, I.P., *Synlett*, 2005, p. 1393.
74. Kabachnik, M.M., Zobnina, E.V., Pavlov, V.Yu., Konstantinov, I.O., Ponomarev, G.V., and Beletskaya, I.P., *Izv. Ross. Akad. Nauk, Ser. Khim.*, 2005, p. 256.
75. Chakrabarty, M., *J. Indian Chem. Soc.*, 2001, vol. 78, p. 761.
76. Jiang, X. and Smith, K.M., *J. Chem. Soc., Perkin Trans. 1*, 1996, p. 1601.
77. Morgan, A.R. and Kohli, D.H., *Tetrahedron*, 1995, vol. 36, p. 7603.
78. Grandadam, M., Ingrand, D., Huraux, J.-M., Aveline, B., Delgado, O., Vever-Bizet, C., and Brault, D., *J. Photochem. Photobiol. B: Biol.*, 1995, vol. 31, p. 171.
79. Iakovides, P. and Smith, K.M., *Tetrahedron*, 1996, vol. 52, p. 1123.
80. Bourtine, A.S., Brault, D., Takasugi, M., Delgado, O., and Helene, C., *J. Am. Chem. Soc.*, 1996, vol. 118, p. 9469.
81. Brault, D., Aveline, B., Delgado, O., and Martin, M.T., *Photochem. Photobiol.*, 2001, vol. 73, p. 331.
82. Pavlov, V.Yu., Kabachnik, M.M., Zobnina, E.V., Timofeev, V.P., Konstantinov, I.O., Kimel, B.G., Ponomarev, G.V., and Beletskaya, I.P., *Synlett*, 2003, p. 2193.
83. Pavlov, V.Yu., Konstantinov, I.O., Ponomarev, G.V., Timofeev, V.P., and Kimel', B.G., *Russ. J. Org. Chem.*, 2005, vol. 41, p. 1824.
84. Brunner, H. and Schellerer, K.-M., *Monatsh. Chem.*, 2002, vol. 133, p. 679.
85. Monte, W.T., Johnston, A.A., Lindbeck, A.C., and Wang, X.C., Int. Patent no. 9737994, 1997; *Chem. Abstr.*, 1997, vol. 127, no. 346235.
86. Kyagova, A.A., Mansurova, G.V., Kozir, L.A., Ponomarev, G.V., Pavlov, V.Y., Konstantinov, I.O., and Potapenko, A.Ya., *Photochem. Photobiol.*, 2005, vol. 81, p. 1380.
87. Desjardins, A., Flemming, J., Sternberg, E.D., and Dolphin, D., *Chem. Commun.*, 2002, p. 2622.
88. Kozyrev, A.N., Alderfer, J.L., and Robinson, B.C., *Tetrahedron*, 2003, vol. 59, p. 499.
89. Sternberg, E.D., Desjardins, A.M., and Dolphin, D., Int. Patent no. 02076453, 2002; *Chem. Abstr.*, 2002, vol. 137, no. 279028.
90. Fischer, H. and Medick, H., *Justus Liebigs Ann. Chem.*, 1935, vol. 517, p. 245.
91. Jiang, X., Pandey, R.K., and Smith, K.M., *Tetrahedron Lett.*, 1995, vol. 36, p. 365.
92. Jiang, X., Pandey, R.K., and Smith, K.M., *J. Chem. Soc., Perkin Trans. 1*, 1996, p. 1607.
93. Castella, M., Calahorra, F., Sainz, D., and Velasco, D., *Org. Lett.*, 2001, vol. 3, p. 541.
94. Mettich, S., Munson, B.R., and Pandey, R.K., *Bioconjugate Chem.*, 1999, vol. 10, p. 94.
95. Ponomarev, G.V. and Kirillova, G.V., *Porfiriny: struktura, svoystva, sintez* (Porphyrins: Structure, Properties, and Synthesis), Enikolopyan, N.S., Ed., Moscow: Nauka, 1985, p. 238.
96. Gauler, R. and Risch, N., *Eur. J. Org. Chem.*, 1998, p. 1193.
97. Ponomarev, G.V., *Khim. Geterotsikl. Soedin.*, 1994, vol. 30, p. 1669.
98. Rumyantseva, V.D., Konovalenko, L.I., Nagaeva, E.A., and Mironov, A.F., *Bioorg. Khim.*, 2005, vol. 31, p. 103.

99. Omote, M., Ando, A., Tagaki, T., Koyama, M., Kumadaki, I., and Sato, H., *Chem. Pharm. Bull.*, 1995, vol. 47, p. 1107.
100. Ando, A., Yamazaki, M., Komura, M., Sano, Y., Hattori, N., Omote, M., and Kumadaki, I., *Heterocycles*, 1999, vol. 50, p. 913.
101. Sharman, W.M. and van Lier, J.E., *J. Porphyrins Phthalocyanines*, 2000, vol. 4, p. 441.
102. Castella, M., Trull, F.R., Lopez Calahorra, F., Velasco, D., and Gonzalez, M.M., *Tetrahedron*, 2000, vol. 56, p. 4017.
103. Shigeoka, T., Kuwahara, Y., Watanabe, K., Sato, K., Omote, M., Ando, A., and Kumadaki, I., *J. Fluorine Chem.*, 2000, vol. 103, p. 99.
104. Gauler, R., Keuper, R., Winter, A., and Risch, N., *ARKIVOC*, 2004, vol. 13, p. 48; *Chem. Abstr.*, 2005, vol. 142, no. 328283.
105. Liu, X., Sternberg, E., and Dolphin, D., *Chem. Commun.*, 2004, p. 852.
106. Sakata, I., Nakajima, S., Koshimizu, K., Takada, H., and Inui, Y., JPN Patent no. 07-330773, 1995; *Chem. Abstr.*, 1996, vol. 124, no. 289108.
107. Lee, J., Park, Y., Sung, K., Yong, I., and Kang, H.C., *Bull. Korean Chem. Soc.*, 1999, vol. 20, p. 1371.
108. Hill, J.S., Kahl, S.B., Stylli, S.S., Nakamura, Y., Koo, M.-S., and Kaye, A.H., *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, p. 12126.
109. Sakata, I., Nakajima, S., Koshimizu, K., Takada, H., and Inui, Y., JPN Patent no. 08-067682, 1996; *Chem. Abstr.*, 1996, vol. 124, no. 342987.
110. Hikida, M., Mori, M., Sakata, I., Nakajima, S., and Takata, H., Int. Patent no. 9814453, 1998; *Chem. Abstr.*, 1998, vol. 128, no. 282741.
111. Yumita, N., Nishigaki, R., Sakata, I., Nakajima, S., and Umemura, S.-I., *Jpn. J. Cancer Res.*, 2000, vol. 91, p. 255.
112. Nakae, Y., Fukusaki, E.-i., Kajiya, S.-i., Kobayashi, A., Nakajima, S., and Sakata, I., *J. Photochem. Photobiol. A: Chem.*, 2005, vol. 174, p. 187.
113. Zheng, G., Alderfer, J.L., Senge, M.O., Shibata, M., Dougherty, T.J., and Pandey, R.K., *J. Org. Chem.*, 1998, vol. 63, p. 6434.
114. Zheng, G., Shibata, M., Dougherty, T.J., and Pandey, R.K., *J. Org. Chem.*, 2000, vol. 65, p. 543.
115. Shigeoka, T., Kuwahara, Y., Watanabe, K., Sato, K., Omote, M., Ando, A., and Kumadaki, I., *Heterocycles*, 2000, vol. 52, p. 383.
116. Kumadaki, A., Ando, A., and Omote, M., *J. Fluorine Chem.*, 2001, vol. 109, p. 67.
117. Poliart, C., Briand, J.-F., Tortevoie, F., Leroy, J., Simonneaux, G., and Bondon, A., *Magn. Reson. Chem.*, 2001, vol. 39, p. 615.
118. Chen, Z.-L., Wan, W.-Q., Chen, J.-R., Zhao, F., and Xu, D.-Y., *Heterocycles*, 1998, vol. 48, p. 1739.
119. Chen, Z.-L., Chen, J.-R., Wan, W.-Q., and Xu, D.-Y., *Chinese J. Chem.*, 1998, vol. 16, p. 542; *Chem. Abstr.*, 1999, vol. 130, no. 223096.
120. Brunner, H. and Schellerer, K.-M., *Z. Naturforsch., Teil B*, 2002, vol. 57, p. 751.
121. Brunner, H., Arndt, M.R., and Treitinger, B., *Inorg. Chim. Acta*, 2004, vol. 357, p. 1649.
122. Wedel, M., *Dissertation*, Bremen, 2000.
123. Wedel, M., Walter, A., and Montforts, F.-P., *Eur. J. Org. Chem.*, 2001, p. 1681.
124. Kim, Y.-S., Song, R., Kim, D.H., Jun, M.J., and Sohn, Y.S., *Bioorg. Med. Chem.*, 2003, vol. 11, p. 1753.
125. Jin, G., Seo, E.-Y., Kang, S.-K., Ahn, B.-Z., and Shim, Y.K., *Korean J. Med. Chem.*, 1999, vol. 9, p. 2.
126. Zakharkin, L.I., Ol'shevskaya, V.A., Panfilova, S.Yu., Petrovskii, P.V., Luzgina, V.N., and Evstigneeva, R.P., *Russian Chem. Bull.*, 1999, vol. 48, p. 2312.
127. Evstigneeva, R.P., Luzgina, V.N., Timashov, P.S., Ol'shevskaya, V.A., and Zakharkin, L.I., *Russ. J. Gen. Chem.*, 2003, vol. 73, p. 1648.
128. Ol'shevskaya, V.A., Nikitina, R.G., Zaitsev, A.V., Gyl'malieva, M.A., Luzgina, V.N., Kononova, E.G., Morozova, T.G., Drozhzhina, V.V., Kaplan, M.A., Kalinin, V.N., and Shtil', A.A., *Dokl. Ross. Akad. Nauk*, 2004, vol. 399, p. 783.
129. Evstigneeva, R.P., Pashchenko, V.Z., Luzgina, V.N., Larkina, E.A., Gribkov, A.A., Tusov, V.B., and Gorokhov, V.V., *Dokl. Ross. Akad. Nauk*, 1999, vol. 369, p. 57.
130. Larkina, E.A., Luzgina, V.N., and Evstigneeva, R.P., *Bioorg. Khim.*, 2002, vol. 28, p. 357.
131. Larkina, E.A., Balashova, T.A., Luzgina, V.N., Konovalova, N.V., and Evstigneeva, R.P., *Mendeleev Commun.*, 2005, p. 234.
132. Murakami, H., Matsumoto, R., Okusa, Y., Sagara, T., Fujitsuka, M., Ito, O., and Nakashima, N., *J. Mater. Chem.*, 2002, vol. 12, p. 2026.
133. Cosnier, S., Walter, A., and Montforts, F.-P., *J. Porphyrins Phthalocyanines*, 1998, vol. 2, p. 39.
134. Hamachi, I., Tanaka, S., Tsukiji, S., Shinkai, S., and Oishi, S., *Inorg. Chem.*, 1998, vol. 37, p. 4380.
135. Evstigneeva, R.P., Zheltukhina, G.A., Khalil', V., and Efimova, E.I., *Bioorg. Khim.*, 1999, vol. 25, p. 572.
136. Hayashi, T., Ando, T., Matsuda, T., Yonemura, H., Yamada, S., and Hisaeda, Y., *J. Inorg. Biochem.*, 2000, vol. 82, p. 133.
137. Sondhi, S.M., Rajvanshi, S., Johar, M., and Dastidar, S.G., *Indian J. Chem., Sect. B*, 2002, vol. 41, p. 388.
138. Liang, Z.-X., Nocek, J.M., Huang, K., Hayes, R.T., Kurnikov, I.V., Beratan, D.N., and Hoffman, B.M., *J. Am. Chem. Soc.*, 2002, vol. 124, p. 6849.

139. Lombardi, A., Nastri, F., Marasco, D., Maglio, O., De Sanctis, G., Sinibaldi, F., Santucci, R., Coletta, M., and Pavone, V., *Chem. Eur. J.*, 2003, vol. 9, p. 5643.
140. Sol, V., Branland, P., Chaleix, V., Granet, R., Guilloton, M., Lamarche, F., Verneuil, B., and Krausz, P., *Bioorg. Med. Chem. Lett.*, 2004, vol. 14, p. 4207.
141. Nebol'sin, V.E., Zheltukhina, G.A., and Lobanova, T.N., Russian Patent no. 2250906, 2005; *Chem. Abstr.*, 2005, vol. 142, no. 411 147.
142. Sato, H., Watanabe, M., Hisaeda, Y., and Hayashi, T., *J. Am. Chem. Soc.*, 2005, vol. 127, p. 56.
143. Robinson, B., Morgan, A.R., and Narciso, H.L., Int. Patent no. 9705127, 1997; *Chem. Abstr.*, 1997, vol. 126, no. 211973.
144. Platzek, J., Niedballa, U., Raduechel, B., Weinmann, H.-J., Frenzel, T., Misselwitz, B., and Ebert, W., Int. Patent no. 0005235, 2000; *Chem. Abstr.*, 2000, vol. 132, no. 145791.
145. Sondhi, S.M., Singhal, N., Verma, R.P., Arora, S.K., and Dastidar, S.G., *Indian J. Chem., Sect. B*, 2001, vol. 40, p. 113.
146. Evstigneeva, R.P., Zarubina, T.V., Zheltukhina, G.A., Nebol'sin, V.E., Kaliberda, E.N., and Rumsh, L.D., *Dokl. Ross. Akad. Nauk*, 2001, vol. 381, p. 630.
147. Sahoo, S.K., Sawa, T., Fang, J., Tanaka, S., Miyamoto, Y., Akaike, T., and Maeda, H., *Bioconjugate Chem.*, 2002, vol. 13, p. 1031.
148. Sondhi, S.M., Johar, M., Singh, N., and Dastidar, S.G., *Indian J. Chem., Sect. B*, 2004, vol. 43, p. 162.
149. Kim, Y.-S., Song, R., Lee, C.O., and Sohn, Y.S., *Bioorg. Med. Chem. Lett.*, 2004, vol. 14, p. 2889.
150. Tanabe, S., Yamaguchi, M., Iijima, M., Nakajima, S., Sakata, I., Miyaki, S., Takemura, T., Furuoka, H., Kobayashi, Y., Matsui, T., Uzuka, Y., and Sarashina, T., *Veterinary J.*, 2004, vol. 167, p. 286.
151. Rogovina, S.Z., Solov'eva, A.B., Aksenova, N.A., and Zharov, A.A., *Vysokomol. Soedin. A, B.*, 2004, vol. 46, p. 421.
152. Mitina, V.Kh., Nechaeva, I.S., Ponomarev, G.V., Reshetnikov, A.V., and Klyashchitskii, B.A., *Bioorg. Khim.*, 1995, vol. 21, p. 301.
153. Mironov, A.F., Rumyantseva, V.D., and Ponamareva, O.N., *Mendeleev Commun.*, 1998, p. 187.
154. Torpey, J.W. and Ortiz de Montellano, P.R., *J. Org. Chem.*, 1995, vol. 60, p. 2195.
155. Torpey, J., Lee, D.A., Smith, K.M., and Ortiz de Montellano, P.R., *J. Am. Chem. Soc.*, 1996, vol. 118, p. 9172.
156. Torpey, J. and Ortiz de Montellano, P.R., *J. Biol. Chem.*, 1996, vol. 271, p. 26067.
157. Torpey, J. and Ortiz de Montellano, P.R., *J. Biol. Chem.*, 1997, vol. 272, p. 22008.
158. Vázquez, J., González, M.M., Martí, C., Nonell, S., and Trull, F.R., *Monatsh. Chem.*, 1998, vol. 129, p. 69.
159. Kalish, H.R., Latos-Grażyński, L., and Balch, A.L., *J. Am. Chem. Soc.*, 2000, vol. 122, no. 12478.
160. Robert, A., Cazelles, J., and Meunier, B., *Angew. Chem., Int. Ed.*, 2001, vol. 40, p. 1954.
161. Niemevz, F. and Buldain, G.Y., *J. Porphyrins Phthalocyanines*, 2004, vol. 8, p. 989.
162. Gerlach, B. and Montforts, F.-P., *Justus Liebigs Ann. Chem.*, 1995, p. 1509.
163. Chen, Y., Medforth, C.J., Smith, K.M., Alderfer, J., Dougherty, T.J., and Pandey, R.K., *J. Org. Chem.*, 2001, vol. 66, p. 3930.
164. Immoos, C.E., Bhaskar, B., Cohen, M.S., Barrows, T.P., Farmer, P.J., and Poulos, T.L., *J. Inorg. Biochem.*, 2002, vol. 91, p. 635.
165. Pont, F., Jacobs, N.J., and Montforts, J.-P., *Tetrahedron Lett.*, 1997, vol. 38, p. 6383.
166. Jacobs, M.J., Jacobs, N.J., Kuhn, C.B., Gorman, N., Dayan, F.E., Duke, S.O., Sinclair, J.F., and Sinclair, P.R., *Biochem. Biophys. Res. Commun.*, 1996, vol. 227, p. 195.
167. Jacobs, N.J., Kruszyna, H.G., Hier, J.S.L., Dayan, F.E., Duke, S.O., Pont, F., and Montforts, F.-P., *Biochem. Biophys. Res. Commun.*, 1999, vol. 259, p. 195.
168. Dayan, F.E., Duke, S.O., Faibis, V., Jacobs, J.M., and Jacobs, N.J., *Arch. Biochem. Biophys.*, 1998, vol. 351, p. 27.
169. Fischer, H. and Neumann, W., *Justus Liebigs Ann. Chem.*, 1932, vol. 494, p. 225.
170. Treibs, A., *Justus Liebigs Ann. Chem.*, 1969, vol. 728, p. 225.
171. Tyulyaeva, E.Yu., Lomova, T.N., and Andrianova, L.G., *Zh. Neorg. Khim.*, 2001, vol. 46, p. 432.
172. Lomova, S.T. and Andrianova, L.G., *Russ. J. Gen. Chem.*, 2003, vol. 73, p. 283.
173. Ishida, Y., Konishi, K., Nagamune, T., and Aida, T., *J. Am. Chem. Soc.*, 1999, vol. 121, p. 7947.
174. Hoeper, F. and Montforts, F.-P., *Justus Liebigs Ann. Chem.*, 1995, p. 1033.
175. Bats, J.W., Haake, F., Meier, M., Montforts, F.-P., and Scheurich, G., *Justus Liebigs Ann. Chem.*, 1995, p. 1617.