

SHORT
COMMUNICATIONSThermal Decomposition of 1-Dinitromethyl-3-nitro-1,2,4-triazole
Derivatives in Solution


R. S. Stepanov, L. A. Kruglyakova, and A. M. Astakhov

Siberian State University of Technology, pr. Mira 82, Krasnoyarsk, 660049 Russia

Received November 26, 2006

DOI: 10.1134/S1070428007030293

In continuation of our previous studies [1, 2] on the relations between the structure and the rate of thermal decomposition of polynitro compounds, we examined a series of 1-dinitromethyl-3-nitro-1H-1,2,4-triazole derivatives **I–V**.



I, R = Me; **II**, R = NO₂; **III**, R = Cl; **IV**, R = Br; **V**, R = I.

Compounds **I–V** were synthesized and purified by known methods [3]; they contained no less than 99% of the main substance (according to chromatographic data). The kinetics of their thermal decomposition were studied by the manometric technique [2] in dibutyl phthalate. The rate constants of their thermal decomposition do not depend on the concentration in solution in the range 1–5 wt %. The reactions follow the first-order kinetics up to a conversion of 40–50%. The activation parameters of the thermal decomposition of compounds **I–V** in dibutyl phthalate (2 wt %) are given below.

Comp. no.	I	II	III	IV	V
ΔT , °C	140–180	100–140	110–150	110–150	100–140
$k_{100^\circ\text{C}} \times 10^5$, s ⁻¹	0.02	1.1	3.9	9.4	18.0
E_a , kJ/mol	156	143	133	133	131
log A	15.1	15.1	14.2	14.7	14.6
E_s^a , kJ/mol	0	-2.14	-2.81	-3.44	-3.62

^a E_s is the substituent steric constant [4].

It follows from these data that the rate constant of thermal decomposition increases by three order of

magnitude in going from compound **I** to **V** and that the energy of activation decreases by more than 20 kJ/mol. An analogous pattern was observed in the decomposition of 5-dinitromethyl-2-methyltetrazole [2], for which homolytic mechanism with initial cleavage of the C–NO₂ bond was proved. Presumably, the thermal decomposition of compounds **I–V** follows a similar mechanism. This assumption is confirmed by the existence of correlations (1) and (2).

$$\log k_{100^\circ\text{C}} = -(6.69 \pm 0.05) - (0.80 \pm 0.02)E_s; \quad (1)$$

$$r = 0.999, s_y = 0.051, n = 5.$$

$$E_a = (156.05 \pm 2.18) + (6.98 \pm 0.80)E_s; \quad (2)$$

$$r = 0.975, s_y = 1.05, n = 5.$$

The negative value of δ (–0.80) in equation (1) indicates that steric effect of substituents favors the process. Low values of the energy of activation suggest intermediacy of a semirigid activated complex [5].

REFERENCES

- Stepanov, R.S., Kruglyakova, L.A., and Astakhov, A.M., *Russ. J. Org. Chem.*, 2001, vol. 37, p. 1793.
- Stepanov, R.S., Astakhov, A.M., Kruglyakova, L.A., and Golubtsova, O.A., *Russ. J. Gen. Chem.*, 2000, vol. 70, p. 934.
- Kofman, T.P., Kartseva, G.Yu., Glazkova, E.Yu., and Krasnov, K.N., *Russ. J. Org. Chem.*, 2005, vol. 41, p. 753.
- Shan'ko, V.N., Stepanov, R.S., and Gidasov, B.V., *Materialy konferentsii po itogam NIR. Sektsiya organicheskoi khimii* (Proc. Conf. on the Results of Research Work), Krasnoyarsk: Sib. Tekhnol. Inst., 1971, p. 41.
- Nazin, G.M. and Manelis, G.B., *Usp. Khim.*, 1994, vol. 63, p. 327.