

SHORT COMMUNICATIONS

Synthesis of 1-Alkylamino-3*H*-naphtho-[1,2,3-*de*]quinoline-2,7-diones

M. S. Sokolova, T. I. Lavrikova, and L. M. Gornostaev

Astaf'ev Krasnoyarsk State Pedagogical University, ul. A. Lebedevoi 89, Krasnoyarsk, 660049 Russia


e-mail: gornostaev@kspu.ru

Received November 15, 2006

DOI: 10.1134/S1070428007040252

1-Alkylamino-3*H*-naphtho[1,2,3-*de*]quinoline-2,7-diones **I** exhibit luminescence properties [1]. Known methods for the synthesis of these compounds are based on nucleophilic replacement of hydrogen or chlorine atom or nitro or sulfo group in the corresponding 1-substituted 3*H*-naphtho[1,2,3-*de*]quinoline-2,7-diones **Ia–Id** (Scheme 1) [2–4].

Scheme 1.



R = Me, X = H (**a**), O₂N (**b**); R = H, X = SO₃Na (**c**), Cl (**d**);
R' = Alk, Ar.

However, initial compounds **Ia–Id** are difficultly accessible, and the yields of the target alkylamino derivatives are often poor. The alkylation of 1-amino-3*H*-naphtho[1,2,3-*de*]quinoline-2,7-dione (**I**, R = H, X = NH₂) does not give compounds **II** but involves the endocyclic nitrogen atom [5].


We have found that readily available 1-(*N*-nitroso-*N*-alkylaminoacetamido)-9,10-anthraquinones **IIIa** and **IIIb** undergo cyclization to alkylaminonaphthoquinolines **IIa** and **IIb** under mild conditions (Scheme 2). Presumably, the intramolecular ring closure is followed by elimination of the nitroso group, for analogous cyclization of the precursors of nitrosamines **IIIa** and **IIIb**, 1-(alkylaminoacetamido)-9,10-anthraquinones **IVa** and **IVb**, does not occur under similar conditions.

The reaction is also accompanied by decomposition of nitrosamines **IIIa** and **IIIb** with formation of 1-amino-9,10-anthraquinone which was detected by TLC in the reaction mixture.

It is known that electron-withdrawing substituents in the acetylamino group of acetylaminoanthraquinones facilitate their cyclization to the corresponding 3*H*-naphtho[1,2,3-*de*]quinoline-2,7-diones [6]. In the ¹H NMR spectra of both *E* and *Z* isomers of **IIIa** and **IIIb**, singlets from the methylene protons are displaced downfield by 1.2–2.0 ppm relative to the corresponding signals of 1-(alkylaminoacetamido)-9,10-anthraquinones **IVa** and **IVb**, so that they appear approximately in the same region as the methylene proton signal of 1-chloroacetyl-amino-9,10-anthraquinone (δ 4.55 ppm). The *N*-nitrosoalkylamino groups in **IIIa** and **IIIb** are stronger electron acceptors than chlorine atom, and this factor is likely to favor the cyclization of **III** into **II**.

1-Ethylamino-3*H*-naphtho[1,2,3-*de*]quinoline-2,7-dione (IIa**).** A mixture of 0.58 g (1.7 mmol) of compound **IIIa**, 8 ml of dimethyl sulfoxide, 1.5 g of anhydrous sodium sulfate, and 1 g of potassium carbonate was stirred for 1 h at 60°C. The mixture was cooled and poured onto 100 g of ice, the precipitate was filtered off and treated with 6 ml of benzene at 60–70°C, the mixture was cooled, and the precipitate was filtered off. This procedure was repeated once more. Yield 0.50 g (70%), mp 270–273°C. UV spectrum (DMF), λ_{max}, nm (log ε): 345 (3.80), 445 (4.20) (cf. [4]). ¹H NMR spectrum, δ, ppm: 1.06 t (3H, CH₃, *J* = 7 Hz), 3.05 m (2H, CH₂), 7.50–7.62 m and 7.98–8.07 m (7H, H_{arom}), 8.28 m (1H, NH), 12.40 s (1H, NH). Found, %: C 73.84; H 4.79; N 9.32. C₁₈H₁₄N₂O₂. Calculated, %: C 74.48; H 4.82; N 9.65.

Scheme 2.



R = Me (a), Ph (b).

1-Benzylamino-3H-naphtho[1,2,3-de]quinoline-2,7-dione (IIb) was synthesized in a similar way from 1.5 g (4.2 mmol) of compound **IIIb**, 15 ml of DMSO, 2 g of Na₂SO₄, and 2 g of K₂CO₃. Yield 1 g (75%), mp 280–283°C. UV spectrum (DMF), λ_{max} , nm (log ϵ): 345 (3.70), 445 (4.17). ¹H NMR spectrum, δ , ppm: 4.32 d (2H, CH₂, J = 7 Hz), 7.01–7.18 m (5H, C₆H₅), 7.90 t (1H, NH, J = 8 Hz), 7.50–7.51 m and 8.00–8.40 m (7H, H_{arom}), 12.35 s (1H, NH). Found, %: C 78.01; H 4.41; N 8.15. C₂₃H₁₆N₂O₂. Calculated, %: C 78.40; H 4.54; N 7.95.

N-(9,10-Dioxo-9,10-dihydroanthracen-1-yl)-2-[ethyl(nitroso)amino]acetamide (IIIa). Sodium nitrite, 0.8 g (11 mmol), was added under stirring at 20–22°C to a solution of 1.23 g (4 mmol) of compound **IVa** in 30 ml of acetic acid. After 30–40 min, the precipitate was filtered off and washed with water (10 ml), ethanol (10 ml), and diethyl ether (10 ml). Yield 1.20 g (89%), mp 134–136°C. According to the ¹H NMR data (taking into account that signals of *E* isomers appear in a weaker field [7]), compound **IIIa** was a mixture of *Z* and *E* isomers at a ratio of 3:1. ¹H NMR spectrum, δ , ppm: 1.06 t (0.76H, CH₃, J = 8 Hz); 1.45 t (2.25H, CH₃, J = 8 Hz); 3.70 q (0.5H, CH₂CH₃, *Z* isomer, J = 8 Hz); 4.40 q (1.5H, CH₂CH₃, *E* isomer, J = 8 Hz); 4.57 s (1.5H, COCH₂, *Z* isomer); 5.31 s (0.5H, COCH₂, *E* isomer); 7.85–8.20 m, 8.18–8.26 m, and 8.86–8.95 m (7H, H_{arom}); 12.09 s (0.75H, NH, *Z* isomer); 12.15 s (0.25H, NH, *E* isomer). Found, %: C 64.40; H 4.33; N 11.95. C₁₈H₁₅N₃O₄. Calculated, %: C 64.09; H 4.45; N 12.46.

2-[Benzyl(nitroso)amino]-N-(9,10-dioxo-9,10-dihydroanthracen-1-yl)acetamide (IIIb). Sodium nitrite, 1.2 g (17 mmol), was added under stirring at 20–22°C to a solution of 3.00 g (7.5 mmol) of compound **IVb** in 50 ml of acetic acid. The product was isolated as described above for **IIIa**. Yield 2.80 g (87%) (a mixture of *Z* and *E* isomers at a ratio of 1:2), mp 178–180°C. ¹H NMR spectrum, δ , ppm: 4.48 s (1.32H, CH₂Ph, *E* isomer); 4.91 s (0.66H, CH₂Ph, *Z* isomer); 5.30 s (0.66H, COCH₂, *Z* isomer); 5.59 s (1.32H, COCH₂, *E* isomer); 7.40–7.52 m, 7.89–7.97 m, 8.23–8.30 m, and 8.79–8.90 m (12H, H_{arom}); 12.01 s (0.75H, NH, *Z* isomer); 12.20 s (0.25H, NH, *E* isomer). Found, %: C 69.17; H 4.25; N 10.44. C₂₃H₁₇N₃O₄. Calculated, %: C 69.17; H 4.26; N 10.52.

N-(9,10-Dioxo-9,10-dihydroanthracen-1-yl)-2-(ethylamino)acetamide (IVa). A mixture of 2.00 g (6.6 mmol) of 2-chloro-*N*-(9,10-dioxo-9,10-dihydroanthracen-1-yl)acetamide, 30 ml of dioxane, and 5 ml of 70% aqueous ethylamine was heated for 2 h at 60°C. After cooling, the yellow precipitate was filtered off and purified by recrystallization from ethanol. Yield 1.63 g (79%), mp 138–140°C. ¹H NMR spectrum, δ , ppm: 1.19 t (3H, CH₃, J = 6 Hz); 2.66 q (2H, CH₂CH₃, J = 6 Hz); 3.25 br.s (1H, NHCH₂); 3.37 br.s (2H, CH₂NH); 7.83–7.92 m, 8.13–8.21 m, and 9.10–9.12 m (7H, H_{arom}); 12.10–13.60 br.s (1H, NHCO). Found, %: C 70.42; H 5.07; N 8.72. C₁₈H₁₆N₂O₃. Calculated, %: C 70.12; H 5.19; N 9.09.

The ¹H NMR spectra were recorded on a Bruker DRX-500 spectrometer from solutions in DMSO-*d*₆

using tetramethylsilane as internal reference. The UV spectra were measured on a Helios ϵ spectrophotometer. The progress of reactions and the purity of products were monitored by TLC on Silufol plates (toluene–acetone, 10:1). 2-Chloro-*N*-(9,10-dioxo-9,10-dihydroanthracen-1-yl)acetamide was synthesized by the procedure described in [8].

This study was performed under financial support by the V.P. Astaf'ev Krasnoyarsk State Pedagogical University (project nos. 42-06-1/FP, 22-06-1/FP).

REFERENCES

1. Krasovitskii, B.M. and Bolotin, B.M., *Organicheskie lyuminofory* (Organic Luminophores), Leningrad: Khimiya, 1976, p. 154.
2. Kazankov, M.V. and Ufimtsev, V.N., *Khim. Geterotsikl. Soedin.*, 1972, no. 3, p. 373.
3. Allen, C.F.H. and Wilson, C.V., *J. Org. Chem.*, 1945, vol. 10, p. 594.
4. Sadchenko, L.S. and Gudzenko, V.I., *Zh. Org. Khim.*, 1976, vol. 12, no. 5, p. 1106.
5. Elizbarashvili, E.N., Lagvilava, I.V., and Samsoniya, Sh.A., *Khim. Geterotsikl. Soedin.*, 2005, p. 1868.
6. *Methoden der organischen Chemie (Houben–Weyl)*, Müller, E., Bayer, O., Meerwein, H., and Ziegler, K., Eds., Stuttgart: Georg Thieme, 1979, vol. 7/3c, p. 351.
7. *Comprehensive Organic Chemistry*, Barton, D. and Ollis, W.D., Eds., Oxford: Pergamon, 1979, vol. 2. Translated under the title *Obshchaya organicheskaya khimiya*, Moscow: Khimiya, 1985, vol. 3, p. 453.
8. Kazankov, M.V., Putsa, M.V., and Mukhina, L.L., *Khim. Geterotsikl. Soedin.*, 1972, p. 1651.