

SHORT
COMMUNICATIONS

3,4-Dibromotetrahydro- λ^6 -thiophene 1,1-Dioxide in S,N-Tandem Heterocyclizations. Synthesis of Tetrahydrothieno-thiazolopyrimidines


A. A. Shklyarenko and V. V. Yakovlev

“Applied Chemistry” Russian Research Center, pr. Dobrolyubova 14, St. Petersburg, 197198 Russia
e-mail: synthon@inbox.ru

Received November 17, 2006

DOI: 10.1134/S1070428007040318

Pyrimidine derivatives possess important pharmacological properties [1]. Among these, the most prominent are such naturally occurring alkaloids as batzelladines [2], ptilomycalins [3], saxitoxins [4], and luotonins [5]; their molecules include five-membered saturated rings fused to a pyrimidine base. We previously showed [6] that 2,3-dibromopropyl sulfones readily undergo heterocyclization with 2-thiouracils. While continuing studies on tandem heterocyclizations, we have synthesized tetrahydrothienothiazolopyrimidine derivatives via S,N-tandem alkylation of 2-thiouracils **Ia** and **Ib** with 3,4-dibromotetrahydro- λ^6 -thiophene 1,1-dioxide. The reactions were carried out in ethanol at room temperature, the reactant ratio dibromosulfolane–2-thiouracil **I**–potassium hydroxide being 1:2:4; the reactions were complete in 8 h, and the yields of products **IIa** and **IIb** were 78–80%.



Products **IIa** and **IIb** were isolated as individual substances, and their structure was confirmed by the

¹H NMR and mass spectra. The ¹H NMR spectra of **IIa** and **IIb** contained signals from the CH₂CHCHCH₂ fragment as an ABMM'NN' spin system having four chiral centers. In the mass spectra of **IIa** and **IIb**, the most characteristic were peaks from the following fragment ions: [M – SO₂H]⁺, [M – SO₂]⁺, [M – SO₂ – SH]⁺, [M – SO₂ – SH – C₂H₂]⁺. The fragmentation pattern was proposed on the basis of published data [7, 8]. The absence of isomeric compounds among the products indicates that the reaction is chemo- and regioselective and that the process follows a concerted mechanism typical of tandem reactions [9].

2-Amino-5a,6,8,8a-tetrahydro-4H-7 λ^6 -thieno-[3',4':4,5][1,3]thiazolo[3,2-a]pyrimidin-4-one 7,7-dioxide (IIa). Yield 78%, mp 276–277°C (from EtOH–DMF). ¹H NMR spectrum, δ , ppm: 3.38 m (2H, CH), 3.65 m (2H, CH), 4.56 m (1H, CH), 4.85 s (1H, H_{arom}), 5.33 m (1H, CH), 6.38 br.s (2H, NH₂). Mass spectrum, m/z (I_{rel} , %): 259 (100) [M]⁺, 194 (30.5), 195 (51.1), 162 (62.2), 136 (58.8).

2-Methoxy-5a,6,8,8a-tetrahydro-4H-7 λ^6 -thieno-[3',4':4,5][1,3]thiazolo[3,2-a]pyrimidin-4-one 7,7-dioxide (IIb). Yield 80%, mp 263–264°C (from EtOH–DMF). ¹H NMR spectrum, δ , ppm: 3.56 m (2H, CH), 3.68 m (2H, CH), 3.77 s (3H, CH₃), 4.79 m (1H, CH), 5.33 s (1H, H_{arom}), 5.43 m (1H, CH). Mass spectrum, m/z (I_{rel} , %): 274 (100) [M]⁺, 209 (36.6), 210 (52.4), 177 (64.9), 151 (48.6).

The ¹H NMR spectra were recorded from solutions in DMSO-*d*₆ on a Bruker AM-500 spectrometer (500.13 MHz) using the residual proton signal of the solvent as reference. The mass spectra (electron

impact, 70 eV) were obtained on an MKh-1321 mass spectrometer with direct sample admission into the ion source. LC–MS analysis was performed on a Micro-mass ZDM-2000 instrument (electrospray ionization, positive ion registration).

REFERENCES

1. Rovnyak, G.C., Kimball, S.D., Beyer, B., Cucinotta, G., DiMarco, J.D., Gougoutas, J., Hedberg, A., Malley, M., McCarthy, J., Zhang, R., and Moreland, S., *J. Med. Chem.*, 1995, vol. 38, p. 119.
2. Snider, B.B., Chen, J., Patil, A.D., and Freyer, A., *Tetrahedron Lett.*, 1996, vol. 37, p. 6977.
3. Heys, L., Moore, C.G., and Murphy, P.J., *Chem. Soc. Rev.*, 2000, vol. 29, p. 57.
4. Kishi, Y., *Heterocycles*, 1980, vol. 14, p. 1477.
5. Santosh, B.M. and Narshinha, P.A., *Synthesis*, 2002, p. 323.
6. Shklyarenko, A.A., Chernitsa, B.V., and Yakovlev, V.V., *Russ. J. Org. Chem.*, 2005, vol. 41, p. 1097.
7. Ponomarev, D.A. and Takhistov, V.V., *Recent Advances in Analytical Techniques*, Attaur-Rahman, M., Ed., Reading: Hardwood Academic, 2002, p. 369.
8. Ponomarev, D.A., Golovin, A.V., and Takhistov, V.V., *Eur. J. Mass Spectrom.*, 2002, vol. 8, p. 409.
9. Nicolaou, K.C., Tamsyn, M., and Snyder, S.A., *Chem. Commun.*, 2003, p. 551.