

Synthetic Approach to Preparation of Indole Derivatives Fused with a Bicyclo[3.3.1]nonane Framework

T. Yu. Baranova, O. N. Zefirova, N. V. Averina, V. V. Boyarskikh,
G. S. Borisova, N. V. Zyk, and N. S. Zefirov

Moscow State University, Moscow, 119992 Russia

Received July 21, 2006

Abstract—Synthetic procedures for *N*-(2-methoxy-6,7,8,9,10,11-hexahydro-5*H*-6,10-methanocycloocta[*b*]-indol-9-yl)acetamide were investigated. The desired product was obtained from the mono ethylene ketal of bicyclo[3.3.1]nonane-2,6-dione in a six-stage procedure involving as the key point the formation of an indole fragment on the bicyclo[3.3.1]nonane framework by Fischer reaction.


DOI: 10.1134/S1070428007080192

Melatonin [*N*-acetyl-5-methoxytryptamine (**I**)] is an endogeneous hormone endowed with chronobiotic, neuroprotection, immunostimulating, and anticancer activity; it is bonded in the body with three subtypes of protein receptor molecules. Recently intensive research is carried out worldwide directed on the preparation of synthetic substances possessing a high affinity and selectivity with respect to each of the melatonin receptors subtypes. In designing these compound a most common method is hampering of the conformational mobility of the melatonin side chain.

The target of the present study was a development of approaches to the synthesis of a “conformationally limited” melatonin analog with an indole fragment fused to a bicyclo[3.3.1]nonane framework: *N*-(2-methoxy-6,7,8,9,10,11-hexahydro-5*H*-6,10-methanocycloocta[*b*]-indol-9-yl)acetamide (**II**).


The published procedures for preparation of similar fused systems involve mainly building up of the polycyclic fragment on the indole structure (see, e.g., [1–4]). However an inverse method was described in [5–

7]: indole fragment was built on the bicyclo- and azabicyclo[3.3.1]nonane framework by Fischer reaction; just this approach we chose for the synthesis of compound **II**.




We studied first different methods of acetamide substituent formation in the position 6 of the bicyclo[3.3.1]nonane skeleton fused with an unsubstituted indole (Scheme 1).

Initial ethylene ketal **III** was obtained by procedure [8] using toluene as solvent that resulted in some



improvement of the yield of the desired product. The attempt to carry out Fischer reaction with **III** as in [5] with phenylhydrazine in acetic acid at 80°C gave final product **IVa** with an impurity of the corresponding hydrazone. A more convenient preparation process was the reaction carried out in methanol in the presence of zinc chloride as a catalyst; this method provided indole derivative **IVa** in 53% yield.

We tried two procedures for preparation of amine **Va**: a) a reductive amination of ketone **IVb**, and b) through oxime **IVc** with its subsequent reduction into amine. The reaction of ketone **IVb** with sodium borohydride in water-alcoholic ammonia solution resulted however in a product with an IR spectrum lacking the carbonyl group absorption band and with a broad absorption band at 3200–3400 cm⁻¹. The mass spectrum of the compound contained a peak with M^+ 227 and also a peak 209 [$M - H_2O$]⁺, and the ¹H NMR spectrum, a broad singlet at δ 2.02 (OH) and a multiplet at δ 3.84 ppm. (HCOH). The signal of C–OH in the ¹³C NMR spectrum appeared at δ 73 ppm. These spectral characteristics and the elemental analysis data indicate that in the reaction in question forms alcohol **Vc** previously described in [6].



Next we tested the possibility to synthesize amine **Va** via oxime **IVc**. We also tried to prepare the latter by two alternative ways: by Fischer reaction of bicyclo[3.3.1]nonane-2,6-dione monooxime (**VI**), and from ketone **IVb**.

Compound **VI** was prepared by a standard procedure from ketal **III** and hydroxylamine hydrochloride in a water-alcohol medium in the presence of NaOH leading to the formation of mono ethylene ketal oxime of bicyclo[3.3.1]nonane-2,6-dione* with the subsequent removal of the protective group. Fischer reaction with monooxime **VI** neither in the presence of acetic acid nor catalyzed by zinc chloride provided the desired oxime **IVc**. In the first case mainly the bicyclo[3.3.1]nonane-2,6-dione phenylhydrazone was obtained (mp 141–142°C [5] and m/z 332) and also the oxime of bicyclo[3.3.1]nonane-2,6-dione hydrazone [mass spectrum: 257 [M^+], 240 [$M - OH$]⁺; IR spectrum: 3330 (NH of hydrazone), 1605 cm⁻¹ (C=N)].

* Yield of mono ethylene ketal oxime of bicyclo[3.3.1]nonane-2,6-dione was 67%. IR spectrum, cm⁻¹: 3100–3300 (OH), 1660 (C=N). Found, %: C 62.15; H 7.86; N 6.24. C₁₁H₁₇NO₃. Calculated, %: C 62.54; H 8.11; N 6.63.

Oxime **IVc** was obtained in a quantitative yield from ketone **IVb** in a water-alcohol medium with hydroxylamine hydrochloride preliminary treated with K₂CO₃. The IR spectrum of oxime **IVc** contained a characteristic absorption band at 3400 cm⁻¹ (NH of indole) on the background of the OH group absorption (3200–3500 cm⁻¹). In the ¹H NMR spectrum of oxime **IVc** a signal was observed at δ 4.12 ppm corresponding to the NOH proton, and in the ¹³C NMR spectrum a signal at δ 163.77 ppm corresponded to C=N.


The oxime was reduced to the corresponding amine with lithium aluminum hydride in anhydrous THF. The oily amine **Va** obtained was further acetylated. The structure of compound **Va** was confirmed by mass spectra and by ¹H and ¹³C NMR spectra.

The acetylation of amine **Va** was carried out by treating with acetic anhydride in the presence of pyridine. The IR spectrum of compound **Vb** purified by chromatography contained the absorption band of the NH group from the indole fragment (3400 cm⁻¹) and absorption bands of amide groups C=O (1650) and NH (3330 cm⁻¹). The formation of amide **Vb** was also confirmed by ¹H NMR spectrum where appeared a broad singlet of the amide group NH proton at δ 5.67 ppm and a broad singlet at δ 4.07 ppm of $W_{1/2}$ 28.8 Hz corresponding to the H⁶ proton of the *endo*-isomer of compound **Vb** (the assignment is based on [9]). A single isomer of compound **Vb** was yielded by chromatographic purification.

In the next stage we developed the way of indole fragment functionalization aiming at the preparation of the target compound **II**. We attempted to introduce a methoxy group into the position 5' by the classic method of 5-substituted indoles synthesis via bromination of unsubstituted ethylene ketal **IVa** followed by replacement of the bromine by a methoxy group [10, 11] (Scheme 2, a).

After the bromination of compound **IVa** with bromine vapor in sulfuric acid in the presence of silver sulfate [10, 11] we isolated two main products from the reaction mixture, the first one being oxindole **IVb**. The second product gave a positive response to Beilstein's test, and its mass spectrum contained ion peaks of m/z 305 and 303 (30 and 27% respectively) thus corresponding to the empirical formula C₁₅H₁₄BrNO, and also peaks of ions formed on elimination of isotopes ⁷⁹Br and ⁸¹Br from M^+ (226, 225, 224 of intensity 35.83 and 75% respectively). According to IR and ¹H NMR spectra the indole fragment did not suffer any changes in the reaction process, and the substitution occurred in the α -position with respect to the carbonyl. Obviously the bromination

Scheme 2.



did not proceed at the tertiary position in the bridgehead; moreover, in the ¹H NMR spectrum a signal was observed of the H–C–Br proton at 4.5 ppm. Besides, in the region 2.7–3.45 ppm signals appeared belonging to protons H¹, H^{4a}, H^{4ε}, and H⁵ of the bicyclo[3.3.1]nonane framework (assignment based on [9]) that are close in position to those observed in the spectra of compounds **IVa**, **IVb** and **Va**, **Vb**. All the data obtained confirm, that the bromination of ethylene ketal **IVa** occurred not in the aromatic ring but at a methylene group adjacent to a carbonyl (structure **VII**, Scheme 2, a).

The methoxy group was more successfully introduced into the indole fragment by Fischer reaction of compound **III** with 1-(4-methoxyphenyl)hydrazine hydrochloride in the presence of ZnCl₂. Compound **VIIIa** was obtained as a result in 68% yield (Scheme 2, b). Its further treatment with 2 N H₂SO₄ at 50–60°C led to the formation of the corresponding ketone **VIIIb**. The signals of protons of OCH₃ groups were observed in the ¹H NMR spectra of both compounds **VIIIa** and **VIIIb** in the region 3.87 ppm.

Acetamino group formation in compound **VIIIb** in order to obtain desired structure **II** was carried out similarly to the above procedure via oxime **VIIIc** prepared quantitatively from compound **VIIIb**. The reduction of compound **VIIIc** with lithium aluminum hydride gave amine **IX** that was without isolation brought into reaction with acetic anhydride in the presence of pyridine. We synthesized as a result the target product **II** whose NMR spectra show the formation of two (*endo*- and *exo*-) isomers of compound **II** with respect to acetamino group in a ratio 6:1 as confirmed also by chromato-mass spectral data. In the ¹H and ¹³C NMR

spectra of compound **II** the signals of all protons and all carbon atoms of the bicyclic fragment of both isomers of compound **II** are clearly seen (see EXPERIMENTAL). The value *W*_{1/2} 29 Hz suggested [9] the prevalence of the *endo*-isomer **II** that we isolated in the individual state by chromatography.

Hence we developed in this study a preparative procedure for the synthesis of *N*-(2-methoxy-6,7,8,9,10,11-hexahydro-5*H*-6,10-methanocycloocta-*[b]*indol-9-yl)acetamide (**II**), potential analog of melatonin “with a limited conformational mobility”.

EXPERIMENTAL

¹H and ¹³C NMR spectra were registered on a spectrometer Bruker AMX-400 at operating frequencies 400 and 100 MHz respectively using HMDS as internal reference. IR spectra were recorded on spectrophotometers UR-20 and Specord 75 IR from mulls in mineral oil. Mass spectra were measured on a GC-MS instrument JMS-D300. The reaction progress was monitored by TLC on Silufol UV-254 plates. Separation by column chromatography was carried out on silica gel Lancaster (60–200 μm).

N-(2-Methoxy-6,7,8,9,10,11-hexahydro-5*H*-6,10-methanocycloocta-*[b]*-indol-9-yl)acetamide (**II**) was obtained in the same way as amide **Vb**, yield 30% calculated on oxime **VIIIc**. The product was purified by chromatography (eluent methanol–chloroform, 1:20), mp 140°C. IR spectrum, cm^{−1}: 1660 (C=O), 3300 (NH of amide), 3390 (NH of indole).

Endo-isomer. ¹H NMR spectrum (CDCl₃), δ, ppm: 7.62 br.s (1H, NH of indole), 7.20 d (1H, H⁷, *J* 8.60 Hz),

6.96 d (1H, $H^{4'}$, J 2.16 Hz), 6.80 d.d (1H, $H^{6'}$, J 2.54, 8.60 Hz), 5.51 br.s (1H, NH of amide), 4.08 m (1H, H^6 , $W_{1/2}$, 29.0 Hz), 3.89 s (3H, OCH_3), 3.00 br.s (1H), 2.83 d.d (1H, $HC^{4\epsilon}$, J 6.66, 16.63 Hz), 2.64 d (1H, H^{4a} , J 17.02 Hz), 2.60 br.s (1H), 2.10 m (1H), 2.03 s [3H, $NC(O)CH_3$], 2.00–1.07 m (5H). ^{13}C NMR spectrum ($CDCl_3$), δ , ppm: 169.41 (C=O), 153.94, 1137.52, 130.66, 127.47, 111.35, 110.67, 109.59, 100.10, 55.93 (C^6), 32.08, 31.13, 30.73, 28.42, 25.16, 20.88. Mass spectrum, m/z (I_{rel} , %): 298 (100) [M] $^+$, 281 (3), 254 (24), 238 (46), 224 (16), 210 (47), 198 (73), 183 (18), 167 (34).

Exo-isomer. 1H NMR spectrum ($CDCl_3$), δ , ppm: 7.62 br.s (1H, NH of indole), 7.19 d (1H, $H^{7'}$, J 8.84 Hz), 6.93 d (1H, $H^{4'}$, J 2.53 Hz), 6.80 d.d (1H, $H^{6'}$, J 2.54, 8.60 Hz), 5.90 br.s (1H, NH of amide), 4.04 m (1H, H^6 , $W_{1/2}$ 23.5 Hz), 3.89 s (3H, OCH_3), 3.05 d.d (1H, $CH^{4\epsilon}$, J 7.32, 16.68 Hz), 3.04 br.s (1H), 2.65 d (1H, HC^{4a} , J 16.93 Hz), 2.41 br.s (1H), 2.10 m (1H), 2.06 s [3H, $NC(O)CH_3$], 2.00–1.24 m (5H). ^{13}C NMR spectrum ($CDCl_3$), δ , ppm: 196.44 (C=O), 136.54, 135.63, 127.13, 121.06, 119.18, 117.65, 110.76, 109.69, 51.84 (C–NH), 32.13, 31.16, 30.70, 28.32, 25.11, 23.64, 20.85. Mass spectrum, m/z (I_{rel} , %): 298 (33) [M] $^+$, 280 (1), 253 (1), 239 (100), 224 (6), 210 (13), 198 (54), 182 (10), 167 (14).

5',6',7',8',10',11'-Hexahydrospiro(1,3-dioxolane-2,9')-[6,10]methanocycloocta[b]indole) (IVa). To a solution of 0.3 g (1.53 mmol) of mono ethylene ketal **III** in 3 ml of methanol was added at stirring 0.165 g (1.53 mmol) of phenylhydrazine and 0.75 g (5.50 mmol) of anhydrous $ZnCl_2$. The mixture was boiled for 6 h, washed with water and methanol. Yield 0.22 g (53%), mp 184–186°C (EtOH) (185–187°C [5]). 1H NMR spectrum ($CDCl_3$), δ , ppm: 7.66 br.s (1H, NH), 7.48 d (1H, $H^{7'}$, J 7.58 Hz), 7.30 d (1H, $H^{4'}$, J 7.33 Hz), 7.16–7.07 m (2H, $H^{5',6'}$), 4.00 m [4H, (OCH_2) $_2$], 3.05–1.45 m (10H). ^{13}C NMR spectrum ($CDCl_3$), δ , ppm: 136.00, 135.52, 127.54, 120.84, 119.06, 117.90, 111.30, 110.44, 109.06, 64.30 (OCH_2CH_2O), 38.52, 30.38, 29.87, 28.52, 28.32, 22.24. Mass spectrum, m/z (I_{rel} , %): 269 (100) [M] $^+$, 236 (5), 254 (71), 210 (50), 200 (40).

5',6',7',8',10',11'-Hexahydro-9H-6,10-methanocycloocta[b]indole-9-one (IVb) was obtained as described in [5], yield 88%, mp 145–147°C (EtOH) (151–153°C [5]). 1H NMR spectrum ($CDCl_3$), δ , ppm: 7.49 br.s (1H, NH), 7.47 d (1H, $H^{7'}$, J 7.82 Hz), 7.33 d (1H, $H^{4'}$, J 7.82 Hz), 7.18 t, 7.13 t (2H, $H^{5',6'}$, J 7.05 Hz), 3.24 br.s (1H), 3.13 d.d (1H, $H^{4\epsilon}$, J 6.65, 16.44 Hz), 3.03 br.s (1H), 2.83 d (1H, H^{4a} , J 16.63 Hz), 2.37–2.10 m (6H).

Mass spectrum, m/z (I_{rel} , %): 225 (65) [M] $^+$, 197 (37), 168 (100), 167(175), 156 (12).

5',6',7',8',10',11'-Hexahydro-9H-6,10-methanocycloocta[b]indole-9-one oxime (IVc). To a solution of 0.14 g (2.00 mmol) of hydroxylamine hydrochloride in 5 ml of ethanol was added at stirring 0.14 g (1.00 mmol) of K_2CO_3 and then dropwise, a solution of 0.40 g (1.78 mmol) of compound **IVb** in 10 ml of ethanol. The reaction mixture was stirred for 3 h, then 20 ml of water was added, extracted with CH_2Cl_2 , the organic layer was dried over Na_2SO_4 , and the solution was evaporated on a rotary evaporator. The crude product was subjected to chromatography (eluent benzene–ethyl acetate, 3:1) to isolate 0.30 g (70%) of oxime **IVc** as colorless crystals, mp 125°C. IR spectrum, cm^{-1} : 3400 (NH of indole) on the background 3200–3500 (OH). 1H NMR spectrum ($CDCl_3$), δ , ppm: 7.78 br.s (1H, NH), 7.48 d (1H, $H^{7'}$, J 8.02 Hz), 7.34 d (1H, $H^{4'}$, J 7.83 Hz), 7.18, 7.12 t (2H, $H^{5',6'}$, J 7.24 Hz), 4.12 br.s (1H, NOH), 3.25–1.60 m (10H). ^{13}C NMR spectrum ($CDCl_3$), δ , ppm: 163.77 (C=N), 135.67, 134.86, 127.25, 121.41, 119.41, 117.97, 110.66, 109.21 (C^3), 34.64, 32.53, 30.25, 29.08, 27.44, 26.25. Found, %: C 75.03; H 6.55; N 11.37. $C_{15}H_{16}N_2O$. Calculated, %: C 75.00; H 6.67; N 11.67.

6',7',8',9',10',11'-Hexahydro-5H-6,10-methanocycloocta[b]indol-9-amine (Va). To a dispersion of 0.28 g (7.37 mmol) of $LiAlH_4$ in 10 ml of anhydrous THF was added dropwise at stirring a solution of 0.6 g (2.5 mmol) of oxime **IVc** in 10 ml of anhydrous THF, the reaction mixture was boiled for 6.5 h. On cooling with ice 1.5 ml of water–THF mixture, 1:1, was added, the mixture was boiled for 1 h, the precipitate was filtered off and washed with hot THF. The combined filtrates were dried over Na_2SO_4 , and the solution was evaporated on a rotary evaporator to obtain 0.56 g (99%) of yellow oily substance. On treating with ethyl acetate solid amine **Va** crystallized (yield ~30%). IR spectrum, cm^{-1} : 3390 br (NH) on a background of wide absorption in the region 3100–3450 (NH_2). 1H NMR spectrum ($DMSO-d_6$), δ , ppm: 10.52 br.s (1H, NH), 7.35 d (1H, $H^{7'}$, J 7.33 Hz), 7.23 d (1H, $H^{4'}$, J 8.08 Hz), 6.97, 6.91 t (2H, $H^{5',6'}$, J 7.58, 7.08 Hz), 2.96 m (2H, H^{4a}), 2.84 m (1H, H^6), 2.55 d.d (1H, $H^{4\epsilon}$, J 6.83, 16.43 Hz), 2.08–0.80 m (10H, framework, NH_2). ^{13}C NMR spectrum ($DMSO-d_6$), δ , ppm: 137.72, 135.95, 127.39, 120.16, 118.26, 117.67, 111.13, 109.37, 53.99 (C^6), 34.79, 32.92, 30.95, 28.63, 28.36, 19.37. Mass spectrum, m/z (I_{rel} , %): 226 (100) [M] $^+$, 209 (70), 194 (20), 180 (75), 168 (95).

N-(6',7',8',9',10',11'-Hexahydro-5H-6,10-methanocycloocta[b]indol-9-yl)acetamide (Vb).

A mixture of 0.11 g of oily amine **Va**, 2 ml of pyridine, and 1 ml of Ac₂O was stirred for 12 h at room temperature, then it was evaporated, and Et₂O was added. The ether solution was washed with 1 N HCl and 5% NaOH solution, dried over Na₂SO₄, and evaporated in a vacuum. The oily residue was subjected to chromatography (eluent methanol–chloroform, 1:9) to isolate 0.05 g (38%) of yellow crystals, mp 170°C. IR spectrum, cm⁻¹: 1650 (C=O), 3280 (NH of amide), 3400 (NH of indole). ¹H NMR spectrum (CDCl₃), δ, ppm: 7.89 br.s (1H, NH of indole), 7.49 d (1H, H^{7'}, *J* 6.82 Hz), 7.31 d (1H, H^{4'}, *J* 7.58 Hz), 7.13 m (2H, H^{5',6'}), 5.67 br.s (1H, NHAc), 4.07 br.s (1H, H⁶, *W*_{1/2} 28.8 Hz), 3.00 br.s (1H), 2.85 d.d (1H, H^{4e}, *J* 6.32, 16.68 Hz), 2.67 d (1H, H^{4a}, *J* 16.68 Hz), 2.57 br.s (1H), 2.07 m (1H), 2.01 s (3H, CH₃), 1.93–1.10 m (5H). ¹³C NMR spectrum (CDCl₃), δ, ppm: 169.44 (C=O), 136.54, 135.63, 127.13, 121.06, 119.18, 117.65, 110.76, 109.69, 51.84 (C⁶), 32.13, 31.16, 30.70, 28.32, 25.11, 23.64, 20.85. Mass spectrum, *m/z* (*I*_{rel}, %): 268 (100) [*M*]⁺, 253, 225, 209, 194, 180, 168.

Monooxime of bicyclo[3.3.1]nonane-2,6-dione (VI) was prepared from the oxime of mono ethylene ketal of bicyclo-[3.3.1]nonane-2,6-dione (see the text) by procedure [5], yield 94%. IR spectrum, cm⁻¹: 3400 br (OH), 1700 (C=O) br, sh 1650 (C=N). Mass spectrum, *m/z* (*I*_{rel}, %): 167 [*M*]⁺, 150 [*M*–OH]⁺. Found, %: C 64.18; H 7.57; N 7.97. C₉H₁₃NO₂. Calculated, %: C 64.65; H 7.83; N 8.37. *M* 167.21.

8-Bromo-5,6,7,8,10,11-hexahydro-9H-6,10-methanocycloocta[b]indol-9-one (VII). To 10 ml of concn. H₂SO₄ was added 1 g (3.70 mmol) of ethylene ketal **IVa** and 1.16 g of silver sulfate, and a flow of bromine vapor (0.6 g) with air was passed through the mixture. The reaction mixture was poured on ice, the precipitate was separated and washed with CHCl₃. The filtrate was extracted with CHCl₃, the organic layer was passed through a bed of silica gel. The solution was evaporated in a vacuum to obtain 0.96 g of dark-brown oily substance which was subjected to chromatography (eluent hexane–ethyl acetate, 2:1) to give ketone **IVb** and compound **VII**. IR spectrum, cm⁻¹: 3400 (NH), 1720 (C=O). ¹H NMR spectrum (CDCl₃), δ, ppm: 8.08 br.s (1H, NH), 7.46 d (1H, *J* 7.80 Hz), 7.37 d (1H, *J* 8.00 Hz), 7.22 t (1H, *J* 7.54 Hz), 7.13 t (1H, *J* 7.34 Hz), 4.49 m (1H, H⁷), 3.39 br.s (1H), 3.23 br.s (1H), 3.15 d.d (1H, H^{4e}, *J* 6.85, 16.38 Hz), 2.83 d (1H, H^{4a}, *J* 16.20 Hz), 2.80–2.18 m (4H). Mass spectrum, *m/z* (*I*_{rel}, %): 305 (30) [*M* + 1]⁺, 303 (27), 226 (35), 225 (83), 224 (75).

2'-Methoxy-5',6',7',8',10',11'-hexahydrospiro-(1,3-dioxolane-2,9'-[6,10]methanocycloocta[b]-indole)

(VIIIa). To a solution of 0.97 g (4.95 mmol) of mono ethylene ketal **VI** in 12 ml of methanol under argon was added at stirring 0.95 g (5.44 mmol) of 4-methoxyphenylhydrazine hydrochloride, 0.376 g (2.72 mmol) of K₂CO₃, and 2.28 g (16.72 mmol) of anhydrous ZnCl₂. The mixture was boiled for 4 h, the precipitate was filtered off, washed with water and methanol. Yield 1.0 g (68%), colorless crystals, mp 160–162°C. IR spectrum, cm⁻¹: 3400 (NH). ¹H NMR spectrum (CDCl₃), δ, ppm: 7.56 br.s (1H, NH), 7.19 d (1H, H^{7'}, *J* 8.84 Hz), 6.95 d (1H, H^{4'}, *J* 2.27 Hz), 6.79 d.d (1H, H^{6'}, *J* 2.28, 8.59 Hz), 4.00 m (4H, OCH₂CH₂O), 3.87 s (3H, OCH₃), 2.97 m (2H), 2.86 d.d (1H, H^{4e}, *J* 6.83, 16.68 Hz), 2.24–1.48 m (7H). ¹³C NMR spectrum (CDCl₃), δ, ppm: 153.90, 137.26, 130.69, 127.92, 111.23, 111.14, 110.67, 109.26, 100.35, 64.33 (OCH₂CH₂O), 55.95 (OCH₃), 36.49, 30.44, 29.80, 28.81, 28.55, 22.52. Found, %: C 72.32; H 7.10; N 4.72. C₁₈H₂₁NO₃. Calculated, %: C 72.24; H 7.02; N 4.68.

2-Methoxy-5,6,7,8,10,11-hexahydro-9H-6,10-methanocycloocta[b]indol-9-one (VIIIb) was obtained by procedure [5] from compound **VIIIa**. Yield 82%, mp 170–172°C. IR spectrum, cm⁻¹: 3390 (NH), 1690 (C=O). ¹H NMR spectrum (CDCl₃), δ, ppm: 7.80 br.s (1H, NH), 7.19 d (1H, H^{7'}, *J* 8.81 Hz), 6.95 d (1H, H^{4'}, *J* 1.96 Hz), 6.79 d.d (1H, H^{6'}, *J* 2.34, 8.81 Hz), 3.87 s (3H, OCH₃), 3.20 br.s (1H), 3.09 d.d (1H, H^{4e}, *J* 6.85, 16.43 Hz), 3.02 br.s (1H), 2.80 d (1H, H^{4'a}, *J* 16.24 Hz), 2.36–2.06 m (6H). ¹³C NMR spectrum (CDCl₃), δ, ppm: 213.68 (C=O), 154.18, 136.04, 130.85, 127.49, 111.55, 111.38, 108.15, 100.24, 55.79 (CH₃), 44.58, 36.65, 32.25, 31.96, 28.45, 25.13. Mass spectrum, *m/z* (*I*_{rel}, %): 255 (100) [*M*]⁺, 238 (5), 227 (12), 212 (10), 198 (100), 183 (25), 167 (20). Found, %: C 75.05; H 7.00; N 5.24. C₁₆H₁₇NO₂. Calculated, %: C 75.29; H 6.67; N 5.49. *M* 255.

(9E)-2-Methoxy-5,6,7,8,10,11-hexahydro-9H-6,10-methanocycloocta[b]indol-9-one oxime (VIIIc) was obtained similarly to oxime **IVc** in 96% yield. The compound was purified by chromatography (eluent benzene–ethyl acetate, 3:1) through a thin layer of silica gel (1 cm). mp 170°C. IR spectrum, cm⁻¹: 3400 (NH), 3200–3450 (OH). ¹H NMR spectrum (CDCl₃), δ, ppm: 7.67 br.s (1H, NH), 7.21 d (1H, H^{7'}, *J* 8.34 Hz), 6.92 d (1H, H^{4'}, *J* 2.53 Hz), 6.82 d.d (1H, H^{6'}, *J* 2.53, 8.59 Hz), 4.12 br.s (1H, NOH), 3.86 s (3H, OCH₃), 3.24–1.61 m (10H). ¹³C NMR spectrum (CDCl₃), δ, ppm: 163.66 (C=N), 154.03, 136.43, 130.78, 128.35, 111.30, 109.09, 109.09, 100.25, 55.91, 34.64, 32.51, 30.31, 29.72, 29.19, 27.51. Found, %: C 71.30; H 6.81; N 10.42. C₁₆H₁₈N₂O₂. Calculated, %: C 71.11; H 6.67; N 10.37.

2-Methoxy-6,7,8,9,10,11-hexahydro-5H-6,10-methanocycloocta[b]indol-9-amine (IX). To a dispersion of 0.14 g (5.00 mmol) of LiAlH_4 in 10 ml of anhydrous THF was added dropwise at stirring a solution of 0.45 g (1.67 mmol) of compound **VIIIc** in 10 ml of THF, and the mixture was boiled for 6.5 h. On cooling with ice 0.8 ml of water–THF mixture, 1:1, was added, the mixture was boiled for 1 h, the precipitate was filtered off and washed with hot THF. The combined filtrates were dried over Na_2SO_4 , and the solution was evaporated on a rotary evaporator to obtain 0.38 g of yellow oily substance that was used in the synthesis of compound **II** without isolation.

The study was carried out under a financial support of the Russian Foundation for basic Research (grant no. 06-03-32843a) and of the Division of Chemistry and Material Sciences of the Russian Academy of Sciences.

REFERENCES

1. Trudell, M.L. and Cook, J.M., *J. Am. Chem. Soc.*, 1989, vol. 111, p. 7504.
2. Fu, X. and Cook, J.M., *J. Org. Chem.*, 1993, vol. 58, p. 661.
3. Bailey, P.D., Morgan, K.M., Smith, D.I., and Vernon, J.M., *J. Chem. Soc., Perkin Trans. 1*, 2000, p. 3566.
4. Bailey, P.D., Morgan, K.M., and Smith, D.I., *J. Chem. Soc., Perkin Trans. 1*, 2000, p. 3578.
5. Kadzyauskas, P.P., Butkus, E.P., Vasyulite, Yu.A., Averina, N.V., and Zefirov, N.S., *Khim. Geterotsikl. Soedin.*, 1979, p. 315.
6. Butkus, E., Berg, U., Malinausiene, J., and Sandstrom, J., *J. Org. Chem.*, 2000, vol. 65, p. 1353.
7. Klusacek, H. and Musso, H., *Chem. Ber.*, 1970, vol. 103, p. 3066.
8. Amat, M., Sanfeliu, E., Bonjoch, J., and Bosch, J., *Tetrahedron Lett.*, 1989, vol. 30, p. 3841.
9. Gilbert, I.M., Hewett, C.L., Ral, D.R., Redpath, J., Savage, D.S., and Sleight, T., *J. Chem. Soc., Perkin Trans. 1*, 1995, 133.
10. Kost, A.N., Yudin, L.G., and Budylin, V.A., *Khim. Geterotsikl. Soedin.*, 1967, p. 16.
11. Kost, A.N., Yudin, L.G., Budylin, V.A., and Abdullaev, M.R., *Khim. Geterotsikl. Soedin.*, 1971, p. 512.