

This article was downloaded by:

On: 25 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Separation Science and Technology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713708471>

Erratum

Tsutomu Kawasaki

To cite this Article Kawasaki, Tsutomu(1982) 'Erratum', Separation Science and Technology, 17: 6, 877

To link to this Article: DOI: 10.1080/01496398208058328

URL: <http://dx.doi.org/10.1080/01496398208058328>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ERRATUM

Gradient Hydroxyapatite Chromatography with Small Sample Loads. III. Fundamental Differential Equation for Gradient Chromatography

TSUTOMU KAWASAKI

[article in *Separation Science and Technology*, 16(7), 817-851 (1981)]

On page 819, the fourth paragraph reads incorrectly. The corrected paragraph is as follows:

For each component of the mixture the ratio $(R_F)_\lambda$ of the migration rate of molecules to the migration rate of the solvent in an elementary volume δV at any column position L' should be equal to the ratio B_λ of the amount of molecules existing in the interstitial liquid (i.e., the mobile phase) to the total amount in that elementary volume. This is a first principle of chromatography in the case of the quasi-static process (1,3). B_λ and the mean value B of B_λ within the column section are both independent of the total amount of molecules in the column section if the amount of molecules is small because, in this situation, the linear section of the adsorption isotherm should be realized. With stepwise chromatography the migration of molecules on the column can, in fact, be described by using Eq. (1) (see Ref. 1, Appendix III).