

This article was downloaded by:

On: 30 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Spectroscopy Letters

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597299>

Preface

N. A. Borisevich; B. I. Stepanov

To cite this Article Borisevich, N. A. and Stepanov, B. I. (1975) 'Preface', *Spectroscopy Letters*, 8: 9, 605

To link to this Article: DOI: 10.1080/00387017508067367

URL: <http://dx.doi.org/10.1080/00387017508067367>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Preface

The present special issue of the journal is concerned with the problem of laser action of complex organic molecules. This scientific problem stands on the borderline of spectroscopy and quantum electronics and is of great interest for scientists in different fields of research. Specialists in lasers are attracted by the possibility of the continuous tuning of laser radiation frequency, of the achievement of laser action in a wide frequency range, and of the creation of broad-band amplifiers. On the other hand, spectroscopists consider this phenomenon from the point of view of utilizing qualitatively new possibilities for determining spectroscopic parameters of substances and especially their nonlinear optical properties.

The editors of the issue endeavoured to draw the largest number of outstanding names in this field of science to the issue. However, not all our attempts proved successful. Nevertheless, we believe that practically all the most important aspects of the problem under consideration have been reflected in the issue. Most of the papers are of a survey nature and generalize the results obtained by large scientific teams.

N.A.Borisevich

B.I.Stepanov