

This article was downloaded by:

On: 30 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Spectroscopy Letters

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597299>

A Study of *cis* - *trans* Isomerism of Benzalide and Thiobenzanilide

B. Bogdanov^a; I. Petrov^a

^a Faculty of Science-Skopje, Institute of Chemistry, Macedonia, Yugoslavia

To cite this Article Bogdanov, B. and Petrov, I.(1992) 'A Study of *cis* - *trans* Isomerism of Benzalide and Thiobenzanilide', Spectroscopy Letters, 25: 2, 239 — 243

To link to this Article: DOI: 10.1080/00387019208020689

URL: <http://dx.doi.org/10.1080/00387019208020689>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A STUDY OF *cis* - *trans* ISOMERISM OF BENZALIDE AND THIOBENZANILIDE

Keywords: Benzanilide, Thiobenzanilide, *cis-trans* Isomerism,
Infrared spectra.

B. BOGDANOV and I. PETROV

Institute of Chemistry, Faculty of Science-Skopje,
91000 Skopje, Macedonia, Yugoslavia

ABSTRACT

The conformations of the benzanilide and thiobenzanilide are studied using IR spectral data and Extended Huckel MO calculations. The results indicate that the benzanilide in CCl₄ dilute solution exist only in a *trans*-conformation, while the thiobenzanilide exist in a *cis-trans* equilibrium. The observed relative stabilities of the *cis*-thiobenzanilide can be explained on the basis of the calculated torsional barriers around the thiocarbonyl-nitrogen bond.

INTRODUCTION

The structure behavior of the amides and thioamides have been the subject of much investigation using the spectroscopic and theoretical studies [1-6]. A more detailed study of their configurations is of importance in any attempt of structure - activity correlation.

The benzanilide (I) and thiobenzanilide (II) could possibly take the *trans* - and *cis* - isomeric forms (Fig. 1). ϕ_1 and ϕ_2 are torsional angles between the plane of the functional group and plane of the phenyl rings bound to N and C atoms. The partially double bond character of the C-N bond from the Ph-C(X)-NH-Ph (X=O or S) which is due to the delocalization of the electronic pair at the nitrogen atom determines the hindered rotation around this




Fig. 1. Molecular conformations of benzanilide (I, X=O) and thiobenzaldehyde (II, X=S). The *cis*- and *trans*-planar conformations ($\phi_1 = \phi_2 = 0^\circ$) are shown.

bond and the possibilities for a relative stabilization of different conformational isomers. Suzuki et al. [1] examined a series of thioformanilides and thioacetanilides and showed that the thioformanilides have predominantly a *cis*-conformation while the thioacetanilides exist in both *cis* and *trans* forms. In this respect the thioanilides evidently differ from anilides which normally have a *trans*-conformation. This difference is attributed on increased steric repulsion between the phenyl ring and the sulfur atom in the thioamides. The torsional barriers in thioamides are in general 0.04-0.13 eV/mol higher than in corresponding amides [4].

In the present work IR spectral data and Extended Hückle MO (EHMO) calculations for the existence of conformational isomerism in benzanilide and thio benzanilide are analysed.

EXPERIMENTAL

Infrared spectra

The benzanilide and the thiobenzaldehyde were synthesized using known method [7]. The infrared spectra were recorded on a Perkin-Elmer 580 IR Spectrophotometer. The benzanilide and the thiobenzanilide as dilute solution in CCl_4 (at concentrations $1.2 \cdot 10^{-3}\text{M}$ and $1.3 \cdot 10^{-3}\text{M}$, respectively) were recorded. The IR spectra in the N-H stretching frequencies range are shown in Fig. 2.




Fig. 2. IR spectrum of benzanilide (I) and thiobenzanilide (II) in CCl_4 .

Method of calculations

The total energies were calculated by means of EHMO method using QCPE No 358. Adopted values of integrals and parametrization are the same as in ref. [8]. The sulfur 3d orbitals were included in the basis set. The bond lengths and angles were set to values given by X-ray analysis [9]; regular benzene rings are assumed.

RESULTS AND DISCUSSION

In the infrared spectrum of benzanilide in carbon tetrachloride dilute solution, the N-H stretching vibration appears as a single monomeric band at 3440 cm^{-1} , while the spectrum of thiobenzaldehyde gives rise to two monomeric bands, a strong one at 3390 cm^{-1} and a weak one at 3365 cm^{-1} (Fig. 2). The appearance of two bands in thiobenzaldehyde indicates that it exists in a *cis-trans* equilibrium. The strong band belongs to *trans*- and the weak to *cis*- configuration.

The conformational preference exhibited by these molecules is clearly governed by the relative importance of two factors, the π -interactions which tend to keep the molecules planar (the conjugative factor), and the non-bonding interactions which may be alleviated by deviation from planarity (the steric factor). We

Table 1

Torsional barriers and potential constants (in eV) of benzanilide (I) and thiobenzanilide (II)

Molecule	conformation Φ_1, Φ_2 (deg.)	<i>cis-trans</i> barrier	V_1	V_2	V_3
I	90, 90 ^a	0.999	-0.105	1.094	-0.085
II	90, 90 ^a	1.130	-0.013	1.176	-0.079
I	90, 90 ^b	1.835	1.568	0.879	0.342
II	90, 90 ^b	2.085	1.857	0.923	-0.396

^aMost stable *cis*-conformation (see Fig. 1).

^bMost stable *trans*-conformation (see Fig. 1).

have carried out MO calculations employing EHMO method to estimate the relative stabilities and the barrier highs to rotation. The torsional barriers around C-N bond for benzanilide and thiobenzanilide are presented in Table 1.

The expansion to a Fourier series of the potential function

$$V(\Phi) = \sum \frac{1}{2} V_{Ni} (1 - \cos Ni \Phi)$$

deserves some interest since it allows various physical effects on the mechanism of rotational barriers to be separated [10]. In particular, the physical effects reflected by the potential constants V_i for the present compounds are the interaction of local dipole and steric hindrance (V_1), the tendency to adopt an uniplanar conformation in order to maximize conjugative stability (V_2), and the bond-bond repulsion (V_3). For most stable *cis*-conformation, the results prove that the two-fold component (Table 1) for benzanilide and thiobenzanilide is most important and largely responsible for the rotational barrier. Furthermore, the V_1 is more important than V_3 , and although smaller than V_2 , plays a decisive role in assessing the relative stabilities of the *cis*-and

trans-conformations. The energy barrier and potential constant V_2 of benzanilide are smaller than those of thiobenzanilide, indicating less mesomeric inter action between oxygen and C-N bond, than those between sulfur and C-N bond. However, as can be seen from Table 1, the most stable *trans*-conformation of benzanilide and thiobenzanilide have V_1 values greater than V_2 indicating that the steric factor play dominant role in the total torsional barrier of those molecules.

ACKNOWLEDGEMENT

This work has been carried out thanks to the support of the Republic Macedonian Science Found.

REFERENCES

- 1 I. Suzuki, M. Tsuboi, T. Shimanouchi, S. Muzishima, *Spectrochim. Acta*, 16 (1960) 471.
- 2 G. Isaksson and J. Sandstrom, *Acta Chem. Scand.*, 21 (1967) 1605.
- 3 C. N. Rao, K. G. Rao, A. Goel and D. Balasubramanian, *J. Chem. Soc. (A)*, (1971) 3077.
- 4 C. Piccinni-Leopardi, O. Fabre, D. Zimmermann and J. Reissue, *Can. J. Chem.* 55 (1977) 2649.
- 5 J. Sandstrom, *Tetrahedron Lett.*, 7 (1979) 639.
- 6 I. Petrov and O. Grupce, *J. Mol. Struct.*, 115 (1984) 481.
- 7 O. Grupce and I. Petrov, *J. Mol. Struct.*, 115 (1984) 119.
- 8 V. Galasso, G. De Alti and A. Bigotto, *Tetrahedron*, 27 (1970) 6151.
- 9 J. Zabicky, *The Chemistry of Amides*, Interscience Publishes, London 1970, p.4 and 387.
- 10 L. Radom, W. J. Hehre and J. A. People, *J. Am. Chem. Soc.*, 94 (1972) 2371.

Date Received: 10/01/91
Date Accepted: 10/30/91