

This article was downloaded by:

On: 30 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Spectroscopy Letters

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597299>

## Subject Index to Volume 32

**To cite this Article** (1999) 'Subject Index to Volume 32', *Spectroscopy Letters*, 32: 6, 1105 — 1117

**To link to this Article: DOI:** 10.1080/00387019909350053

**URL:** <http://dx.doi.org/10.1080/00387019909350053>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SUBJECT INDEX TO VOLUME 32

## A

Absorption spectra, of Schiff bases, role of solvent acidity and basicity in the solvent effect in, 57–71

Acetonitrile-D<sub>3</sub>-cation interactions, FTIR investigation of, 783–792

Acidity constants, of 2-monosulfohydroxybenzene, 333–346

3-Acylaminoquinazolin-4(3H)-one derivatives, proton NMR spectroscopy of barriers to rotation about N-N bonds in, 671–677

Adsorption, of  $\omega$ -mercaptoalkanols on silver surfaces, FT-Raman SERS of, 281–288

Alkaline isomerization, of cytochrome c NMR spectroscopy of the effects of nitrogenous ligands on, 519–533

in the presence of  $\gamma$ -picoline, <sup>1</sup>H NMR of, 305–314

Amino- and aminoalkylpyridines, fluorescence characteristics of Schiff bases derived from, 181–196

2-Aminopyridine, infrared spectra of complexes with palladium(II), 165–180

Aniline, infrared spectra of complexes with palladium(II), 165–180

Antibacterial sesquiterpene glycoside, from *Biebersteinia heterostemon*, 1005–1012

Antihistamines, spectrophotometric analysis of, 913–930

Artificial neural networks multivariate calibration, application to near-infrared spectrophotometric determination of powdered pharmaceutical metronidazole, 431–442

use in FTIR spectroscopy for simultaneous determination of multicomponents in air toxic organic compounds, 421–429

6-Aryl-3-(1-*p*-chlorophenyl-5-methyl-1,2,3-triazol-4-yl)-*s*-triazolo-[3,4-*b*]-1,3,4-thiadiazoles, <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy of, 1025–1032

4-Arylidene-2-aryl-5-oxazolones(az-lactones), structure and luminescent properties in solution and the crystalline state, 553–569

## B

Benzylidenemalononitrile series, infrared studies of the validity of constants of ionic substituents in, 793–802

BiD molecule, transition probabilities and dissociation energy of, 31–34

*Biebersteinia heterostemon*, antibacterial sesquiterpene glycoside and other bioactive compounds from, 1005–1012

Binuclear copper(II) imidazole saccharinato complex, FTIR spectra of, 237–256

Bioactive compounds, from *Biebersteinia heterostemon*, 1005–1012

Biological materials, texture of stretched and unstretched polymer films with and without, 629–637

4,4'-Bipyridine, structure and conformation of, 35–45

Bis(ethylenediamine)-diisothiocyanatonickel(II)-benzene, spectroscopic and structural studies of clathrate compound of, 397–405, 763–771

Bond moments, of nitromethane, 347–359

## C

$C_{60}$ , thermal lens spectra of, 535–541

Cadmium infrared study of the heating effects of complexes with sulfadimethoxine, 729–735

nuclei, IBM-2 calculation of, 803–817

simultaneous determination with lead in cigarettes and cigarette smoke by impaction-graphite furnace and graphite furnace atomic absorption spectrometry, 615–621

cAMP, laser Raman spectroscopy of effects on membrane phospholipid of human erythrocytes, 197–203

Casting-on-water method, photo-acoustic studies of PMMA very thin films prepared by, 371–382

$CCl_4$  solution, aqueous, nanosecond laser flash photolysis transient spectra of, 983–992

Cellulose, ultraviolet spectroscopy of functionalization with silanes, 993–1003

Charge transfer complexes, formed in the reaction of  $\pi$ -electron acceptors with the donor cyclic base 1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, 115–124

Chemiluminescence, inverted micellar enhanced organic peroxyxalate, 273–279

Cigarettes and cigarette smoke, impaction-graphite furnace and graphite furnace atomic absorption spectrometry simultaneous determination of cadmium and lead in, 615–621

Clathrate compound, of bis(ethylenediamine)-diisothiocyanatonickel(II)-benzene, spectroscopic and structural studies of, 397–405, 763–771

Cobalt, infrared study of the heating effects of complexes with sulfadimethoxine, 729–735

Cobalt(II), cationic, complexes with isatin/triphenylphosphine mixed ligands, 315–319

Copper(II) imidazole saccharinato complex, FTIR spectra of, 237–256

Copper<sup>2+</sup>-doped sodium hydrogen oxalate monohydrate single crystal, EPR study of, 571–579

Coumarin(VII), substituted, thermodynamics of, 383–395

Cresylviolet perchlorate, Stark hole-burning spectroscopy in amorphous hosts, 125–137

Cyanoiron(II)-polyimine supermolecule, pentanuclear, and its corresponding Prussian blue film, spectroscopic and electrochromic behavior of, 963–971

Cytochrome c  
<sup>1</sup>H NMR of alkaline isomerization in the presence of  $\gamma$ -picoline, 305–314  
 NMR spectroscopy of the effects of nitrogenous ligands on the alkaline isomerization of, 519–533

**D**

1,4-Diaminoanthraquinone dye, SERS in silver colloid, 103–113

1,2:5,6-Dibenzanthracene, application of time-resolved linear dichroism spectroscopy to, 747–762

Differential cross sections, for coherent scattering of 59.5 keV  $\gamma$ -rays, 623–628

10,11-Dihydro-5H-dibenzo[a,d]cycloheptene-5-one and 9,10-dihydro-4H-benzo [4,5]cyclohepta[1,2-b]thiophene-4-one, substituted, mass spectral fragmentation patterns of, 601–613

Dipeptide Schiff bases, metal complexes with, 679–688

Di-2-pyridyl ketone benzoylhydrazone, for solvent extraction and spectrophotometric determination of nickel(II) in natural waters, 257–271

Dissociation energy, of BiD molecule, 31–34

*N*-(2,6-Di-tert-butyl-1-hydroxyphenyl) salicylaldimines, synthesis, spectroscopic characterization, and redox reactivity of, 485–495

DNA, binding of ruthenium(II) complex to, 931–939

Double layer, secondary photoemission through, 205–212

Dye, 1,4-diaminoanthraquinone, SERS in silver colloid, 103–113

**E**

EDXRF. *See* Energy dispersive X-ray fluorescence

$\pi$ -Electron acceptors, spectroscopy of the charge transfer complexes formed in the reaction with the donor cyclic base 1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, 115–124

Electron paramagnetic resonance (EPR), of copper<sup>2+</sup>-doped sodium hydrogen oxalate monohydrate single crystal, 571–579

Electronic interaction studies, of *p*-substituted  $\alpha$ -phenoxyacetones, 1041–1056

Electronic spectra, of the *B*–*X* system of isotopic silver oxide molecules, 875–882

Electrospray mass spectrometry, for investigation of heteronuclear terbium–zinc cryptate, 1033–1039

Energy dispersive X-ray fluorescence (EDXRF), of lead pollution, 661–669

2,3-Epoxypropyl or 3-amino-2-hydroxypropyl moiety, NMR characterization of purine nucleoside analogues with, 649–660

EPR. *See* Electron paramagnetic resonance

Equilibrium dialysis, of the interaction between trypanocide 3-nitrobenzaldehyde guanyl hydrazone and bovine serum albumin, 941–951

Europium

- ions, variation in the luminescent lifetimes of, 737–746
- luminescent properties of hexakis(nitrito)europate(III) ion, 155–163

Extended X-ray absorption fine structure spectroscopy, of the structure of platinothioneins, 883–894

F

Films

- N*-octadecanoyl-L-alanine and zinc octadecanoyl-L-alaninate LB, FTIR spectroscopy of phase transitions of, 1–16
- polymer mixtures, multivariate modeling with data augmentation and infrared, Raman, and near-infrared spectroscopy for quantification of polyethylene types in, 1057–1071
- Prussian blue, of a pentanuclear cyanoiron(II)-polyimine supermolecule, 963–971
- stretched and unstretched polymer, texture with and without embedded biological materials, 629–637
- very thin PMMA prepared by a casting-on-water method, photoacoustic studies of, 371–382

Fluorescence behavior, of the lithium-tetracycline system, influence of solvent on, 73–82

Fluorescence characteristics, of Schiff bases derived from amino- and aminoalkylpyridines, 181–196

Fourier transform infrared (FTIR) intensities, of nitromethane, 347–359

spectra

- of binuclear copper(II) imidazole saccharinato complex, 237–256
- of glutarimide, 47–56

spectroscopy  
 of acetonitrile-D<sub>3</sub>-cation interactions, 783–792  
 of the hydrolysis of titanium tetrabutoxide, 289–304  
 of phase transitions of *N*-octadecanoyl-L-alanine and zinc octadecanoyl-L-alaninate LB films, 1–16  
 using artificial neural networks, for simultaneous determination of multicomponents in air toxic organic compounds, 421–429

Fourier transform vibrational spectra, of magnesium hydrogenphosphate trihydrate, 703–717

Fourier transform-Raman SERS, of adsorption of  $\omega$ -mercaptolkanols on silver surfaces, 281–288

FTIR. *See* Fourier transform infrared

Fulvic acid, adsorbed on copper electrodes, SERS of, 689–701

**G**

Gadolinium(III) diethyltriamine-pentaacetic acid protonation constants, determination by solvent extraction and ICP-AES, 463–467

Glutarimide, FTIR and Raman spectra of, 47–56

Glycoside, antibacterial sesquiterpene, from *Biebersteinia heterostemon*, 1005–1012

Graphite furnace atomic absorption spectrometry, simultaneous determination of cadmium and lead in cigarettes and cigarette smoke by, 615–621

**H**

Hafnium monohalides, spectroscopy of, 719–728

Haloacetylated compounds, 1,1,1-trichloro-4-methoxy-3-alken-2-ones, 973–981

Halochromic properties, of *cis*-Ru(bpy)<sub>2</sub>Cl<sub>2</sub> in organic solvents, 497–504

Heating effects, of nickel, cadmium, and cobalt complexes of sulfadimethoxine, infrared study of, 729–735

Heterocycles, molecular structure of, 851–865

Heterocyclic aldehydes, nitrogen containing, stereochemistry of, 581–600

Heterocyclic compounds, condensed, 1025–1032

Heteronuclear terbium–zinc cryptate, electrospray mass spectrometric investigation of, 1033–1039

Hexakis(nitrito)europate(III) ion, luminescent properties of, 155–163

Humic acid, adsorbed on copper electrodes, SERS of, 689–701

Hydrogen bonding, and the protonated water dimer, 819–828

Hydrogen-bond proton acceptors, tetrafluoroberrylyate(2<sup>-</sup>) ions as, 361–369

Hydrolysis, of titanium tetrabutoxide, FTIR spectroscopy of, 289–304

**I**

IBM-2. *See* Interacting boson model-2

ICP-AES. *See* Inductively coupled plasma-atomic emission spectrometry

Impaction-graphite furnace atomic absorption spectrometry, simultaneous determination of cadmium and lead in cigarettes and cigarette smoke by, 615-621

Inductively coupled plasma-atomic emission spectrometry (ICP-AES), for determination of gadolinium(III) diethyltriaminepentaacetic acid protonation constants, 463-467

Infrared spectra, of 2-aminopyridine and aniline complexes with palladium(II), 165-180

Infrared spectroscopy, for average molecular weight determination of polysulfide polymers, 773-782

Infrared study  
of complex between salicylalbenzoylhydrazone and iron(III), 505-517  
of the heating effects of nickel, cadmium, and cobalt complexes of sulfadimethoxine, 729-735  
of the validity of constants of ionic substituents in the benzylidenemalononitrile series, 793-802

Interacting boson model-2 (IBM-2), for calculation of cadmium nuclei, 803-817

Inverted micellar enhanced organic peroxyoxalate chemiluminescence, 273-279

Ionic substituents, in the benzylidenemalononitrile series, infrared studies of the validity of constants of, 793-802

Iron-di- and poly-carboxylated-pyrocatechol and pyrogallol mixed ligands chelates, 407-419

Iron(III)  
cationic complexes with isatin/triphenylphosphine mixed ligands, 315-319  
infrared study of complex with salicylalbenzoylhydrazone, 505-517

Isatin/triphenylphosphine mixed ligands, cationic iron(III), cobalt(II), and nickel(II) complexes with, 315-319

Isotopic silver oxide molecules, vibrational structure of the *B-X* system of, 875-882

**J**

Judd-Ofelt analysis, of photoacoustic spectra of powdered neodymium compounds, 543-551

**L**

Langmuir-Blodgett (LB) films, *N*-octadecanoyl-L-alanine and zinc octadecanoyl-L-alaninate, FTIR spectroscopy of phase transitions of, 1-16

Laser flash photolysis, nanosecond, of the transient spectra of aqueous  $\text{CCl}_4$  solution, 983-992

Laser Raman spectroscopy, of effects of cAMP on membrane phospholipid of human erythrocytes, 197–203

LB. *See* Langmuir-Blodgett

Lead

- EDXRF of pollution by, 661–669
- simultaneous determination with cadmium in cigarettes and cigarette smoke by impaction-graphite furnace and graphite furnace atomic absorption spectrometry, 615–621

Lithium-tetracycline system, influence of solvent on the fluorescence behavior of, 73–82

Luminescence

- of strontium borates, 895–910
- of terbium<sup>3+</sup>-doped strontium quinolate, 867–873

Luminescence lifetimes, variation for europium<sup>3+</sup> ion, 737–746

Luminescent properties of 4-arylidene-2-aryl-5-oxazolones(azlactones), 553–569

of hexakis(nitrito)europate(III) ion, 155–163

**M**

Magnesium hydrogenphosphate trihydrate, Fourier transform vibrational spectra of, 703–717

Magnetic field profiling, for SSMRS and MGRS in condensed matter, 1013–1024

Manganate(VI) intermediate, short lived, spectrophotometric detection through oxidation of natural and synthetic polymers by alkaline permanganate, 17–29

Mass spectral fragmentation patterns, of substituted 10,11-dihydro-5H-dibenzo[a,d]cycloheptene-5-one and 9,10-dihydro-4H-benzo[4,5]cyclohepta[1,2-b]thiophene-4-one, 601–613

Membrane phospholipid, of human erythrocytes, laser Raman spectroscopy of effects of cAMP on, 197–203

ω-Mercaptoalkanols, FT-Raman SERS of adsorption on silver surfaces, 281–288

Metal complexes

- with dipeptide Schiff bases, 679–688
- with Schiff bases derived from sulfa drugs, 213–236

Methacrylic acid, liquid polysulfide oligomers based on, 321–331

Methylene blue method, for spectrophotometric determination of on-line dissolution of zinc sulfide for sulfide determination in stabilized water, 469–484

4-Methylesculetin azo compounds, synthesis and analytical application with transition metal ions, 829–840

Methylpyridinocobaloxime, optical probe detection method for study in aqueous solution, 639–647

Metronidazole, powered pharmaceutical, application of artificial neural network multivariate calibration to near-infrared spectrophotometric determination of, 431–442

**MGRS.** *See* Magnetic field gradient spectroscopy, in condensed matter, magnetic field profiling for, 1013–1024

**Mixed ligand chelates, iron,** 407–419

**complexes, of isatin/triphenylphosphine mixed ligands and cationic iron(III), cobalt(II), and nickel(II),** 315–319

**Molecular weight, average determination for polysulfide polymers using infrared spectroscopy,** 773–782

**Monohydroxybenzene, sulfonation of,** 333–346

**2-Monosulfohydroxybenzene, acidity constants of,** 333–346

**Multicomponents, simultaneous determination in air toxic organic compounds using artificial neural networks in FTIR spectroscopy,** 421–429

**Multivariate modeling with data augmentation and infrared, Raman, and near-infrared spectroscopy, for quantification of polyethylene types in polymer film mixtures,** 1057–1071

**N**

**Near-infrared spectrophotometry, for determination of powdered pharmaceutical metronidazole, application of artificial neural network multivariate calibration to,** 431–442

**Neodymium compounds, powdered, Judd-Ofelt analysis of photoacoustic spectra of,** 543–551

**Nickel, infrared study of the heating effects of complexes with sulfadimethoxine,** 729–735

**Nickel(II)**

- cationic complexes with isatin/triphenylphosphine mixed ligands,** 315–319
- solvent extraction and spectrophotometric determination in natural waters using di-2-pyridyl ketone benzoylhydrazone,** 257–271

**3-Nitrobenzaldehyde guanyl hydrazone trypanocide, equilibrium dialysis and NMR of the interaction with bovine serum albumin,** 941–951

**Nitrogen-containing heterocyclic aldehydes, stereochemistry of,** 581–600

**Nitrogenous ligands, NMR spectroscopy of the effects on the alkaline isomerization of cytochrome c,** 519–533

**Nitromethane, FTIR intensities, bond moments, and its derivatives,** 347–359

**NMR.** *See* Nuclear magnetic resonance

**Nuclear magnetic resonance (NMR) for characterization of purine nucleoside analogues with 2,3-epoxypropyl or 3-amino-2-hydroxypropyl moiety,** 649–660

**of the effects of nitrogenous ligands on the alkaline isomerization of cytochrome c,** 519–533

**$^1\text{H}$ , of alkaline isomerization of cytochrome c in the presence of  $\gamma$ -picoline,** 305–314

<sup>1</sup>H and <sup>13</sup>C  
6-aryl-3-(1-*p*-chlorophenyl-5-  
methyl-1,2,3-triazol-4-yl)-s-tri-  
azolo[3,4-*b*]-1,3,4-thiadiazoles,  
1025-1032  
of disubstituted thiophenes,  
443-462

of the interaction between trypano-  
cide 3-nitrobenzaldehyde  
guanyl hydrazone and bovine  
serum albumin, 941-951

<sup>17</sup>O chemical shifts, of 1,1,1-tri-  
chloro-4-methoxy-3-alken-2-  
ones, 973-981

proton, of barriers to rotation about  
N-N bonds in 3-acylaminoquin-  
azolin-4(3H)-one derivatives,  
671-677

## O

*N*-Octadecanoyl-L-alanine LB films,  
FTIR spectroscopy of phase  
transitions of, 1-16

Oligomers, liquid polysulfide, based  
on sulfur, styrene, and  
methacrylic acid, 321-331

On-line dissolution, of zinc sulfide for  
sulfide determination in  
stabilized water samples,  
469-484

Optical probe detection, of photoly-  
sis of methylpyridinocobal-  
oxime in aqueous solution,  
639-647

Organic compounds, simultaneous  
determination in the air by  
FTIR spectroscopy using  
artificial neural networks,  
421-429

Oxidation, of natural and synthetic  
polymers by alkaline perman-  
ganate for spectrophotometric  
detection of short-lived  
manganate(VI), 17-29

## P

Paint pigments, ancient, Raman  
microscopy of, 841-850

Palladium(II), infrared spectra of  
complexes with 2-amino-  
pyridine and aniline, 165-180

Pentanuclear cyanoiron(II)-polyimine  
supermolecule, and its  
corresponding Prussian blue  
film, spectroscopic and  
electrochromic behavior of,  
963-971

Phase transitions, of *N*-octadecanoyl-  
L-alanine and zinc octa-  
decanoyl-L-alaninate LB films,  
FTIR spectroscopy of, 1-16

$\alpha$ -Phenoxyacetones, *p*-substituted,  
conformational and electronic  
interaction studies of,  
1041-1056

3-Phenyl-5-hydroxy-5-trichlorometh-  
yl-4,5-dihydro-1*H*-pyrazole-1-  
carboxyamide, molecular  
structure of, 851-865

Photoacoustic spectra, of powdered  
neodymium compounds, Judd-  
Ofelt analysis of, 543-551

Photoacoustic studies, of very thin  
PMMA films prepared by a  
casting-on-water method,  
371-382

Photoemission, secondary, through  
the double layer, 205-212

Photolysis, of methylpyridinocobaloxime in aqueous solution, an optical probe detection method for study of, 639–647

$\gamma$ -Picoline,  $^1\text{H}$  NMR of alkaline isomerization of cytochrome c in the presence of, 305–314

Piroxicam, vibrational spectrum of a platinum complex of, 953–961

Platinothioneins, extended X-ray absorption fine structure spectroscopy of structure of, 883–894

Platinum complex, of piroxicam, vibrational spectrum of, 953–961

PMMA. *See* Poly(methyl methacrylate)

Polyethylene, low density, linear low density, and high density, multivariate modeling with data augmentation and infrared, Raman, and near-infrared spectroscopy, quantification in polymer film mixtures, 1057–1071

Polymer films, stretched and unstretched, texture with and without embedded biological materials, 629–637

Polymers, natural and synthetic, spectrophotometric detection of short-lived manganate(VI) intermediate through alkaline permanganate oxidation of, 17–29

Poly(methyl methacrylate) (PMMA) very thin films, prepared by a casting-on-water method, photoacoustic studies of, 371–382

Polysulfide liquid oligomers, based on sulfur, styrene, and methacrylic acid, 321–331

polymers, infrared spectroscopy determination of average molecular weight of, 773–782

Protonated water dimer, and hydrogen bonding, 819–828

Protonation constants, of gadolinium(III) diethyltriaminepentacetic acid, determination by solvent extraction and ICP-AES, 463–467

Purine nucleoside analogues, NMR characterization with 2,3-epoxypropyl or 3-amino-2-hydroxypropyl moiety, 649–660

**R**

Raman microscopy, of the ancient paint pigments, 841–850

Raman spectra, of glutarimide, 47–56

Raman spectroscopy, laser, of effects of cAMP on membrane phospholipid of human erythrocytes, 197–203

Rare earth ions, in strontium borates, reduction of, 895–910

Rhodanine(IV), substituted, thermodynamics of, 139–153

Ruthenium complex, halochromic properties in organic solvents, 497–504

Ruthenium(II) complex, binding to DNA, 931–939

## S

Salicylalbenzoylhydrazone, infrared study of complex with iron(III), 505–517

Schiff bases  
derived from amino- and aminoalkylpyridines, fluorescence characteristics of, 181–196  
derived from sulfa drugs, studies of metal complexes with, 213–236  
dipeptide, metal complexes with, 679–688  
role of solvent acidity and basicity in the solvent effect in absorption spectra of, 57–71

Secnidazole, stability-indicating assay in the presence of its degradation products, 1073–1098

Secondary photoemission, through the double layer, 205–212

Selected-states magnetic-resonance spectroscopy (SSMRS), in condensed matter, magnetic field profiling for, 1013–1024

SERS. *See* Surface enhanced Raman scattering

Silanes, ultraviolet spectroscopy of cellulose functionalization with, 993–1003

Silver  
colloid, SERS of 1,4-diamino-anthraquinone dye in, 103–113  
surfaces, FT-Raman SERS of the adsorption of  $\omega$ -mercapto-alkanols on, 281–288

Silver oxide molecules, isotopic, vibrational structure of the  $B$ – $X$  system of, 875–882

Sodium hydrogen oxalate mono-hydrate single crystal, copper<sup>2+</sup>-doped, EPR study of, 571–579

Sodium hydrogen sulfate crystals, vibrational studies of modifications of, 83–102

Solvent  
acidity and basicity, role in the solvent effect in the absorption spectra of Schiff bases, 57–71  
effects on the <sup>17</sup>O NMR chemical shifts of 1,1,1-trichloro-4-methoxy-3-alken-2-ones, 973–981  
influence on the fluorescence behavior of the lithium-tetracycline system, 73–82  
organic, halochromic properties of ruthenium complex in, 497–504

Solvent extraction  
and ICP-AES, for determination of the protonation constants of gadolinium(III) diethyltriaminepentaacetic acid, 463–467  
and spectrophotometry, for determination of nickel(II) in natural waters using di-2-pyridyl ketone benzoylhydrazone, 257–271

Spectrophotometry  
for analysis of antihistamines, 913–930  
for determination of nickel(II) in natural waters using di-2-pyridyl ketone benzoylhydrazone, 257–271  
near infrared, for determination of powdered pharmaceutical metronidazole, 431–442

SSMRS. *See* Selected-states magnetic-resonance spectroscopy

Stability-indicating assay, of secnidazole in the presence of its degradation products, 1073–1098

Stark hole-burning spectroscopy, of cresylviolet perchlorate in amorphous hosts, 125–137

Stereochemistry, of nitrogen-containing heterocyclic aldehydes, 581–600

Strontium borates, luminescence of, 895–910

Strontium quinolinate, terbium<sup>3+</sup> doped, luminescence of, 867–873

Styrene, liquid polysulfide oligomers based on, 321–331

Sulfa drugs, study of metal complexes with Schiff bases derived from, 213–236

Sulfadimethoxine, infrared study of heating effects of nickel, cadmium, and cobalt complexes of, 729–735

Sulfide, determination in stabilized water samples by on-line dissolution of zinc sulfide, 469–484

Sulfonylation, of monohydroxybenzene, 333–346

Sulfur, liquid polysulfide oligomers based on, 321–331

Surface enhanced Raman scattering (SERS) of 1,4-diaminoanthraquinone dye in silver colloid, 103–113

Fourier transform-Raman, of the adsorption of  $\omega$ -mercapto-alkanols on silver surfaces, 281–288

of fulvic and humic acids adsorbed on copper electrodes, 689–701

**T**

Terbium<sup>3+</sup>-doped strontium quinolate, luminescence of, 867–873

Terbium–zinc cryptate, heteronuclear, electrospray mass spectrometric investigation of, 1033–1039

Tetrafluoroberrylyate(2<sup>-</sup>) ions, as hydrogen-bond proton acceptors, 361–369

1,4,10,13-Tetraoxa-7,16-diazacyclooctadecane donor cyclic base, spectroscopy of the charge transfer complexes formed in the reaction with  $\pi$ -electron acceptors, 115–124

Texture, of stretched and unstretched polymer films with and without embedded biological materials, 629–637

Thermal lens spectra, of C<sub>60</sub>, 535–541

Thermodynamics of substituted coumarin(VII), 383–395

of substituted rhodanine(IV), 139–153

Thiophenes, disubstituted, synthesis, NMR, and X-ray crystal structure determination of, 443–462

Time-resolved linear dichroism spectroscopy, application to excited singlet state of 1,2:5,6-dibenzanthracene, 747–762

Titanium tetrabutoxide, FTIR spectroscopy of hydrolysis of, 289–304

3-(*p*-Tolylsulfonamido)rhodanine  
transition metal complexes,  
potentiometric studies of,  
139–153

Transition metal  
complexes, 3-(*p*-tolylsulfonamido)-  
rhodanine, potentiometric  
studies of, 139–153

ions, synthesis and analytical  
application of the complexes of  
4-methylesculetin azo com-  
pounds with, 829–840

Transition probabilities, of BiD  
molecule, 31–34

1,1,1-Trichloro-4-methoxy-3-alken-2-  
ones, solvent effects on the  $^{17}\text{O}$   
NMR chemical shifts of,  
973–981

Trypanocide 3-nitrobenzaldehyde  
guanyl hydrazone, equilibrium  
dialysis and NMR of the  
interaction with bovine serum  
albumin, 941–951

U

Ultraviolet spectroscopy, of cellulose  
functionalization with silanes,  
993–1003

V

Vibrational spectrum, of a platinum  
complex of piroxicam, 953–961

Vibrational studies, of sodium hydro-  
gen sulfate crystals, 83–102

X

X-ray crystal structure, of disubsti-  
tuted thiophenes, 443–462

Z

Zinc octadecanoyl-L-alaninate LB  
films, FTIR spectroscopy of  
phase transitions of, 1–16