

This article was downloaded by:

On: 30 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## **Spectroscopy Letters**

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597299>

## **Subject Index to Volume 33**

**To cite this Article** (2000) 'Subject Index to Volume 33', *Spectroscopy Letters*, 33: 6, 979 — 990

**To link to this Article:** DOI: 10.1080/00387010009350174

**URL:** <http://dx.doi.org/10.1080/00387010009350174>

**PLEASE SCROLL DOWN FOR ARTICLE**

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SUBJECT INDEX TO VOLUME 33

A

- Absorption spectra  
 charge-transfer complex, application of the Mulliken two-state model to, 595-614  
 of mixed chiral amino acid porphyrin Langmuir-Blodgett films, 633-642  
 Acetazolamide,  $^1\text{H}$  NMR spectroscopy for quantitative analysis in solid dosage forms, 661-669  
 Adipokinetic hormones, insect, NMR of, 875-891  
 Adriamycin, spectroscopic characterization of binding to DNA, 693-704  
 Adsorption states, SERS for study of, 69-81  
 Aggregation behavior, of azoanthracene compounds in solution, 359-367  
 Alkali feldspars, luminescence spectra of, 103-113  
 $\alpha$ -Amylase, porcine pancreatic, bromine-81 NMR of anion binding to, 893-899  
 Analgetic mixture, multicomponent, the second-derivative spectrophotometric assay of, 173-183

- Anion binding, to porcine pancreatic  $\alpha$ -amylase, bromine-81 NMR of, 893-899  
 Antarctica, determination of heavy metals in lichens growing on different ecological habitats in, 435-444  
 Antibacterial bisabolane sesquiterpene, assignment of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra by two-dimensional NMR techniques, 235-243  
 Azoanthracene compounds, aggregation behavior in solution, 359-367  
 Azobenzene derivatives, mercaptotended, RA-IR studies in self-assembled monolayers, 949-962

B

- Barium(II) complexes, spectral techniques for study of, 909-916  
 Benzene chromophore, intensity behavior in the near-ultra-violet spectra of chloro and fluoro derivatives of, 681-692

Benzene liquid derivatives, viscosity and density versus temperature compared with  $^1\text{H}$  NMR relaxation time results for, 917-930

Benzophenones, polymethoxylated,  $^1\text{H}$  and  $^{13}\text{C}$  NMR determination of formation of polysubstituted diphenylmethane dimers by reduction of, 755-775

*N*-(1-Benzoyl-3-pyrrolidinyl) benzamide, structural study of, 777-793

Benzylidene aniline, vibrational spectra of complexes with copper(II) and nickel(II) metal ions, 523-533

Binding, of adriamycin to DNA, 693-704

Bisabolane sesquiterpene, antibacterial, assignment of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra by two-dimensional NMR techniques, 235-243

Blood, NMR comparative study of water  $T_1$  and  $T_2$  relaxation times in, 115-125

4,4'-Butane-(1,4,7-three-*p*-tolylsulphonyl-1,4,7-three amine) diphthalonitrile, structure of, 245-254

## C

Cadmium, separated selectively with ion exchange method from solution by inductively coupled plasma atomic emission spectrometry, 195-200

Calixresorcin[4]arene, rare earth ions encapsulated water soluble functional, photophysical properties of, 705-715

Carbon dioxide molecule, sequence and hot transitions in, 283-300

Cephalosporins, use of cerium(IV) in the spectrophotometric and spectrofluorimetric determination in pharmaceutical preparations of, 931-948

Cerium(IV), use in the spectrophotometric and spectrofluorimetric determinations of penicillins and cephalosporins in their pharmaceutical preparations, 931-948

Charge-transfer complex, absorption spectra of, 595-614

Chiral amino acid porphyrin Langmuir-Blodgett films, mixed, absorption spectral study of, 633-642

$\text{C}_2\text{H}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  single crystal, gamma irradiated, electron spin resonance of, 735-742

Chromophores

benzene, intensity behavior in the near-ultraviolet spectra of chloro and fluoro derivatives of, 681-692

pyridine, intensity behavior of the long wavelength UV transition under substitution in the para position to the nitrogen atom, 615-623

Cobalt clathrate compound, spectroscopic and structural studies of, 147-155

Cobalt(II) cupferronato derivatives, Raman studies of, 857-865

- Complex spin system, from a *spiro*-  
chamigrene, NMR analysis of,  
457–467
- Conducting surfaces, spectral de-  
pendencies of surface photo-  
charge effect at, 393–398
- Copper complex, electronic absorp-  
tion spectrum of, 185–194, 971
- Copper(II) complexes  
of bidentate Schiff base ligands,  
comparison of the ultraviolet-  
visible spectra of, 445–455  
dinuclear, photoacoustic spec-  
troscopy of a pair of hydrate  
isomers for, 847–856  
with *o*-hydroxybenzylidene  
*o*-hydroxyaniline, vibrational  
spectra of, 523–533
- Correlation spectroscopy (COSY),  
gradient selected constant time,  
1–8
- COSY. *See* Correlation spectroscopy
- Crude oils, three-dimensional fluor-  
escence spectra of, 963–970
- Crushing, influence on the ultraviolet  
emission band, 103–113
- CTAB micelle aqueous solutions,  
electric spectra of tetra-hydro-  
phenyl porphyrin with two  
hexadecyl chains in, 157–172
- Cupferranoto complexes, transition  
metal, vibrational behavior of,  
857–865

## D

- Diphenylbicyclo[2.2.1]hept-5-ene-  
2,3-dicarboxylic anhydrides,  
isomeric,  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  spec-  
tral parameters and geometries  
of, 717–734

- Diphenylmethanes  
dimers, polysubstituted,  $^1\text{H}$  and  $^{13}\text{C}$   
NMR determination of the  
mechanism of formation by  
reduction of polymethoxylated  
benzophenones, 755–775  
1-naphthyl-polymethoxylated,  $^1\text{H}$   
and  $^{13}\text{C}$  NMR determination of,  
211–226
- DNA, spectroscopic characterization  
of binding of adriamycin to,  
693–704

## E

- Ecological habitats in Schirmacher  
Oasis, East Antarctica,  
determination of heavy  
metals in lichens growing on,  
435–444
- EDTA titration, for spectrophoto-  
metric determination of di- and  
trihydric phenols, 347–357
- Electric spectra, of tetra-hydrophenyl  
porphyrin with two hexadecyl  
chains in CTAB micelle  
aqueous solutions, 157–172
- Electron paramagnetic resonance, of  
the chemical form of mangan-  
ese<sup>2+</sup> in the Chinese loess  
samples, 201–209
- Electron spin resonance, of gamma  
irradiated  $\text{C}_2\text{H}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  single  
crystal, 735–742
- Electronic absorption spectrum  
of cobalt clathrate compound,  
147–155  
of a copper complex, 185–194, 971
- Electronic spectra, of 6-(2-pyridylzo)-  
3-acetamidophenol and its  
metal complexes, 821–832

- Electrospray mass spectrometry  
 of complexation of transitional  
 metal ions with oligobipyridine  
 ligand, 35–46  
 of europium(III) supramolecules,  
 399–406  
*See also* Mass spectrometry  
 Enalapril maleate, spectrophotometric  
 determination of, 91–102  
 Erythrocyte membrane, human,  
 effects of *N*-phosphoprolin on  
 the phospholipid of, 653–659  
 1-Ethyl-1,2,4-triazole, spectral and  
 magnetic studies of complexes  
 with Co(II), Ni(II), Cu(II),  
 Zn(II), and Cd(II), 23–34  
 Europium(III) supramolecules,  
 electrospray mass spectro-  
 metric study of, 399–406

## F

- Far infrared spectroscopy, of  
 complexes of Co(II), Ni(II),  
 Cu(II), Zn(II), and Cd(II) with  
 1-ethyl-1,2,4-triazole and  
 1-propyl-1,2,4-triazole, 23–34  
 Fast atom bombardment mass spec-  
 trometry, formation of the  
 $[M + MNH_4]^+$  ion of oligosac-  
 charides in, 469–473  
 Fats, influence of lanolin and white  
 soft paraffin on determination  
 of vitamin A in, 227–234  
 Feldspars, alkali, luminescence  
 spectra of, 103–113  
 Films, Langmuir-Blodgett  
 of amphiphile with Schiff base as  
 headgroup and its copper  
 complex, Fourier transform  
 infrared studies of, 301–321  
 mixed chiral amino acid porphyrin,  
 absorption spectral study of,  
 633–642  
 Fluorescence spectra  
 of (Ln, Zn)OF:Eu<sup>3+</sup> powder  
 phosphors, effect of zinc ions  
 on, 423–434  
 three dimensional, of crude oils,  
 963–970  
 Fluorimetry, for determination  
 of methoxamine HCl in  
 bulk drugs and in ampoules,  
 9–21  
 4-Fluorophenylpyruvic acid, X-ray  
 crystallography and vibrational  
 spectroscopy of polymorphism  
 of, 509–522  
 Fourier transform infrared spectra,  
 of 4-vinylpyridine and its  
 transition metal(II) tetra-  
 cyanonickelate complexes,  
 475–483  
 Fourier transform infrared studies  
 on Langmuir-Blodgett film of  
 amphiphile with Schiff base as  
 headgroup and its copper  
 complex, 301–321  
 of structural phase transition in  
 $[N(CH_3)_4]_3Bi_2Br_9$ , 269–282  
 Fourier transform Raman spectroscopy  
 for quantification of the number of  
 silanol groups in silicalite and  
 mesoporous MCM-41,  
 569–584  
 of synthetic solid solutions formed  
 by orthorhombic sulfates,  
 323–336  
 Frequency domain, power spectral  
 estimation for signal  
 processing in, 369–383

## G

- Gamma irradiated  $C_2H_2O_4 \cdot 2H_2O$   
single crystal, electron spin  
resonance of, 735–742
- 3-Geranyl-4-hydroxybenzoate deriva-  
tive from *Xerophyta plicata*,  
isolation and determination of  
xerophytolic acid as, 643–651
- Gradient selected constant time  
COSY, 1–8

## H

- Heavy metals, determination in  
lichens growing on ecological  
habitats in Schirmacher Oasis,  
East Antarctica, 435–444
- Heptakis-(2,3,6-tri-*o*-methyl)- $\beta$ -  
cyclodextrin, conformational  
analysis by two-dimensional  $^1H$   
NMR of inclusion complex  
with 3-hydroxy-2-naphthalene  
carboxylate anion, 407–413
- High pressures, Raman spectra of  
selenium dioxide at, 555–567
- 3-Hydroxy-2-naphthalenecarboxylate  
anion, conformational analysis  
by two-dimensional  $^1H$  NMR  
of inclusion complex with  
heptakis-(2,3,6-tri-*o*-methyl)- $\beta$ -  
cyclodextrin, 407–413

## I

- Inclusion complex, of 3-hydroxy-2-  
naphthalenecarboxylate anion  
with heptakis-(2,3,6-tri-*o*-meth-  
yl)- $\beta$ -cyclodextrin, conforma-  
tional analysis by two-dimen-  
sional  $^1H$  NMR, 407–413

- Inductively coupled plasma atomic  
emission spectrometry  
for determination of cadmium  
separated selectively with  
ion exchange method,  
195–200
- for line selection and determina-  
tion of trace amounts of  
elements in high-purity iron,  
127–145
- using power spectral estimation for  
signal processing in the  
frequency domain, 369–383
- Infrared spectra  
of pendant double bonds in mono-  
disperse micron macroporous  
resin, 867–874
- polarized, of 2-(methylthio)benzo-  
thiazole, 535–545
- vapor phase, of analogs of the  
nerve agent Sarin, 795–819
- Infrared spectroscopy, of complexes  
of Co(II), Ni(II), Cu(II), Zn(II),  
and Cd(II) with 1-ethyl-1,2,4-  
triazole and 1-propyl-1,2,4-  
triazole, 23–34
- Insect adipokinetic hormones, NMR  
of, 875–891
- Integerrimine, molecular con-  
formational analysis of,  
495–507
- Intensity behavior  
of chloro and fluoro derivatives of  
the benzene chromophore in  
the near-ultraviolet spectra,  
681–692
- of long wavelength UV transition  
of the pyridine chromophore  
under substitution in the para  
position to the nitrogen atom,  
615–623

- Interaction vector model, for study of the intensity behavior in the near-ultraviolet spectra of chloro and fluoro derivatives of the benzene chromophore, 681–692
- Ion exchange method, inductively coupled plasma atomic emission spectrometry determination of cadmium separated selectively with, 195–200
- Ion guide source, optogalvanic properties in the recoil nucleus selective photoionization, 83–90
- Iron, high purity, inductively coupled plasma atomic emission spectrometry for line selection and determination of trace amounts of elements in, 127–145
- <sup>57</sup>Iron-Mössbauer studies, on loess-palaeosol samples from central China, 833–846

## L

- Langmuir-Blodgett films of amphiphile with Schiff base as headgroup and its copper complex, Fourier transform infrared studies of, 301–321
- mixed chiral amino acid porphyrin, absorption spectral study of, 633–642
- Lanolin, influence on the determination of vitamin A in fats, 227–234

- Lichens, growing on different ecological habitats in Schirmacher Oasis, East Antarctica, determination of heavy metals in, 435–444
- Ligand field theory, of a copper complex, 185–194, 971
- Loess, Chinese, electron paramagnetic resonance determination of manganese<sup>2+</sup> in, 201–209
- Loess-palaeosol samples, from central China, <sup>57</sup>iron-Mössbauer studies on, 833–846
- Long wavelength UV transition, of the pyridine chromophore under substitution in the para position to the nitrogen atom, intensity behavior of, 615–623
- Luminescence spectra, of alkali feldspars, 103–113
- Luminescence study, high resolution, of salts of triethylenetetra-aminehexaacetatouropate(III) anion, 585–594

## M

- Macroporous resin, monodisperse micron, infrared spectra of pendent double bonds in, 867–874
- Magnetic resonance spectroscopy, continuous wavelet transform as a method for postacquisition water suppression in, 47–67
- Manganese<sup>2+</sup>, electron paramagnetic resonance in the Chinese loess samples, 201–209
- Mass spectrometry, of oligosaccharides, 469–473
- See also* Electrospray mass spectrometry

- MCM-41, mesoporous, Fourier transform Raman spectroscopy quantification of the number of silanol groups in, 569–584
- 2-Mercaptobenzothiazole, Fourier transform SERS study of pH dependence of, 901–907
- Metal complexes, of 6-(2-pyridylazo)-3-acetamidophenol, preparation, chemical characterization, and electronic spectra of, 821–832
- Methoxamine HCl, spectrophotometric and fluorimetric determination in bulk drugs and in ampoules, 9–21
- $\beta$ -Methyl- $\beta$ -nitrostyrene, role of triethylamine in the reversible photoisomerization of, 385–391
- 2-(Methylthio)benzothiazole, polarized infrared and Raman spectra of, 535–545
- Micelle aqueous solutions, CTAB, electric spectra of tetra-hydro-phenyl porphyrin with two hexadecyl chains in, 157–172
- Molecular conformational analysis, of integerrimine, 495–507
- Mulliken two-state model, charge-transfer complex absorption spectra and applicability of, 595–614
- Multicomponent analgetic mixture, the second-derivative spectrophotometric assay of, 173–183

## N

- Naphthalenes, substituted, statistical substituent chemical shift values for prediction of the  $^{13}\text{C}$  NMR signal positions in, 255–267
- 1-Naphtyl-polymethoxylated diphenylmethanes,  $^1\text{H}$  and  $^{13}\text{C}$  NMR determination of, 211–226
- $[\text{N}(\text{CH}_3)_4]_3\text{Bi}_2\text{Br}_9$ , Fourier transform infrared studies of structural phase transition in, 269–282
- Near-ultraviolet spectra, intensity behavior of chloro and fluoro derivatives of the benzene chromophore in, 681–692
- Nerve agent Sarin, vapor-phase infrared spectral study of analogs of, 795–819
- Nickel(II) metal ions, vibrational spectra of complexes with *o*-hydroxybenzylidene *o*-hydroxyaniline, 523–533
- $\beta$ -Nitrostyrenes,  $^{13}\text{C}$  NMR spectroscopy of, 337–345
- NMR. *See* Nuclear magnetic resonance
- $\text{NO}_2$ , rotational analysis of the vibronic bands in the range 5050–5200  $\text{\AA}$ , 743–753
- Nuclear magnetic resonance (NMR) for analysis of a complex spin system from a *spiro*-chamigrene, 457–467
- bromine-81, for probe of anion binding to porcine pancreatic  $\alpha$ -amylase, 893–899
- $^{13}\text{C}$ , of  $\beta$ -nitrostyrenes, 337–345
- $^{13}\text{C}$  signal positions, statistical substituent chemical shift values for prediction in substituted naphthalenes, 255–267
- for characterization of  $\text{Pd}_2(\text{PPh}_3)_2\text{Cl}_2\text{O}$ , 671–680
- comparative study of water  $T_1$  and  $T_2$  relaxation times by, 115–125



[Nuclear magnetic resonance (NMR)  
*continued*]

- $^1\text{H}$ ,  
for quantitative analysis of  
acetazolamide in solid dosage  
forms, 661–669  
relaxation time results,  
compared with viscosity  
and density versus temperature  
for liquid derivatives of  
benzene and pyridine,  
917–930
- $^1\text{H}$  and  $^{13}\text{C}$   
for determination of the  
mechanism of formation of  
polysubstituted diphenyl-  
methane dimers by reduction of  
polymethoxylated benzo-  
phenones, 755–775  
for determination of 1-naphthyl-  
polymethoxylated diphenyl-  
methanes, 211–226
- $^1\text{H}$  and  $^{13}\text{C}$  spectra, of an anti-  
bacterial bisabolane ses-  
quiterpene by two-  
dimensional NMR  
techniques, 235–243
- $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$ , spectral  
parameters and geometries of  
isomeric diphenylbicy-  
clo[2.2.1]hept-5-ene-2,3-  
dicarboxylic anhydrides,  
717–734
- of insect adipokinetic hormones,  
875–891
- two dimensional  $^1\text{H}$ , for con-  
formational analysis of  
3-hydroxy-2-naphthalene  
carboxylate anion with  
heptakis-(2,3,6-tri-*o*-methyl)-  
 $\beta$ -cyclodextrin, 407–413

## O

- o*-Hydroxybenzylidene *o*-hydroxy-  
aniline, vibrational spectra  
of complexes with copper(II)  
and nickel(II) metal ions,  
523–533
- Oligobipyridine ligand, electrospray  
mass spectrometry of com-  
plexation with transitional  
metal ions, 35–46
- Oligosaccharide  $[\text{M} + \text{MNH}_4]^+$  ion,  
formation in fast atom  
bombardment mass spec-  
trometry, 469–473
- Optogalvanic properties, of the ion  
guide source in the recoil  
nucleus selective photo-  
ionization, 83–90
- Orthorhombic sulfates, Fourier  
transform Raman study of  
synthetic solid solutions  
formed by, 323–336
- Oxazolidinone derivative, structural  
study of *N*-(1-benzoyl-3-  
pyrrolidinyl) benzamide,  
777–793
- Oxine, photoacoustic phase spectra of  
rare earth complexes with,  
547–554

## P

- $\text{Pd}_2(\text{PPh}_3)_3\text{Cl}_3\text{O}$ , spectroscopic  
characterization and X-ray  
crystal structure of, 671–680
- Pendent double bonds, in mono-  
disperse micron macroporous  
resin, infrared spectra of,  
867–874

- Penicillins, use of cerium(IV) in the spectrophotometric and spectrofluorimetric determination in pharmaceutical preparations of, 931–948
- pH dependence, of 2-mercaptobenzo-thiazole, Fourier transform SERS of, 901–907
- Phase transition, structural, Fourier transform infrared studies in  $[\text{N}(\text{CH}_3)_4]_3\text{Bi}_2\text{Br}_9$ , 269–282
- Phenols, di- and trihydric, spectrophotometric determination by EDTA titration, 347–357
- Phospholipid, of human erythrocyte membrane, effects of *N*-phosphoproline on, 653–659
- N*-Phosphoproline, effects on the phospholipid of human erythrocyte membrane, 653–659
- Photoacoustic phase spectra, of rare earth complexes with oxine, 547–554
- Photoacoustic spectroscopy, of a pair of hydrate isomers for dinuclear copper(II) complexes, 847–856
- Photoisomerization, reversible, of  $\beta$ -methyl- $\beta$ -nitrostyrene, 385–391
- Photophysical properties, of rare earth ions encapsulated water soluble functional calix-resorcin[4]arene, 705–715
- Phthalocyanine complexes, mono and bis, SERS spectra deposited on silver and gold supports, 625–631
- Platyphylline, crystal structure of, 495–507
- Polymorphism, of 4-fluorophenyl-pyruvic acid, X-ray crystallography and vibrational spectroscopy of, 509–522
- Powder phosphors,  $(\text{Ln}, \text{Zn})\text{OF}:\text{Eu}^{3+}$ , effect of zinc ions on the fluorescence spectra of, 423–434
- Power spectral estimation, for signal processing in the frequency domain, 369–383
- Precious metals  
  recycle procedure for laboratory rejects, 485–494  
  spectroscopic characterization and X-ray crystal structure of  $\text{Pd}_2(\text{PPh}_3)_3\text{Cl}_2\text{O}$ , 671–680
- 1-Propyl-1,2,4-triazole, spectral and magnetic studies of complexes with Co(II), Ni(II), Cu(II), Zn(II), and Cd(II), 23–34
- Pyridine chromophore long wavelength UV transition, intensity behavior under substitution in the para position to the nitrogen atom, 615–623
- Pyridine liquid derivatives, viscosity and density versus temperature compared with  $^1\text{H}$  NMR relaxation time results for, 917–930
- 6-(2-Pyridylazo)-3-acetamidophenol and its metal complexes, preparation, chemical characterization, and electronic spectra of, 821–832
- R
- RA-IR studies, of the structure of mercapto-ended azobenzene derivatives in self-assembled monolayers, 949–962

## Raman spectra

- of cobalt(II) cupferronato derivatives, 857–865
- polarized, of 2-(methylthio)benzothiazole, 535–545
- of selenium dioxide at high pressures, 555–567
- of 4-vinylpyridine and its transition metal(II) tetracyanonickelate complexes, 475–483

Ramipril, spectrophotometric determination in dosage forms, 91–102

## Rare earth

- ions encapsulated water-soluble functional calixresorcin[4]-arene, photophysical properties of, 705–715
- photoacoustic phase spectra of complexes with oxine, 547–554

Recoil nucleus selective photoionization, optogalvanic properties of the ion guide source in, 83–90

Recycling, of precious metals, 485–494

Rotational analysis, of the vibronic bands of NO<sub>2</sub> in the range 5050–5200 Å, 743–753

Ruthenium(II) complexes, spectral characterization of, 415–421

## S

Sarin, vapor-phase infrared spectral study of analogs of, 795–819

## Schiff base

- bidentate ligands, comparison of the ultraviolet-visible spectra of the copper(II) complexes of, 445–455

- as headgroup in amphiphile, Fourier transform infrared studies of copper complex with, 301–321

Second-derivative spectrophotometry, of a multicomponent analgetic mixture, 173–183

Selenium dioxide, Raman spectra at high pressures, 555–567

Self-assembled monolayers, RA-IR studies of the structure of mercapto-ended azobenzene derivatives in, 949–962

SERS. *See* Surface-enhanced Raman Spectroscopy

Silanol, Fourier transform Raman spectroscopy quantification of the number in silicalite and mesoporous MCM-41, 569–584

Silicalite, Fourier transform Raman spectroscopy quantification of the number of silanol groups in, 569–584

Solid dosage forms, <sup>1</sup>H NMR spectroscopy for quantitative analysis of acetazolamide in, 661–669

Spectral dependencies, of the surface photo-charge effect at conducting surfaces, 393–398

Spectral techniques, for study of barium(II) complexes, 909–916

Spectrofluorimetry, for determination of penicillins and cephalosporins in their pharmaceutical preparations, use of cerium(IV) in, 931–948

## Spectrophotometry

- for determination of di- and trihydric phenols by EDTA titration, 347–357

- for determination of enalapril maleate and ramipril, 91–102
- for determination of methoxamine HCl in bulk drugs and in ampoules, 9–21
- for determination of penicillins and cephalosporins in their pharmaceutical preparations, use of cerium(IV) in, 931–948
- second derivative, for assay of a multicomponent analgetic mixture, 173–183
- Spiro*-chamigrene, NMR analysis of a complex spin system from, 457–467
- Statistical substituent chemical shift values, for prediction of the  $^{13}\text{C}$  NMR signal positions in substituted naphthalenes, 255–267
- Surface photo-charge effect, spectral dependencies at conducting surfaces, 393–398
- Surface-enhanced Raman spectroscopy (SERS)
  - Fourier transform, for study of pH dependence of 2-mercapto-benzothiazole, 901–907
  - spectra, of mono and bisphthalocyanine complexes deposited on silver and gold supports, 625–631
  - for study of adsorption states, 69–81

## T

- Temperature, and sequence and hot transitions in the carbon dioxide molecule, interference of adjacent lines belonging to different transitions as a function of, 283–300

- Tetra-hydrophenyl porphyrin, with two hexadecyl chains, electric spectra study in CTAB micelle aqueous solutions, 157–172
- Transition metal cupferronato complexes, vibrational behavior of, 857–865
- Transition metal(II) tetracyanonickelate complexes, of 4-vinylpyridine, Fourier transform infrared and Raman spectra of, 475–483
- Transitional metal ions, electrospray mass spectrometry of complexation with oligobipyridine ligand, 35–46
- Triethylamine, role in the reversible photoisomerization of  $\beta$ -methyl- $\beta$ -nitrostyrene, 385–391
- Triethylenetetraaminehexaacetato-uroate(III) anion, high-resolution luminescence study of salts of, 585–594

## U

- Ultraviolet emission band, influence of crushing on, 103–113
- Ultraviolet-visible spectra, of the copper(II) complexes of the bidentate Schiff base ligands, 445–455

## V

- Vibrational behavior, of transition metal cupferronato complexes, 857–865

- Vibrational spectra, of benzylidene aniline, *o*-hydroxybenzylidene *o*-hydroxyaniline and its complexes with copper(II) and nickel(II) metal ions, 523–533
- Vibrational spectroscopy, of the polymorphism of 4-fluorophenylpyruvic acid, 509–522
- 4-Vinylpyridine and its transition metal(II) tetracyanonickelate complexes, Fourier transform infrared and Raman spectra of, 475–483
- Vitamin A, influence of lanolin and white soft paraffin on determination in fats, 227–234

## W

- Water suppression, postacquisition, continuous wavelet transform as a method for in biomedical proton magnetic resonance spectroscopy, 47–67
- Water  $T_1$  and  $T_2$  relaxation times, NMR comparative study in healthy and pathological blood, 115–125

- Wavelet transform, continuous, as a potential method for post-acquisition water suppression in biomedical proton magnetic resonance spectroscopy, 47–67
- White soft paraffin, influence on the determination of vitamin A in fats, 227–234

## X

- Xerophyta plicata*, isolation and structural determination of xerophytolic acid as a 3-geranyl-4-hydroxybenzoate derivative from, 643–651
- Xerophytolic acid, isolation and structural determination of, 643–651
- X-ray crystallography, of the polymorphism of 4-fluorophenylpyruvic acid, 509–522

## Z

- Zinc ions, effect on the fluorescence spectra of (Ln, Zn)OF:Eu<sup>3+</sup> powder phosphors, 423–434