

This article was downloaded by:

On: 30 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Spectroscopy Letters

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597299>

Subject Index to Volume 36

Online publication date: 16 December 2003

To cite this Article (2003) 'Subject Index to Volume 36', *Spectroscopy Letters*, 36: 5, 577 — 584

To link to this Article: DOI: 10.1081/SL-120026618

URL: <http://dx.doi.org/10.1081/SL-120026618>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Subject Index to Volume 36

A

N-Acetyl-L-alanine and N-acetyl-L-alaninate, conformation and vibrational frequencies of, 537–550
Acetylbenzene, chiroptical studies of lanthanide(III) complexes of sulphur–nitrogen–oxygen ligands derived from, 239–250
Aerosol supply, power interruption technique for determining the difference between electron and gas temperatures in the argon d.c. arc supplied with, 303–312
Aluminum, trace analysis on silicon surfaces by ultra-soft X-ray emission spectroscopy, 93–98
1,2-bis(*p*-Aminophenoxy)ethane, transition metal complexes with a Schiff based derived from, 429–440

Ammonium, diffuse reflectance infrared spectroscopy quantitative analysis of addition to silica sand and soil, 251–274

Argon radio-frequency inductively coupled plasma, excitation mechanism of chromium emission lines in, 99–115

Aromatic protons, concentration-dependent variation of ^1H -NMR chemical shifts in sampangine derivatives, 477–485

Atomic fluorescence spectrometry, L-cysteine enhanced hydride generation for determination of germanium in geological samples by, 275–285

B

Boron oxide, vibrational analysis of the β system of, 531–536



Bovine serum albumin, mechanism of interaction with bromophenol blue, 197–209
 Bromophenol blue, mechanism of interaction with bovine serum albumin, 197–209

C

Calcium oxalates, quantitative FTIR spectroscopic evaluation in deteriorated mural paintings, 501–513
 Calix[4]arene-crown-6-cation binary complexes, comparison of stability under electrospray mass spectrometry, 327–340
 CARS. *See* Coherent anti-Stokes Raman scattering
 Charge transfer, protonation dependent photoinduced intramolecular, in 2-(*p*-dimethylaminostyryl)benzazoles, 375–386
 Charge transfer molecular complexes, of pyrazoles with σ - and π -electron acceptors, 357–373
 Chemiluminescence flow injection, for assay of fluoride in waters and human urine using immobilized reagents, 117–131
 Chiroptical studies, of lanthanide(III) complexes of sulphur–nitrogen–oxygen ligands derived from L-cysteine and acetylbenzene, 239–250
 Chlorsulfuron, infrared method with reduced solvent con-

Subject Index to Volume 36

sumption for determination in pesticide formulations, 515–529
 Chromium emission lines, excitation mechanism in the argon radio-frequency inductively coupled plasma, nitrogen microwave induced plasma, and argon or nitrogen glow discharge plasmas, 99–115
 Cobalt(III), in symmetric complexes with macrocyclic and β -diketonato ligands, 183–196
 Coherent anti-Stokes Raman scattering (CARS), nanosecond time resolved resonance, of triplet excited polycyclic aromatic compounds, 1–23
 Complexes
 calix[4]arene-crown-6-cation binary, comparison of stability under electrospray mass spectrometry, 327–340
 of cobalt(III) with macrocyclic and β -diketonato ligands, 183–196
 group VI dinuclear oxo metal, of salicylideneimine-2-anisole Schiff base, 71–82
 lanthanide(III), of sulphur–nitrogen–oxygen ligands derived from L-cysteine and acetylbenzene, 239–250
 obtained by mechanochemical reactions of hemin, 83–92
 of oxime ligands, spectroscopic characterization of, 51–70
 transition metal with a Schiff based derived from 4-methoxybenzaldehyde and 1,2-bis(*p*-aminophenoxy)-ethane, 429–440



Concentration-dependent variation, of $^1\text{H-NMR}$ chemical shifts for aromatic protons in sampaagine derivatives, 477–485

Copper electrodes, surface-enhanced Raman spectroscopy of adsorption of cysteamine at, 211–226

Cysteamine, surface-enhanced Raman spectroscopy of adsorption at copper electrodes, 211–226

L-Cysteine

atomic fluorescence spectrometric determination of germanium in geological samples by hydride generation enhanced by, 275–285

chiroptical studies of lanthanide(III) complexes of sulphur–nitrogen–oxygen ligands derived from, 239–250

D

Dental tissue, thermally treated, electron spin resonance–native signal in, 487–499

Diatomite/polyaniline composite, spectral studies of, 25–33

Diffuse reflectance infrared spectroscopy, for quantitative analysis of added ammonium and nitrate in silica sand and soil, 251–274

β -Diketonato ligands, spectral assignments of symmetric cobalt(III) complexes with, 183–196

2-(p-Dimethylaminostyryl)benzazoles, protonation depen-

dent photoinduced intramolecular charge transfer in, 375–386

Diode laser-excited optogalvanic and absorption measurements of uranium in a hollow cathode discharge, 167–180

Doubly charged ion emission lines, excitation from a Grimm-style glow discharge plasma, 561–572

E

Electron acceptors, σ and π , charge transfer molecular complexes of pyrazoles with, 357–373

Electron spin resonance–native signal, in thermally treated dental tissue, 487–499

Electrospray mass spectrometry, comparison of stability of calix[4]arene-crown-6-cation binary complexes under, 327–340

Excitation mechanism, of chromium emission lines in the argon radio-frequency inductively coupled plasma, nitrogen microwave induced plasma, and argon or nitrogen glow discharge plasmas, 99–115

F

Films, poly(2-methyl-1-naphthylamine), electrochemical studies and structure of, 151–165



Fluorescence, native, of pyridoxine hydrochloride in pharmaceutical preparations after sorption on Sephadex SP C-25, 133–149

Fluoride, chemiluminescence flow injection assay in waters and human urine using immobilized reagents, 117–131

Fluorometric characteristics, of a wave-guide cell with low refractive index, 551–560

Fourier transform infrared (FTIR) spectroscopy, quantitative, for evaluation of gypsum and calcium oxalates in deteriorated mural paintings, 501–513

FTIR. *See* Fourier transform infrared

Furocoumarinones, isomeric, gas chromatographic ion trap tandem mass spectrometric characterization of, 387–401

2-Furylbenzothiazoles and their vinyllogues, substituted, absorption and fluorescence properties of, 43–50

G

Gas chromatography ion trap tandem mass spectrometry, for characterization of isomeric furocoumarinones, 387–401

Germanium, determination in geological samples by atomic fluorescence spectrometry, 275–285

Subject Index to Volume 36

Glow discharge plasmas
argon or nitrogen, excitation mechanism of chromium emission lines in, 99–115

Grimm style, excitation of doubly charged ion emission lines from, 561–572

1-(β -D-Glucopyranosyl)-1,2,4-triazoles, structural assignments by ^1H - and ^{13}C -NMR, 461–475

Graphite furnace, behavior of volatile elements in the presence of silver as matrix modifier, 441–447

Gypsum, quantitative FTIR spectroscopic evaluation in deteriorated mural paintings, 501–513

H

Hemin, spectroscopic characterization of complexes obtained by mechanochemical reactions of, 83–92

Heroin diluent, identification of, 35–42

Hollow cathode discharge, diode laser-excited optogalvanic and absorption measurements of uranium in, 167–180

Hydrocarbons, higher, methane photo-conversion using 355 nm laser radiation, 313–326

Hydrochlorothiazide, in tablets with spironolactone, quantitative two-component analysis by one-dimensional continuous Daubechies and biorthogo-

nal wavelet analysis of ultraviolet spectra, 341–355
 N-(2-Hydroxy-1-naphthylidene)-anils, keto-enol tautomerism in, 287–301
 Hydroxytrifluoroborate anion, quantum chemical calculations and vibrational spectra of, 227–237

I

Infrared method, with reduced solvent consumption, for the determination of chlorsulfuron in pesticide formulations, 515–529

K

Keto-enol tautomerism, in N-(2-hydroxy-1-naphthylidene)-anils, 287–301

L

Lanthanide(III) complexes, of sulphur–nitrogen–oxygen ligands derived from L-cysteine and acetylbenzene, 239–250

Leukotriene D4 receptor antagonists, stereochemical elucidation of norbornene derivatives synthesized as, 407–418

M

Macrocyclic ligands, spectral assignments of symmetric cobalt(III) complexes with, 183–196

Mechanochemical reactions, of hemin, 83–92

Metal complexes, group VI dinuclear oxo, of salicylidene-imine-2-anisole Schiff base, 71–82

Methane, photo-conversion into higher hydrocarbons using 355 nm laser radiation, 313–326

4-Methoxybenzaldehyde, transition metal complexes with a Schiff base derived from, 429–440

Myrrh, colorimetric determination of crude powdered, purified extract, oily fraction, and pharmaceutical dosage forms (erratum), 181

N

Nitrate, diffuse reflectance infrared spectroscopy quantitative analysis of addition to silica sand and soil, 251–274

Nitrogen microwave induced plasma, excitation mechanism of chromium emission lines in, 99–115

NMR. *See* Nuclear magnetic resonance

Norbornene derivatives, synthesized as leukotriene D4 receptor antagonists, stereochemical elucidation of, 407–418

Nuclear magnetic resonance (NMR)
 ^{13}C , of the conformations of serine-3'- and 5'-thymidine monophosphate conjugates, 419–427



¹H and ¹³C, for structural assignments of 1-(β -D-glucopyranosyl)-1,2,4-triazoles, 461–475

¹H chemical shifts, concentration-dependent variation for aromatic protons in sampanine derivatives, 477–485
one and two dimensional proton and carbon-13, of procaine hydrochloride, 35–42

O

Oxime ligands and their complexes, spectroscopic characterization of, 51–70

P

Pesticide formulations, infrared method with reduced solvent consumption for the determination of chlorsulfuron in, 515–529

Photo-conversion, of methane into higher hydrocarbons using 355 nm laser radiation, 313–326

Plasma diagnostics, power interruption technique for determining the difference between electron and gas temperatures in the argon d.c. arc supplied with aqueous aerosol, 303–312

Polyaniline/diatomite composite, spectral studies of, 25–33

Polycyclic aromatic compounds, triplet excited, nanosecond time-resolved resonance CARS of, 1–23

Subject Index to Volume 36

Poly(2-methyl-1-naphthylamine) films, electrochemical studies and structure of, 151–165

Power interruption technique, for determining the difference between electron and gas temperatures in the argon d.c. arc supplied with aqueous aerosol, 303–312

Procaine hydrochloride, one- and two-dimensional proton and carbon-13 NMR of, 35–42

Product operator theory, for spin-5/2 nuclei, 449–459

Protonation dependent photoinduced intramolecular charge transfer, in 2-(*p*-dimethylaminostyryl)benzazoles, 375–386

Pyrazoles, in charge transfer molecular complexes with σ - and π -electron acceptors, 357–373

Pyridoxine hydrochloride (vitamin B₆), native fluorescence in pharmaceutical preparations after sorption on Sephadex SP C-25, 133–149

Q

Quantitative two-component analysis, of a mixture containing hydrochlorothiazide and spironolactone in tablets by one-dimensional continuous Daubechies and biorthogonal wavelet analysis of ultraviolet spectra, 341–355

Quantum chemical calculations, of the hydroxytrifluoroborate anion, 227–237



S

Salicylideneimine-2-anisole Schiff base, group VI dinuclear oxo metal complexes of, 71–82

Sampangine derivatives, concentration-dependent variation of ^1H -NMR chemical shifts for aromatic protons in, 477–485

Schiff base
derived from 4-methoxybenzaldehyde and 1,2-bis(*p*-aminophenoxy)ethane, spectroscopic and conductance studies of transition metal complexes with, 429–440
salicylideneimine-2-anisole, group VI dinuclear oxo metal complexes of, 71–82

Serine-3'- and 5'-thymidine mono-phosphate conjugates, ^{13}C NMR studies of the conformations of, 419–427

Silicon surfaces, ultra-soft X-ray emission spectroscopy trace analysis of aluminum on, 93–98

Silver, as matrix modifier, behavior of volatile elements in the graphite furnace in the presence of, 441–447

Spin-5/2 nuclei, product operator theory for, 449–459

Spironolactone, in tablets with hydrochlorothiazide, quantitative two-component analysis by one-dimensional continuous Daubechies and bi-orthogonal wavelet analysis of ultraviolet spectra, 341–355

Stability, comparison for calix[4]-arene-crown-6-cation binary complexes under electrospray mass spectrometry, 327–340

Stereochemical elucidation, of norbornene derivatives synthesized as leukotriene D4 receptor antagonists, 407–418

Surface-enhanced Raman spectroscopy, of adsorption of cysteamine at copper electrodes, 211–226

T

Thermally treated dental tissue, electron spin resonance–native signal in, 487–499

Trace analysis, of aluminum on silicon surfaces by ultra-soft X-ray emission spectroscopy, 93–98

Transition metal complexes, with a Schiff base derived from 4-methoxybenzaldehyde and 1,2-bis(*p*-aminophenoxy)ethane, 429–440

U

Uranium, diode laser-excited optogalvanic and absorption measurements in a hollow cathode discharge, 167–180

V

Vibrational spectra of the β system of $^{10}\text{B}^{18}\text{O}$, 531–536



of the hydroxytrifluoroborate anion, 227–237

Vitamin B₆. *See* Pyridoxine hydrochloride

Volatile elements, behavior in the graphite furnace in the presence of silver as matrix modifier, 441–447

W

Wave-guide cell, with low refractive index, fluorometric characteristics of, 551–560

Wavelet analysis of ultraviolet spectra, one dimensional continuous Daubechies and bi-orthogonal, for quantitative two-component analysis of a mixture containing hydrochlorothiazide and spironolactone in tablets, 341–355

X

X-ray emission spectroscopy, ultra-soft, for trace analysis of aluminum on silicon surfaces, 93–98

