



Mercury compounds characterization by thermal desorption

M. Rumayor, M. Diaz-Somoano*, M.A. Lopez-Anton, M.R. Martinez-Tarazona

Instituto Nacional del Carbón (CSIC), C/Francisco Pintado Fe No. 26, 33011 Oviedo, Spain

ARTICLE INFO

Article history:

Received 25 February 2013

Received in revised form

20 May 2013

Accepted 27 May 2013

Available online 31 May 2013

Keywords:

Mercury speciation

Solid samples

Thermal desorption

ABSTRACT

The ability to accurately determine metal mercury content and identify different mercury species in solid samples is essential for developing remediation and control strategies. The aim of the present study is to characterize mercury compounds based on thermal desorption. For this purpose a series of samples was prepared and the operational parameters—heating velocity, carrier gas—were optimized. Fifteen commercial mercury compounds were analyzed for use as fingerprints. The results of the study show that the identification of mercury species by the method of thermal desorption is possible. The temperature of desorption increased according to the following order $HgI_2 < HgBr_2 < Hg_2Cl_2 = HgCl_2 < Hg(CN)_2 < Hg-Cl_2O_8 \cdot H_2O < Hg(SCN)_2 < HgS$ (red) $< HgF_2 < Hg_2(NO_3)_2 \cdot 2H_2O < Hg(NO_3)_2 \cdot H_2O < HgO$ (yellow, red) $< Hg_2-SO_4 < HgSO_4$. A comparison of the estimated total mercury content with the mercury content calculated by integrating the area of the desorption curve shows that recoveries of 79–104% for HgS can be estimated. The proposed method represents a significant step forward in direct mercury analysis in solid samples.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Mercury (Hg) species are potentially toxic compounds, having special concern for human health and the environment. Mercury enters the environment after being released from natural and anthropogenic sources [1]. Because Hg cycles, it can be carried over long distances far away from the source of its emission. It is then deposited in aquatic environments through precipitation and is generally found in organic form as methylmercury.

Mercury is rare in its native state. The main mercury ores include cinnabar and calomel, though there are a few other secondary minerals (corderoite, livingstonite, etc.). Rocks, sediments, water, and soil, all contain small amounts of mercury, which may be released into the environment due to exposure to wind, water, and volcanic activity. However human activity has increased the mobilization and transformations of mercury into the environment [2]. Until now most attention has been directed towards the total amount of mercury. However the behavior of Hg when it is emitted from different sources and its final fate and distribution in byproducts depend to a large extent on the form in which it occurs [3]. Each mercury species in a solid matrix interact in different ways, exhibiting a different degree of solubility and mobility. Moreover the development of mercury remediation and control technologies [4] must take into account mercury associations and the interaction of mercury with solids.

Sequential extraction procedures based on the different solubilities of Hg compounds are commonly used for Hg speciation studies

[5–9]. These methods normally consist of several steps with one or more reagents, making the procedure both tedious and time consuming. Sample contamination or losses of mercury through volatilization are problems generally associated with these methods [10,11]. Other limitations are the poor reproducibility and selectivity of the results. The advantages of these methods, however, are that they are quite easy to carry out in a standard laboratory and they are relatively cheap. As alternatives, instrumental methods such as X-ray adsorption fine structure spectroscopy (EXAFS) [12,13] and X-ray absorption near edge structure (XANES) [14], have been applied for mercury speciation in sediments and soils, but, these techniques are expensive and the detection limits (100 mg kg^{-1}) are not usually appropriate for the analysis of the majority of solid samples. Additionally, for decades, thermal desorption techniques have been applied for mercury speciation to a variety of solid samples such as sediments [15–21], iron-based sorbents for mercury removal [22], fluorescent lamp wastes [23], airborne particulate matter [24] and combustion power station by-products [25–27]. However reproducible results have still not been obtained, in part due to the lack of reference materials for validation purposes. To fill this gap, this study aims to characterize mercury compounds by their behavior during thermal desorption. Although thermo-desorption is not a new technique the proposed application represents a significant step forward for mercury direct analysis in solid samples.

2. Materials and methods

For this study an advanced RA-915 Mercury analyzer coupled to a Pyro-915 furnace, both from Lumex, were used. This mercury

* Corresponding author. Tel.: +34 985119090; fax: +34 985297662.

E-mail address: mercedes@incar.csic.es (M. Diaz-Somoano).

analyzer is a highly versatile equipment that can be used to determine Hg contents in air, liquids and solids both in laboratory and field conditions. The operation of an RA-915 analyzer is based on differential Zeeman atomic absorption spectrometry using high frequency modulation of light polarization. The PYRO-915+ furnace consists of two chambers in series. The first one serves to evaporate liquids and pyrolyze the solid samples. In this section the mercury compounds are released from the solid matrix under a controlled heating mode. The second chamber is kept at approximately 800 °C and is used to reduce the mercury compounds to elemental mercury and to eliminate smoke and interference compounds. The temperature of the first chamber is continuously monitored by means of a thermocouple. Calibration coefficient is set using standard mercury-containing sample. A feature of the equipment is the high calibration stability.

For the purpose of the present study the heating mode was optimized until a good signal resolution was obtained. As can be seen in Table 1 a four-step program was set up. The temperature rate was kept at 40 °C min⁻¹ for 575 s. Then the heating velocity was increased up to 50 °C min⁻¹ where it was held for 200s and then up to 80 °C min⁻¹ where it was held for a further 125s. Air was used as carrier gas at 1 L min⁻¹.

In the first step commercially pure mercury compounds were analyzed in order to determine their desorption temperature and profile. Because of the high Hg content the commercially pure compounds were previously diluted by blending them with an inert material. For this purpose 10 mg of each pure Hg compound was crushed and mixed with silica to obtain the fifteen standard homogenized samples with Hg concentrations of about 800 mg kg⁻¹. Each sample was analyzed three times. The accuracy and precision of the analysis was determined by means of several analysis of a standard sample. HgS was selected for this evaluation. The Hg species were characterized by the temperature range in which they were released. The compounds tested were (i) the most common mercury compounds found in coal and geological samples: HgCl₂, HgS, HgSO₄, HgO, Hg₂Cl₂, Hg₂SO₄, HgBr₂ and (ii) commercially available compounds which are used as pigments or for other industrial applications: HgI₂, Hg(CN)₂, HgCl₂O₈ · H₂O, Hg(SCN)₂, HgF₂, Hg₂(NO₃)₂ · H₂O, Hg(NO₃)₂ · 2H₂O. These compounds were included in the study for use as fingerprints.

In order to detect any potential interferences or interactions resulting from the thermal release of different Hg species, several mixtures of these Hg compounds were analyzed. Due to the lack of standard reference materials for the analysis, the mixtures of Hg species were prepared in the laboratory. Mixtures of pure compounds were prepared by crushing 10 mg of each compound with silica to obtain blends with mercury concentrations of about 1000 mg kg⁻¹. It should be pointed that relatively high mercury concentrations are considered in this work to minimize heterogeneity problems. However the total mercury content of the analyzed mixtures was kept below 10 µg to prevent the equipment from being contaminated. To procure the thermograms, approximately 5 mg of prepared mixture were used, depending on the Hg content. Quantification was carried out by peak integration using Origin 6.0 software.

Table 1
Heating program optimized for the analysis.

| Time (s) | Heating rate (°C/min) |
|----------|-----------------------|
| 0–575 | 40 |
| 576–775 | 50 |
| 776–900 | 80 |

3. Results and discussion

3.1. Mercury species identification: pure compounds

Each Hg species was characterized by its thermo-desorption profile. The desorption profile shows the temperature at which thermal-release starts, reaches a maximum and returns to the baseline. The temperature at which maximal release was reached was considered as the peak temperature, being specific for each compound. The accuracy of the proposed method was tested by determining the average value of 10 replicate determinations of the desorption temperature for HgS red (Table 2). A well defined single peak with an intermediate desorption temperature was observed for this compound. The differences between the expected value and the result obtained being lower than 10%. This indicates a good accuracy of the results. The precision of the analysis was evaluated from the results of standard deviation (SD) and relative standard deviations (%RSD). A good RSD value 6.6% was achieved while SD value is relatively high (Table 2). This fact can be attributed to the low homogeneity obtained during the sample preparation in which a low quantity of standard is diluted with a proportionately high amount of silica.

The profiles obtained are shown in Figs. 1–4. As can be seen Hg halides present desorption peaks at the lowest temperatures (Fig. 1). The decomposition of HgCl₂ occurs at low temperatures, the maximum peak being at 138 °C. The high peak corresponding to Hg₂Cl₂ is at 119 °C and exhibits a shoulder close to the peak temperature corresponding to HgCl₂. This shoulder is probably associated to the liberation of Hg₂Cl₂ [23].

The other Hg halides analyzed, HgI₂ and HgBr₂, have single peaks at approximately 100–110 °C while the profile corresponding to HgF₂ moves to higher temperatures with two peaks at 234 and 450 °C (Fig. 1). The peak at 450 °C is attributed to the total decomposition of HgF₂, while the peak at the lower temperature could be due to the decomposition of Hg compounds formed during the analysis such as Hg₂F₂, or HgFOH [30].

The desorption profiles obtained for the two forms of HgO, yellow and red, are shown in Fig. 2. A maximum peak appears at 471 °C and 284 °C and there are smaller peaks at about 308 °C and 469 °C for red and yellow HgO respectively. Red HgO consists of well-defined crystalline prismatic particles approximately 2 µm in size, which appear to be fused together, to form round multi-particle aggregates around 20 µm in diameter. Yellow HgO consists of smaller particles of around 1 µm that have less well-defined shape and form smaller aggregates about 5 µm in size. The specific surface areas were 0.45 and 0.68 m² g⁻¹ for the red and yellow HgO respectively as measured by N₂ adsorption (BET method). The smaller peak may be due the secondary decomposition of oxide

Table 2
Validation results obtained for HgS red analyzed by thermo-desorption technique. [Hg]=744 mg kg⁻¹.

| Sample | [Hg] (mg kg ⁻¹) |
|--------------------------|-----------------------------|
| 1 | 836 |
| 2 | 809 |
| 3 | 836 |
| 4 | 757 |
| 5 | 769 |
| 6 | 769 |
| 7 | 729 |
| 8 | 779 |
| 9 | 696 |
| 10 | 707 |
| X (mg Kg ⁻¹) | 765 |
| SD _{tot} | 51 |
| RSD (%) | 6.6 |




Fig. 1. Thermal-desorption profiles of the Hg-halides.




Fig. 2. Thermal-desorption profiles of the mercury oxides.




Fig. 3. Thermal-desorption profiles of the S-containing mercury compounds.




Fig. 4. Thermal-desorption profiles of the N-containing mercury compounds.

(I) Hg_2O during the analysis:



The Hg_2O forms at low temperatures ($< 230^\circ\text{C}$) as a result of the oxidation of the free atoms of Hg by atomic oxygen at the interface between the two solid phases ($\text{HgO}/\text{Hg}_2\text{O}$). Its formation is associated, with the formation of a reaction interface that separates the two solid phases ($\text{HgO}/\text{Hg}_2\text{O}$), and with the partial transfer of Hg_2O formation energy to the reactant, which enhances the decomposition of HgO [28].






It should be noted that in the case of yellow HgO , the two peaks have a similar intensity. This can be attributed to the fact that the yellow form of HgO is more reactive than the red form due to the smaller particle size and therefore higher specific surface of the yellow form. Its decomposition occurs more rapidly in this compound whereas the other mercury oxides undergo a change in their internal structure due to the rise in temperature before the total release of mercury.

The desorption profiles corresponding to the S-containing mercury compounds are shown in Fig. 3. Red HgS decomposes in one step at 305°C according to the chemical Eq. (3) whereas HgSO_4 decomposes at a higher temperature, 583°C .



Hg_2SO_4 undergoes a two-step decomposition, with one peak appearing at approximately 300°C and a second peak at 514°C slightly displaced from the HgSO_4 decomposition area. The presence of two intense peaks in the desorption profile suggests that the thermal decomposition of Hg_2SO_4 takes place via reactions (4)–(6). Eqs. (5) and (6) describe the thermal decomposition of HgSO_4 and of the unstable intermediate product $\text{HgSO}_4 \cdot 2\text{HgO}$ [29].



The desorption profiles of the N-containing mercury compounds are shown in Fig. 4. The thermal desorption of the mercury nitrates (I) and (II) is reflected in two peaks, a first peak at $264\text{--}280^\circ\text{C}$ and a second peak at 450°C , probably related to the decomposition of mercury nitrates in HgO which occur between 400 and 500°C (7)–(9). However, it must be taken into account that mercury (I) nitrate exhibits reducing properties. It is partially oxidized by atmospheric oxygen at ambient temperatures forming mercury nitrate (II) and mercury (9) so in its decomposition also could appear $\text{Hg}(\text{NO}_3)_2$.








Fig. 5. Thermal-desorption profiles of mercury(II) perchlorate hydrate.



As can be seen in Fig. 4 the peaks in the desorption profiles of the mercury nitrates (I) and (II) are in similar position but the peak corresponding to the decomposition of HgO (at 450 °C) is more intense in the profile of nitrate mercury (I) because its decomposition is generating more HgO.

The Hg(CN)₂ decomposition profile presents a broad peak at 267 °C. The thermal decomposition profile of Hg(SCN)₂ presents

two peaks, an intense peak at about 177 °C and a small peak at 288 °C close to the position of red HgS (305 ± 12 °C) probably related to the decomposition of Hg(SCN)₂ in HgS as a result of the thermal analysis (Pharaoh serpent reaction).



The desorption profile of mercury (II) perchlorate hydrate is shown in Fig. 5. The desorption of Hg(ClO₄)₂ · H₂O presents an intense peak at about 273 °C and two smaller peaks at 475 and 590 °C. The thermal decomposition of this compound could occur via reactions (11) and (12), decomposing either to form either the chloride and oxygen or the oxide and mixtures of chlorine and oxygen.





Table 3 summarizes the characteristic peak temperatures of each compound. These compounds are used as a reference.

3.2. Mercury species identification: mixtures of pure compounds

In order to evaluate possible interferences or interactions during the thermal release of the different mercury species, several Hg compound mixtures were analyzed. The results obtained (Fig. 6) confirm that different mercury species can be identified in a sample. No important interferences were observed when HgCl₂:HgSO₄; HgCl₂:HgS(red) and HgCl₂:Hg(NO₃)₂ · H₂O: HgSO₄ were tested (Fig. 6a–c). A slight deviation in the case of HgSO₄ was observed in the mixtures HgO:HgSO₄ and HgCl₂:HgO: HgSO₄ (Fig. 6d and e). In these two mixtures the maximum peak

Table 3
Thermal dissociation temperatures corresponding to the pure mercury compounds.

| Mercury compounds | High peak <i>T</i> (°C) | Start <i>T</i> –End <i>T</i> decomposition peak (°C) |
|---|-----------------------------|--|
| HgI ₂ | 100 ± 12 | 60–180 |
| HgBr ₂ | 110 ± 9 | 60–220 |
| Hg ₂ Cl ₂ | 119 ± 9 | 60–250 |
| HgCl ₂ | 138 ± 4 | 90–350 |
| HgS red | 305 ± 12 | 210–340 |
| HgF ₂ | 234 ± 42; 449 ± 12 | 120–350; 400–500 |
| HgO red | 308 ± 1; 471 ± 5 | 200–360; 370–530 |
| HgO yellow | 284 ± 7; 469 ± 6 | 190–380; 320–540 |
| Hg ₂ SO ₄ | 295 ± 4; 514 ± 4 | 200–400; 410–600 |
| HgSO ₄ | 583 ± 8 | 500–600 |
| Hg(SCN) ₂ | 177 ± 4; 288 ± 4 | 100–220; 250–340 |
| Hg(CN) ₂ | 267 ± 1 | 140–360 |
| Hg(NO ₃) ₂ · H ₂ O | 215 ± 4; 280 ± 13; 460 ± 25 | 150–370; 375–520 |
| Hg ₂ (NO ₃) ₂ · 2H ₂ O | 264 ± 35; 427 ± 19 | 120–375; 376–500 |
| HgCl ₂ O ₈ · H ₂ O | 273 ± 1; 475 ± 5; 590 ± 9 | 154–360; 380–510; 520–650 |




Fig. 6. Thermal-desorption profiles of the mixtures: (a) HgCl₂:HgSO₄ (1:1); (b) HgCl₂:HgS red (1:1); (c) HgCl₂:Hg(NO₃)₂ · H₂O: HgSO₄ (1:1:1); (d) HgO red: HgSO₄ (1:1) and (e) HgCl₂:HgO red: HgSO₄ (1:1:1).

Table 4

Comparison of expected and obtained Hg concentrations and recovery values for 5 replicates of red HgS analyzed by means of the thermo-desorption technique.

| Replicate | [Hg] theor. (mg kg ⁻¹) | [Hg] est. (mg kg ⁻¹) | Recovery (%) |
|-----------|------------------------------------|----------------------------------|--------------|
| 1 | 979 | 776 | 79 |
| 2 | 428 | 382 | 89 |
| 3 | 390 | 405 | 104 |
| 4 | 506 | 414 | 82 |
| 5 | 392 | 310 | 79 |

for HgSO₄ appears at a lower temperature. This suggests that an interaction between these compounds occurs. The mechanism of interaction is probably that represented in Eqs. (5) and (6).

3.3. Verification of the method

Due to the lack of reference materials for Hg species, red HgS, was used to verify the proposed method. Its thermal decomposition profile is shown in Fig. 3. Only one peak is produced, so the total mercury concentration obtained from calculation of the peak area (experimental) should fit the expected concentration. Table 4 shows the results obtained for the analysis of 5 sample replicates. Recoveries in the range of 79–104% were estimated suggesting that the method may be explored for quantitative purposes.

4. Conclusions

These results show thermal decomposition to be a promising technique for determining mercury speciation in solid samples. Good accuracy of the results was obtained. The differences between the expected value and the result obtained being lower than 10%. A good RSD value 6.6% was achieved while SD value is relatively high. This fact can be attributed to the low homogeneity obtained during the sample preparation.

The results show it is possible to identify the mercury species from their temperature of decomposition. The temperature decomposition rate of the mercury species was arranged in increasing order as follows: HgI₂ < HgBr₂ < Hg₂Cl₂ = HgCl₂ < Hg(CN)₂ < HgCl₂O₈ · H₂O < Hg(SCN)₂ < HgS (red) < HgF₂ < Hg₂(NO₃)₂ · 2H₂O < Hg(NO₃)₂ · H₂O < HgO (yellow, red) < Hg₂SO₄ < HgSO₄.

No important interferences were observed in the mixtures tested, apart from slight deviations when HgSO₄ and HgO were present together in a sample. This fact is attributed to the formation of an intermediate compound.

Acknowledgments

This work was financed by the PCTI Asturias, the European Regional Development Fund in Asturias (FEDER 2007–2013) and HUNOSA through the project PC10-20. We would like to acknowledge the support of CSIC for providing M. Rumayor with a JAE-Predoc fellowship and M.A. Lopez-Anton with a JAE-Doc contract (European Social Fund).

References

- [1] N. Pirrone, S. Cinnirella, X. Feng, R.B. Finkelman, H.R. Friedli, J. Leaner, R. Mason, A.B. Mukherjee, G.B. Stracher, D.G. Streets, K. Telmer, Atmos. Chem. Phys. 10 (2010) 5951–5964.
- [2] UNEP Global Mercury assessment 2013, Published by UNEP, Available online at www.unep.org/PDF/PressReleases/GlobalMercuryAssessment2013.pdf, 2013.
- [3] A. Kolker, C.L. Senior, J.C. Quick, Appl. Geochem. 21 (2006) 1821–1836.
- [4] J.H. Pavlish, L.L. Hamre, Y. Zhuang, Fuel 89 (2010) 838–847.
- [5] N.S. Bloom, E. Preus, in: Proceedings of the 2nd International Symposium on Contaminated Sediments, 2003, pp. 331–336.
- [6] R. Fernández-Martínez, M.I. Rucandio, Anal. Bioanal. Chem. 375 (2003) 1089–1096.
- [7] N.W. Revis, T.R. Osborne, D. Sedgley, A. King, Analyst 114 (1989) 823–825.
- [8] H. Sakamoto, T. Tomiyasu, N. Yonehara, Anal. Sci. 8 (1992) 35–39.
- [9] R. Rubio, G. Rauret, J. Radioanal. Nucl. Chem. 208 (1996) 529–540.
- [10] J.R. Bacon, C.M. Davidson, Analyst 133 (2008) 25–46.
- [11] A.T. Reis, S.M. Rodrigues, C.M. Davidson, E. Pereira, A.C. Duarte, Chemosphere 81 (2010) 1369–1377.
- [12] C.S. Kim, G.E. Brown, J.J. Rytuba, Sci. Total Environ. 261 (2000) 157–168.
- [13] C.S. Kim, J.J. Rytuba, G.E. Brown, J. Colloid Interface Sci. 271 (2004) 1–15.
- [14] C.S. Kim, S. Bloom, J.J. Rytuba, G.E. Brown, Environ. Sci. Technol. 37 (2003) 5102–5108.
- [15] H. Biester, M. Gosar, G. Müller, J. Geochem. Explor. 65 (1999) 195–204.
- [16] H. Biester, G. Nehrke, J. Fresenius, Anal. Chem. 358 (1997) 446–452.
- [17] A. Bollen, A. Wenke, H. Biester, Water Res. 42 (2008) 91–100.
- [18] H. Biester, C. Scholz, Environ. Sci. Technol. 31 (1997) 233–239.
- [19] H. Biester, M. Gosar, S. Covelli, Environ. Sci. Technol. 34 (2000) 3330–3336.
- [20] Olga V. Shuvaeva, Maria A. Gustaytis, Gennadii N. Anoshin, Anal. Chim. Acta 621 (2008) 148–154.
- [21] A.T. Reis, J.P. Coelho, S.M. Rodrigues, R. Rocha, C.M. Davidson, A.C. Duarte, E. Pereira, Talanta 99 (2012) 363–638.
- [22] M. Ozaki, Md.A. Uddin, E. Sasaoka, S. Wub, Fuel 87 (2008) 3610–3615.
- [23] C. Raposo, C.C. Windmöller, W.A.D. Júnior, Waste Manage. 23 (2003) 879–886.
- [24] X. Feng, J.Y. Lu, D.C. Gregoire, Y. Hao, C.M. Banic, W.H. Schroeder, Anal. Bioanal. Chem. 380 (2004) 683–689.
- [25] J. Li, Y. Yuan, R. Perry, M.M. Maroto-Valer, Prepr. Pap.—Am. Chem. Soc., Div. Fuel Chem. 52 (2) (2007) 511–512.
- [26] M. Rallo, M.A. Lopez-Anton, R. Perry, M.M. Maroto-Valer, Fuel 89 (2010) 2157–2159.
- [27] M.A. Lopez-Anton, Y. Yang, R. Perry, M.M. Maroto-Valer, Fuel 89 (2010) 629–634.
- [28] B.V. Lvov, Thermochim. Acta 333 (1999) 21–26.
- [29] S.A. Tariq, J.O. Hill, J. Therm. Anal. 21 (1981) 277–281.
- [30] B. Brunetti, V. Piacente, A. Latini, P. Scardala, J. Chem. Eng. Data 53 (2008) 2493–2495.