



0040-4039(94)02204-6

Palladium-catalysed Synthesis of α -Diallylated Ketosulphides of Benzothiazole and their Transformation into Diallyl Thiiranes and Trienes

Vincenzo Caló^{a,*}, Vito Fiandanese^a, Angelo Nacci^a and Antonio Scilimati^b


^aCNR Centro di Studio sulle Metodologie Innovative di Sintesi Organiche, Dipartimento di Chimica, ^bDipartimento Farmaco-Chimico, Università di Bari, via Orabona 4-70125 Bari (Italy)

Abstract. α -Ketosulphides of benzothiazole **1** react with allylic carbonates in the presence of palladium acetate in dichloromethane under mild conditions affording α,α -diallylated ketosulphides **2** in high yields. Reduction of **2** with sodium borohydride in isopropanol gives episulphides which in turn can be transformed almost quantitatively into trienes.

Palladium catalysed allylic substitution is a versatile process encompassing a wide range of allyl systems and their nucleophilic partners¹⁻⁴. Among allylic systems, the use of carbonates as leaving groups has gained in popularity since the development of these reagents by Tsuji.³ The initially displaced carbonate loses carbon dioxide generating an alkoxide which is sufficiently basic to deprotonate active methylene compounds as malonates or β -ketoesters. Contrary to silyl enol ethers or enamines,⁵ simple ketones cannot be allylated satisfactorily with allylic carbonates. On the other hand the introduction of two allylic groups at the α -position of ketones, which can be done in principle by regioselective consecutive allylation is generally difficult since the generation of an anion at the same carbon after the first allylation is inhibited and as a consequence α,α' -diallylated products are obtained.⁶


















We wish to report that α -ketosulphides of benzothiazole⁷ **1** react with allylic carbonates in the presence of palladium acetate and triphenylphosphine in dichloromethane under mild conditions to afford α,α -diallylated ketosulphides **2** in high yields. (Scheme 1).

Scheme 1



As reported in table 1, stereoselective allylic rearrangement are observed in the reactions of **1a,b** with some allylic carbonates to give only the E isomers (products **2c-e**). Allylation of these ketosulphides with β -methallyl carbonate afforded only monoallylation products (**2f,g**) even in the presence of large excess of the carbonate. This is probably due to the steric hindrance of the monoallylated ketosulphide. α -Substituted ketone **1c** gave monoallylated products (**2h-j**) as expected.


Table 1. Pd⁰-Mediated Allylation of Ketosulphides of benzothiazole by Allylic Carbonates.^a

Carbonate	Ketone	Product ^b	Yield(%) ^c	Carbonate	Ketone	Product ^b	Yield(%)
	1a		85		1a		88
"	1b		86	"	1b		91
	1a		78 ^d	"			93
	1a		77 ^d		1c		96
"	1b		82 ^d		1c		71

^aAll reactions were run under nitrogen at r.t. in dichloromethane during 1-3 h. [carbonate]: [ketone]: [Pd]: [phosphine] = 3: 1: 0.05: 0.25. ^bAll products gave the expected ¹H, ¹³C NMR, IR and mass spectra. ^cIsolated yields. ^dE-isomer.

The carbonyl reduction of these compounds with sodium borohydride in isopropanol (Scheme 2) gave directly diallylated episulphides together with 2-hydroxybenzothiazole as byproduct, which can be separated by washing the ethereal extract with sodium carbonate solution.

Scheme 2




This reaction is characteristic for β -hydroxy sulphides of benzothiazole as reported previously by us⁷ whereas other different β -hydroxy sulphides did not react under the same reaction conditions. The results are reported in table 2.

Table 2. Synthesis of episulphides from allylated ketosulphides of benzothiazole.^a

Ketone	Episulphide ^b	Yield(%) ^c (E:Z ratio) ^d	Ketone	Episulphide	Yield(%) ^c (E:Z ratio) ^d
2a		88	2b		85
2c		78	2d		90
2e		85	2f		70 (60:40)
2g		75 (87:13)	2h		60 (59:41)
2i		40 (57:43)	2j		65 (85:15)

^aAll reactions were run at r.t. in isopropanol during 5h. [Ketone]: [NaBH₄]= 1: 1. ^bAll products gave the expected ¹H, ¹³C NMR, IR and mass spectra. ^cIsolated yields. ^dEvaluated by ¹H NMR.







Reaction of dieny episulphides with 3-methyl-2-selenoxobenzothiazole^{8,9} **5** in dichloromethane or with triphenyl phosphine in boiling THF afforded trienes almost quantitatively (Scheme 3).



Some results are reported in table 3.

Beside the synthesis of trienes or dienes which can be further elaborated, the allylated ketones (**2a-j**) can be conveniently desulfurized by reaction with tributyltin hydride in boiling toluene^{10,11} to γ,δ -unsaturated ketones which are important intermediates for example for the synthesis of spiro ketals upon peracid oxidation of the double bond.

Table 3. Conversion of allylic episulphides into trienes^a.

Entry	Episulphide	Triene	Yield(%) ^b
1			95
2			97
3			98

^aAll reactions were run in dichloromethane at r.t. [Episulphide]: **5**: Trifluoroacetic acid]: 1: 1: 2. ^bYields evaluated by GC. All products gave the expected ¹H, ¹³C NMR, IR and mass spectra.

General procedure

(a) **Allylation of ketosulphides.** A mixture of palladium acetate (0.05 mmol), triphenylphosphine (0.25 mmol) carbonate (3 mmol) and ketosulphide (1 mmol) in CH₂Cl₂ (15 ml) was stirred under a nitrogen atmosphere at r.t. until the disappearance of the sulphide (TLC). Concentration in vacuo and chromatographic purification (SiO₂) of the residue provided the diallylated ketone(s). (b) **Synthesis of episulphides.** To the allylated ketone (0.9 mmol) dissolved in isopropanol (10 ml) was added portionwise NaBH₄ (0.9 mmol) under stirring at r.t. After the disappearance of the ketone, the solution was evaporated and the residue dissolved in ether was washed with a 5% Na₂CO₃ solution and dried on Na₂SO₄. Evaporation of the solvent leaves almost pure episulphide. (c) **Synthesis of trienes.** The episulphide (0.9 mmol), 2-selenoxobenzothiazole **5** (0.9 mmol) and trifluoroacetic acid (1.8 mmol) dissolved in 5 ml of dichloromethane were stirred at r.t. until the disappearance of the episulphide (TLC). After the filtration of selenium, concentration in vacuo and chromatography (SiO₂) provided the triene.

Acknowledgement: This work was supported in part by grants of MURST (40 and 60%).

References and notes

1. Trost, B. M. *Acc. Chem. Res.* **1980**, *13*, 385-393.
2. Tsuji, J. *Tetrahedron*, **1986**, *42*, 4361-4401 and references therein.
3. Tsuji, J.; Minami, I. *Acc. Chem. Res.* **1987**, *20*, 140-145.
4. Frost, C.G.; Howarth, J.; Williams, J.M.J. *Tetrahedron : Asymmetry*, **1992**, *3*, 1089-1122.
5. Hiroi, K.; Abe, J.; Suya, K.; Sato, S.; Koyama, T. *J. Org. Chem.* **1994**, *59*, 203-213.
6. Bates, R. B.; Taylor, S. R. *J. Org. Chem.* **1994**, *59*, 245-246.
7. Caló, V.; Lopez, L.; Pesce, G. *Gazz. Chim. Ital.* **1979**, *109*, 703-704.
8. Caló, V.; Lopez, L.; Mincuzzi, A.; Pesce, G. *Synthesis*, **1976**, 200-201.
9. This reagent is now commercially available.
10. Ueno, Y.; Okawara, M. *J. Am. Chem. Soc.* **1979**, *101*, 1893-1894.
11. Caló, V. Unpublished results.

(Received in UK 7 October 1994; revised 1 November 1994; accepted 4 November 1994)