



New Approaches to the Asymmetric Synthesis of Dipeptide Isosteres via β -Lactam Synthon Method

Iwao Ojima,* Hong Wang, Tao Wang and Edward W. Ng

Department of Chemistry, State University of New York at Stony Brook, NY 11794-3400, USA.


Received 23 September 1997; accepted 7 November 1997

Abstract: New and efficient synthetic routes to dipeptide isosteres with high enantiomeric purity, e.g., hydroxyethylene, dihydroxyethylene and hydroxyethylamine isosteres, have been developed via oxiranes **6** and formyloxazolines **13** derived from *N*-t-Boc- β -lactams **4**. © 1998 Elsevier Science Ltd. All rights reserved.

It has been shown that hydroxyethylene, dihydroxyethylene and hydroxyethylamine dipeptide isosteres are essential building blocks¹ for the inhibitors of such enzymes as renin and HIV-1 protease.² These dipeptide isosteres provide effective transition state mimics of the substrates for the peptidases, which bind to the enzymes tightly and inhibit their actions.³ Thus, a number of methods has been reported for the synthesis of these dipeptide isosteres.⁴ In the course of our study on the exploration of the " β -Lactam Synthon Method" (β -LSM),⁵ we have developed new and efficient methods for the syntheses of norstatine and its analogs⁶ as well as norstatine-dipeptides and their analogs through novel ring-opening coupling of *N*-acyl- β -lactams,⁷ which is applicable to resin-bound amino acids for solid phase peptide syntheses.⁸ This type of ring-opening coupling of *N*-acyl- β -lactams has also been successfully applied to the efficient and practical syntheses of taxane antitumor agents⁹ and novel hydroxy(keto)ethylene dipeptide isosteres.¹⁰ We have further investigated the useful transformations of isoserines readily obtained by methanolysis of *N*-acyl- β -lactams and synthesized enantiopure 3-t-Boc-aminoalkene-1,2-oxides and 5-formyl-1,3-oxazolines. We communicate here new and efficient routes to hydroxyethylene, dihydroxyethylene and hydroxyethylamine dipeptide isosteres with high enantiomeric purity via these oxiranes and formyloxazolines derived from 1-t-Boc-3-siloxy- β -lactams.


Enantiopure (3*R*,4*S*)-1-(t-Boc)-3-siloxy- β -lactams **4b-d** were readily obtained through efficient chiral ester enolate-imine cyclocondensations, followed by removal of *p*-methoxyphenyl (PMP) and protection as carbamates in the same manner as reported previously from these laboratories. Due to the acidity of the benzylic protons of the aldimine, the corresponding 4-benzyl- β -lactam **4a** could not be obtained directly through the cyclocondensation route.¹¹⁻¹³ However, **4a** can be readily prepared from 4-isobutenyl β -lactam (Scheme 1). The ozonolysis of **1** gave, after reductive workup, the corresponding aldehyde which was subsequently treated with phenylmagnesium bromide. The resulting secondary alcohol was acetylated to give **2** in 66% overall yield. Although only one stereoisomer was obtained, the absolute stereochemical configuration of **2** has not been determined yet. Hydrogenolysis of **2** under 100 psi of hydrogen gave the desired 4-benzyl- β -lactam **3** in 74% yield. Removal of the PMP group of **3**⁹ and protection as a carbamate gave **4a** in 87% overall yield.

Scheme 1



As Scheme 2 shows, methanolysis of *t*-Boc- β -lactams **4a–d** gave *N*-*t*-Boc-O-TIPS-isoserine methyl esters **5a–d** in 91–97% yields. 4-Isobutenylisoserine **5d** was converted to the corresponding isobutylisoserine (norstatine) methyl ester **5e** by hydrogenation on Pd-C in 99% yield. In order to synthesize oxiranes **6**, **5a–e** were reacted with LiAlH_4 (LAH) and the resulting diols were subjected to the Mitsunobu reaction¹⁴ to give oxiranes **6a–e** in 43–77% yields for two steps. Regioselective opening of oxiranes **6a–e** afforded the corresponding alcohols, which were then protected with 2,2-dimethoxypropane to give oxazolines **7a–e** in 36–80% yields for two steps. Ozonolysis of the alkene moieties of **7a–c** and **7e** gave, after oxidative workup, the *N,O*-protected hydroxyethylene isosteres **8a–c** and **8e** in 61–74% overall yields. Due to the presence of another olefin moiety in isobutenylisoserine **7d** which is susceptible to ozonolysis, an alternative procedure was employed, i.e., Wacker oxidation¹⁵ followed by treatment with sodium hypochlorite, to give **8d** in 74% yield. Deprotection of **8a** with *p*-toluenesulfonic acid in MeOH gave *N*-*t*-Boc-hydroxyethylene isostere **9a** in 83% yield. Thus, other oxazolines **8b–e** should give the corresponding *N*-*t*-Boc-hydroxyethylene isostere **9b–e**.


Scheme 2



Hydroxyethylamine isosteres can also be obtained from oxirane **6d**. Ring opening-coupling of **6d** with (*S*)-proline and (*S*)-phenylalanine methyl esters gave the corresponding hydroxyethylamine isosteres **10d** and **11d** in 76% and 74% yields, respectively (Scheme 3).

As Scheme 4 illustrates, the other useful intermediate, 5-formyl-1,3-oxazolines **13**, were also synthesized from **5a–e**. Removal of TIPS of **5a–e** followed by treatment with 2,2-dimethoxypropane / pyridinium *p*-toluenesulfonate (PPTS) gave oxazoline esters **12a–e** in 72–82% yields for two steps. Reduction of the ester moiety with diisobutylaluminum hydride (DIBALH) followed by Swern oxidation gave formyloxazolines **13a–e** in 78–87% overall yields.

Scheme 3



Lewis acid mediated allylation of **13a-e** gave adducts **14a-e** with moderate to good *anti*-selectivities (Table 1). The absolute configuration of **14e** was determined by X-ray analysis (Figure 1). Ozonolysis of **14a-c** and **14e** followed by oxidative workup gave *N,O*-protected dihydroxyethylene isosteres **15a-c** and **15e** in 56-71% yields. Oxazoline **14d** (R^1 = isobutenyl) was subjected to Wacker oxidation followed by treatment with sodium hypochlorite to give **15d** in 53% yield. Deprotection of **15a** (R^1 = PhCH_2) gave dihydroxyethylene isostere **16a** in 77% yield.

Scheme 4





Table 1 Lewis acid mediated allyl addition to formyloxazolines 13

entry	R^1	Lewis acid	allylating reagent	solvent	yield (%)	<i>anti</i> : <i>syn</i>
1	isobutenyl	TiCl_4	allyltributyltin	CH_2Cl_2	50	2:3
2		SnCl_4	allyltributyltin		87	2:3
3 ¹¹			allyltrichlorotin		55	8:1
4		CrCl_2	allyl bromide	THF	84	3:2
5	cyclohexylmethyl	BF_3OEt_2	allyltributyltin	CH_2Cl_2	75	7:1
6		$\text{MgBr}_2\cdot\text{Et}_2\text{O}$			71	3:1
7		CrCl_2	allyl bromide	THF	84	2.2:1
8	benzyl	BF_3OEt_2	allyltributyltin	CH_2Cl_2	70	6.5:1
9		SnCl_4			85	3:1

Figure 1 X-ray structure of 14e 

Further studies on the asymmetric synthesis of a variety of dipeptide isosteres via *N*-acyl- β -lactams 4, oxiranes **6**, formyloxazolines **13** and their applications to the synthesis of enzyme inhibitors are actively underway.

knowledge: This research was supported by a grant from the National Institutes of Health (NIGMS)

REFERENCES AND NOTES

- (a) Thaisrivongs, S.; Pals, D. T.; Kroll, L. T.; Turner, S. R.; Han, F.-S. *J. Med. Chem.* **1987**, *30*, 976-982. (b) Boyd, S. A.; Mantei, R. A.; Hsiao, C.-N.; Baker, W. R. *J. Org. Chem.* **1991**, *56*, 438-442. (c) Rosenberg, S. H.; Boyd, S. A.; Mantei, R. A. *Tetrahedron Lett.* **1991**, *32*, 6507-6508.
- (a) Huff, J. R. *J. Med. Chem.* **1991**, *34*, 2309-2314. (b) Jung, M. J. In *Chemistry and Biochemistry of Amino Acids*; G. C. Barret, Ed.; Chapman and Hall: New York, 1985; pp 227-296.
- (a) Rich, D. H. *J. Med. Chem.* **1985**, *28*, 263-273.
- (a) Ojima, I.; Chen, H. J. C.; Qui, X. *Tetrahedron* **1988**, *44*, 5307-5318. (b) Ojima, I.; Chen, H.-J. C.; K., N. *J. Am. Chem. Soc.* **1988**, *110*, 278-281. (c) Ojima, I.; Pei, Y. *Tetrahedron Letters* **1990**, *31*, 977-980. (d) Ojima, I.; Komata, T.; Qiu, X. *J. Am. Chem. Soc.* **1990**, *112*, 770-774. (e) Ojima, I.; Zhao, M.; Yamato, T.; Nakahashi, K.; Yamashita, M.; Abe, R. *J. Org. Chem.* **1991**, *56*, 5263-5277.
- (a) Georg, G. I.; Ravikumar, V. T. In *The Organic Chemistry of β -Lactam Antibiotics*; G. I. Georg, Ed.; VCH Publishers: New York, 1992; pp 295-368. (b) Ojima, I. *Acc. Chem. Res.* **1995**, *28*, 383-389.
- (a) Ojima, I.; Park, Y. H.; Sun, C. M.; Zhao, M.; Brigaud, T. *Tetrahedron Lett.* **1992**, *33*, 5737-5740.
- (a) Ojima, I.; Sun, C. M.; Park, Y. H. *J. Org. Chem.* **1994**, *59*, 1249-1250.
- (a) Ojima, I.; Sun, C. M.; Zucco, M.; Park, Y. H.; Duclos, O.; Kuduk, S. *Tetrahedron Letters* **1993**, *34*, 4149-4152.
- (a) Ojima, I.; Habus, I.; Zhao, M.; Zucco, M.; Park, Y. H.; Sun, C. M.; Brigaud, T. *Tetrahedron* **1992**, *48*, 6985-7012.
- (a) Ojima, I.; Ng, E. W.; Sun, C. M. *Tetrahedron Letters* **1995**, *36*, 4547-4550.
- (a) Ojima, I. In *The Organic Chemistry of β -Lactam Antibiotics*; G. I. Georg, Ed.; VCH Publishers: New York, 1992; pp 197-255.
- (a) Palomo, C.; Aizpurua, J. M.; Mielgo, A.; Linden, A. *J. Org. Chem.* **1996**, *61*, 9186-9195.
- (a) Kobayashi, Y.; Takemoto, Y.; Ito, Y.; Terashima, S. *Tetrahedron Letters* **1990**, *31*, 3031-3034.
- (a) Castejon, P.; Pasto, M.; Moyano, A.; Pericas, M. A.; Riera, A. *Tetrahedron Letters* **1995**, *36*, 3019-3022.
- (a) Tsuji, J. *Synthesis* **1984**, 369-384.