



Heterogeneous Rhodium-Catalyzed Hydrogenation Conditions for the Highly Effective Synthesis of 1,3-Oxazolidines from 1,2-Amino Alcohols and Nitriles .1#


Stéphane Létinois, Jean-Christophe Dumur, Françoise Hénin*, Jacques Muzart*

Unité Mixte de Recherche "Réactions Sélectives et Applications",
CNRS - Université de Reims Champagne-Ardenne, B.P. 1039, 51687 Reims Cédex 2, France

Received 11 December 1997; accepted 19 January 1998


Abstract: A procedure easily carried out for the synthesis of 1,3-oxazolidines in high yields using 1,2-amino alcohols, nitriles, an atmospheric pressure of hydrogen and catalytic quantities of rhodium on carbon powder is presented. This reaction probably involves the semi-hydrogenation of the nitrile followed by condensation with the amino alcohol. The process may constitute the key step in a two-step sequence for reducing a nitrile into the corresponding aldehyde. © 1998 Elsevier Science Ltd. All rights reserved.

Recently, we reported a one-pot *N*-alkylation procedure of 1,2-amino alcohols through their reaction with nitriles in the presence of palladium on charcoal and hydrogen. Under these conditions, 1,3-oxazolidines are formed as intermediates and they undergo hydrogenolysis of the NC-O bond to afford *N*-alkylated-amino alcohols.² We suspect that the heterocyclization involves firstly hydrogenation of the nitrile to the corresponding imine,^{3,4} and secondly trapping of this imine by the amino alcohol competing with its further reduction (Scheme 1).



Scheme 1

1,3-Oxazolidines are usually obtained from the condensation of 1,2-amino alcohols with either aldehydes or their corresponding acetals.⁵ They are useful as intermediates in organic synthesis.^{5,6} Moreover, it has been recently shown that non-racemic, chiral 1,3-oxazolidines can be effective ligands for metal-catalyzed enantioselective procedures.⁷ Therefore, it would be desirable to find appropriate conditions to selectively and effectively synthesize oxazolidines from amino alcohols and nitriles in a one-pot procedure. With this aim, we have examined the use of heterogeneous catalysts other than palladium in the above reaction. We are now delighted to report that this goal has been attained by using rhodium on carbon powder (Eq. 1).



R: CH₃ (a), CH₂CH₃ (b), (CH₂)₂CH₃ (c), (CH₂)₃CH₃ (d), CH₂Ph (e),


Ph (f), (CH₂)₈CH₃ (g), *c*-C₆H₁₁ (h), CH(CH₃)₂ (i).

This paper is dedicated to Pr J.-P. Pète on the occasion of his 60th birthday.

When a solution of (-)-ephedrine **1** (0.12 M) in acetonitrile containing catalytic quantities of Rh/C⁸ was stirred at room temperature under an atmosphere of hydrogen, complete consumption of the amino alcohol was observed after 7 h. Filtration of the mixture over a short pad of silica gel and evaporation of the solvents gave oxazolidine **4a**⁹ in 95% yield. When using a solvent such as toluene and an amount of **3a** reduced to 10 equiv. (with respect to **1**), the reaction remained very effective, since a 92% yield of the desired product was recovered (Table 1, run 1). As outlined by runs 2-4 and 14-17, similar results were obtained with propionitrile, *n*-butyronitrile and valeronitrile in reaction with either **1** or **2**. In contrast, the use of heavier (runs 5, 7 and 10) or secondary nitriles (run 12) led to a sluggish condensation under analogous conditions: the conversion of **1** was greatly reduced and this was similarly observed when exchanging toluene for methanol or ethyl acetate. Nevertheless, satisfying results were obtained when the reactions in ethyl acetate were performed at reflux instead of room temperature. These conditions provided a 100% consumption of **1** with virtually quantitative formation of **4** even when only 5 equiv. of **3** were used (runs 6, 8, 9, 11 and 13).

The observed diastereomeric ratios obtained for oxazolidines **4** and **5** (Table 1) suggest a thermodynamically controlled reaction: the tautomeric equilibrium between oxazolidines and their open form is in favour of the *cis*-heterocyclic diastereomer as demonstrated by studies on **4a**⁹, **4f**¹¹ and **5f**.¹⁶ In fact, we were unable to identify the open form of **5a-5d** among the reaction products when **2** was used as aminoalcohol.

Nevertheless, careful analysis of the ¹H NMR spectrum of the crude reaction mixture obtained from **1** and **3f** showed the presence of *N*-benzylidenebenzylamine (δ : 4.81 and 8.37 ppm) and dibenzylamine (δ : 3.85 ppm) as side-products while benzylamine (δ : 3.80 ppm) was not detected. As mentioned above, hydrogenation of nitriles and imines into amines are competing reactions to the trapping of imines.¹⁷ In the present reaction, the imine **6f**, formed by hydrogenation of **3f**, is reduced to the primary amine **7f** or trapped either by the amino alcohol giving **4f** or by the amine **7f** leading to **8f** (Scheme 2). Compound **8f** gave **10f** either by hydrolysis⁴ or *via* the imine **9f**.^{4,20} These observations led us to investigate the hydrogenation of **3f** in the absence of **1** (**3f**: 3 mmol, Rh/C: 30 mg, AcOEt, 80°C): a slow reaction took place (the conversion of **3f** was incomplete after 80h) to afford the substituted imine **9f** and the secondary amine **10f**; as above, no primary amine was detected. The large increase in the **10f/9f** ratio with time (16 h: 0.3, 80 h: 4.9) suggested that **10f** was at least in part derived from **9f**.^{4,20} The comparison of the competing reactions showed that the hydrogenation steps were much slower than those of the imine condensation.



Scheme 2

Oxazolidines are masked aldehydes, the latter being produced by hydrolysis of the former.^{5,21} Therefore, we examined the possibility of reducing nitriles into the corresponding aldehydes *via* oxazolidines. For this purpose, an excess of amino alcohol *versus* nitrile was required. Thus, benzonitrile was reacted with

10 equiv. of either (-)-ephedrine or *N*-methylaminoethanol for 48 h (Rh/C: 0.012 equiv., EtOAc, 80°C). After elimination of the catalyst by filtration over Celite and concentration, the crude mixture was hydrolyzed^{21b} leading to benzaldehyde with respectively 71 or 65% overall yields. In conclusion, this two-step procedure offers an efficient alternative to other methods of reducing nitriles.²²

Table 1.: Condensation of various nitriles with (-)-ephedrine (**1**) and (+)-norephedrine (**2**).

Runs	Amino alcohol	RCN (equiv.)	Method ^a	Time h	Conversion % of 1 or 2	Oxazolidine yield %	<i>cis</i> / <i>trans</i> ¹⁰
1	1	3a (10)	A	8	100	4a ⁹ : 92 ^b	93 / 7
2	"	"	"	10	"	4b ¹² : 91 ^b	88 / 12
3	"	3c (10)	"	20	"	4c ¹³ : 94 ^b	92 / 8
4	"	3d (10)	"	"	"	4d : 89 ^b	89 / 11
5	"	3e (5 or 10)	"	16	~ 37	4e : n.d. ^c	n.d. ^c
6	"	3e (5)	B	8	100	4e ¹⁴ : 95 ^d	91 / 9
7	"	3f (5)	A	20	74	4f : n.d. ^c	82 / 18
8	"	3f (3)	B	16	100	4f ¹¹ : 96 ^d	91 / 9
9	"	3g (5)	"	15	"	4g : 97 ^d	91 / 9
10	"	3h (5)	A	24	22	4h : n.d. ^c	86 / 14
11	"	3h (5)	B	21	100	4h : 92 ^d	91 / 9
12	"	3i (10)	A	20	70	4i : n.d. ^c	n.d. ^c
13	"	3i (10)	B	15	100	4i ^{12,15} : 91 ^b	88 / 12
14	2	3a (10)	A	7	"	5a : 99 ^b	73 / 27
15	"	3b (10)	"	7	"	5b : 97 ^b	75 / 25
16	"	3c (10)	"	24	"	5c : 97 ^b	72 / 28
17	"	3d (10)	"	"	"	5d : 99 ^b	68 / 32

^aThe mixture of amino alcohol (0.6 mmol), nitrile (1.8-6 mmol), Rh/C (30 mg) and solvent (5 ml) was stirred under an atmosphere of hydrogen (rubber balloon); A: in toluene at RT; B: in ethyl acetate at 80°C. ^bIsolated yields. ^cNot determined. ^dYields determined by ¹H NMR using dibenzylether (δ : 4.55 ppm) as internal standard.

Acknowledgments: We are grateful to "Ministère de l'Education Nationale" for a PhD studentship to S.L., to Engelhard Company for generous gifts of rhodium on charcoal and to Dr G. Bird (Zeneca Pharma, Reims) for help with the text.

References and notes:

- Presented in part in the 9th IUPAC Organometallic Chemistry directed towards Organic Synthesis, Göttingen, Germany; July, 21-25, 1997 *Poster N° 25*.
- Hénin, F.; Létinois, S.; Muzart, J. *Tetrahedron Lett.* **1997**, *38*, 7187-7190.
- Rylander, P. *Catalytic Hydrogenation in Organic Syntheses* Acad. Press: New York, 1979, p. 138-152.
- Volf, J.; Pasek, J. *Studies in Surface Science and Catalysis* Cerveny, L., Ed.; Elsevier: Amsterdam, 1986, Vol. 27, p. 105-144.
- Bergmann, E.D. *Chem. Rev.* **1953**, *53*, 309-352.
- For recent studies, see a) ref. 5 in². b) Le Bail, M.; Pérard, J.; Aitken, D.J.; Bonin, M.; Husson, H.-P. *Tetrahedron Lett.* **1997**, *38*, 7177-7180. c) Heaney, H.; Papageorgiou, G.; Wilkins, R.F. *Tetrahedron* **1997**, *53*, 14381-14396. d) Enders, D.; Reinhold, U. *Tetrahedron: Asymmetry* **1997**, *8*, 1895-1946.
- a) Simons, K.E.; Wang, G.; Heinz, T.; Giger, T.; Mallat, T.; Pfalz, A.; Baiker, A. *Tetrahedron: Asymmetry* **1995**, *6*, 505-518. b) Dai, W.-M.; Zhu, H.J.; Hao, X.-J. *Tetrahedron: Asymmetry* **1996**, *7*, 1245-1248. c) Prasad, K.R.K.; Joshi, N.N. *J. Org. Chem.* **1997**, *62*, 3770-3771.
- 5% (On dry basis) Rh on carbon powder Escat 30, code: 8000 from Engelhard Company was used throughout this work. This catalyst has a surface area of 1100 m²/g and contains 55 % of water (Technical information from Engelhard Company).
- Beckett, A.H.; Jones, G.R. *Tetrahedron* **1977**, *33*, 3313-3316.
- The *cis*-relative configuration at C-2 for the main diastereoisomer of **4** was deduced from literature data¹¹ and for **5**, by analogy. The *cis/trans* ratios were mainly determined from the relative integrations of the NMR signals corresponding to the hydrogen *gem* to the C-5 phenyl group.
¹H NMR:
 - *cis* **5a**: 0.73 (*Me*(4), d, *J* 6.9), 1.57 (*Me*(2), d, *J* 5.3), 1.84 (NH, m), 3.63 (*H*(4), m), 4.73 (*H*(2), q, *J* 5.3), 4.94 (*H*(5), d, *J* 7.6), 7.25 (*Ph*, m).
 - *trans* **5a**: 0.74 (*Me*(4), d, *J* 6.9), 1.42 (*Me*(2), d, *J* 5.7), 1.84 (NH, m), 3.74 (*H*(4), m), 5.02 (*H*(5), d, *J* 6.5), 5.20 (*H*(2), d, *J* 5.7), 7.25 (*Ph*, m).
 - *cis* **5b**: 0.73 (*Me*(4), d, *J* 6.5), 1.13 (*MeCH*₂, t, *J* 7.6), 1.89 (*CH*₂*Me*, m), 1.89 (NH, m), 3.63 (*H*(4), m), 4.52 (*H*(2), t, *J* 5.7), 4.90 (*H*(5), m), 7.25 (*Ph*, m).
 - *trans* **5b**: 0.74 (*Me*(4), d, *J* 6.9), 1.03 (*MeCH*₂, t, *J* 7.6), 1.69 (*CH*₂*Me*, m), 1.89 (NH, m), 3.69 (*H*(4), m), 4.90 (*H*(5) and *H*(2), m), 7.25 (*Ph*, m).
- Agami, C.; Rizk, T. *Tetrahedron* **1985**, *41*, 537-540 and ref. therein.
- Pfanz, H.; Kirchner, G. *Justus Liebigs Ann. Chem.* **1958**, 149-159.
- Engelhardt, E.L.; Crossley, F.S.; Sprague, J.M. *J. Am. Chem. Soc.* **1950**, *72*, 2718-2723.
- Chiringhelli, D.; Bernardi, L. *Tetrahedron Lett.* **1967**, 1039-1042.
- Bureau, R.; Mortier, J.; Joucla, M. *Bull. Soc. Chim. Fr.* **1993**, *130*, 584-596.
- Fülop, F.; Bernath, G.; Mattinen, J.; Pihlaja, K. *Tetrahedron* **1989**, *45*, 4317-4324.
- The trapping of *in situ* produced imines by primary amines is commonly applied to the synthesis of symmetrical secondary amines.^{3,4,18} Another interesting application concerns the synthesis of heterocyclic compounds.¹⁹
- a) Galan, A.; de Mendoza, J.; Prados, P.; Rojo, J.; Echavarren, A. *J. Org. Chem.* **1991**, *56*, 452-454. b) Giannandrea, R.; Mastroilli, P.; Zaccaria, G.; Nobile, C.F. *J. Mol. Catal. A: Chem.* **1996**, *109*, 113-117.
- Diker, K.; Döe de Maindreville, M.; Lévy, J. *Tetrahedron Lett.* **1995**, *36*, 2497-2500 and 3511-3512.
- Eleved, M. B.; Hogeveen, H.; Schudde, E. P. *J. Org. Chem.* **1986**, *51*, 3635-3642.
- a) Ganter, C.; Brassat, L.; Ganter, B. *Chem. Ber./Recueil* **1997**, *130*, 659-662. b) Myers, A.G.; Yang, B.H.; Chen, H.; McKinstry, L.; Kopecky, D.J.; Gleason, J.L. *J. Am. Chem. Soc.* **1997**, *119*, 6496-6511.
- a) March J. *Advanced Organic Chemistry* 4th edition, J. Wiley: New York **1992**, p. 919-920. b) Toujas, J.L.; Jost, E.; Vaultier, M. *Bull. Soc. Chim. Fr.* **1997**, *134*, 713-717 and ref. therein.