



Pergamon

Tetrahedron Letters 39 (1998) 3225–3228

TETRAHEDRON
LETTERS

Convenient Synthesis of Pyrano[3,2-*c*]quinolines and Indeno[2,1-*c*]quinolines by Imino Diels-Alder Reactions

Govindarajulu Babu and Paramasivan T. Perumal*

Organic Chemistry Division
Central Leather Research Institute, Adyar, Chennai - 600 020, INDIA.

Received 12 January 1998; revised 17 February 1998; accepted 19 February 1998

Abstract: Anhydrous indium trichloride (InCl_3) is found to catalyze the imino Diels-Alder reaction of 3,4-dihydro-2*H*-pyran and indene with Schiff's bases to afford pyrano[3,2-*c*]quinolines and indeno[2,1-*c*]quinolines. Imino Diels-Alder reactions of *N*-benzylidene 1-naphthylamine with 3,4-dihydro-2*H*-pyran and indene results in efficient synthesis of phenanthridine derivatives.


© 1998 Elsevier Science Ltd. All rights reserved.

Pyrano quinoline moiety is present in many alkaloids.¹ Pyrano quinoline derivatives possess wide range of biological activities such as psychotropic activity,² antiallergic activity,³ antiinflammatory activity,⁴ estronegic activity⁵ and are used as a potential pharmaceuticals.⁶ Indeno quinoline derivatives also possess wide range of biological activities such as 5-HT-receptor binding activity,⁷ antiinflammatory activity,⁸ and also act as antitumor agents⁹ and inhibitor for steroid reductase.¹⁰ Imino Diels-Alder reactions provide an easy entry into quinoline derivatives.¹¹ Recently, we have shown that indium trichloride activates the imines¹² in the imino Diels-Alder reaction of Schiff's bases with cyclopentadiene and cycloalkenones.

Few reports are available for the synthesis of pyrano- and indeno quinolines by imino Diels-Alder reactions.¹³ Herein, we wish to report the synthesis of pyrano- and indeno quinolines by imino Diels-Alder reactions of 3,4-dihydro-2*H*-pyran and indene with Schiff's bases catalyzed by indium trichloride (InCl_3).

In the presence of 20 mol % anhydrous indium trichloride (InCl_3), *N*-benzylidene aniline **1a** was treated with 3,4-dihydro-2*H*-pyran **2** in acetonitrile at room temperature. After 30 minutes we obtained pyrano quinolines **3a** and **4a** in a ratio of 41:59 in an overall yield of 80 % (Scheme 1).

Scheme 1



Pyrano quinolines **3** and **4** were obtained in varying ratios for imines derived from *p*-substituted anilines. In the case of imines derived from 2,4- and 2,5-disubstituted anilines, pyrano quinoline derivative **4** was obtained as major compound. The results are summarized in Table 1.

Table 1. Reaction of Schiff's bases with 3,4-dihydro-2*H*-pyran (**2**) employing 20 mol% InCl₃.^a


Schiff's base	Substituents			Time, min.	Product ^b ratio of 3:4	Overall yield (%)
	R ¹	R ²	R ³			
1a	H	H	H	30	41:59	80
1b	H	Cl	H	30	34:66	50
1c	H	OCH ₃	H	240	58:42	70
1d	H	CH ₃	H	120	68:32	70
1e	H	Cl	CH ₃	40	5:95	46
1f	CH ₃	H	CH ₃	180	5:95	45

a: All reactions were conducted at room temperature by addition of 20 mol % InCl₃ to a mixture of Schiff's base and 3,4-dihydro-2*H*-pyran in acetonitrile.

b: Products were characterized by Mass, IR, ¹H NMR and ¹³C NMR. The ratio based on isolation by chromatography.


Indium trichloride also catalyzes effectively the imino Diels-Alder reaction of 3,4-dihydro-2*H*-pyran with *N*-benzylidene 1-naphthylamine **5** and resulted in phenanthridine derivatives **6** and **7** in a ratio of 32:68 in an overall yield of 53 % (Scheme 2).

Scheme 2



Next, *N*-benzylidene aniline **1a** in acetonitrile was stirred at room temperature with indene **8** in the presence of 20 mol % indium trichloride. After 6 h, indenoquinoline **9a** was obtained in moderate yield (Scheme 3). Only one isomer was obtained in the reaction of Schiff's bases with indene and the results are summarized in Table 2.

Scheme 3

Table 2 : Reaction of Schiff's bases with indene (8) employing 20 mol % InCl_3 .^a

Schiff's base	Product ^b	Substituents			Time, h	Yield (%) ^c
		R ¹	R ²	R ³		
1a	9a	H	H	H	6	40
1b	9b	H	Cl	H	6	48
1c	9c	H	OCH ₃	H	24	30
1e	9e	H	Cl	CH ₃	12	48
1f	9f	CH ₃	H	CH ₃	9, reflux	65


a : All reactions were conducted at room temperature by addition of 20 mol % InCl_3 to a mixture of schiff's bases and indene in acetonitrile.

b : Products were characterized by Mass, IR, ^1H NMR and ^{13}C NMR.

c : Isolated yield.

When *N*-benzylidene 1-naphthylamine 5 in acetonitrile was treated with indene 8 in the presence of 20 mol % InCl_3 , indeno phenanthridine¹⁴ 10 was obtained in 50 % yield (Scheme 4).

Scheme 4



In conclusion, we have shown that indium trichloride catalyzes the imino Diels-Alder reaction of Schiff's bases with 3,4-dihydro-2*H*-pyran and indene which provides a convenient synthesis of pyrano

[3,2-*c*]quinolines and indeno[2,1-*c*]quinolines. Application of imino Diels-Alder reaction to *N*-benzylidene 1-naphthylamine results in novel synthesis of phenanthridine derivatives.

Acknowledgement: We thank Council of Scientific and Industrial Research, New Delhi, India, for financial support.

REFERENCES AND NOTES

1. Ramesh, M.; Mohan, P. S.; Shanmugam, P. *Tetrahedron* **1984**, *40*, 4041.
2. Nesterova, I. N.; Alekseeva, L. M.; Andreeva, L. M.; Andreeva, N. I.; Golovira, S. M.; Granik, V. G. *Khim. -Farm. Zh.* **1995**, *29*, 31 (Russ); *Chem. Abstr.* **1996**, *124*, 117128t.
3. a) Takahashi, K. *et al. Oyo Yakuri* **1986**, *32*, 233(Jpn); *Chem. Abstr.* **1986**, *105*, 218691j b) Yamada, N.; Kadowaki, S.; Takahashi, K.; Umezu, K. *Biochem. Pharmacol.* **1992**, *44*, 1211.
4. Faber, K.; Stueckler, H.; Kappe, T. *J. Heterocycl. Chem.* **1984**, *21*, 1177.
5. Akhmed Khodzhaeva, Kh. S.; Bessonova, I. A. *Dokl. Akad. Nauk Uzh. SSR*, **1982**, 34-36 (Russ); *Chem. Abstr.* **1983**, *98*, 83727q.
6. Mohamed, E. A. *Chem. Pap.* **1994**, *48*, 261; *Chem. Abstr.* **1995**, *123*, 9315x.
7. Anzini, M.; Cappelli, A.; Vomero, S.; Cagnotto, A.; Skorupska, M. *Med. Chem. Res.* **1993**, *3*, 44.
8. Quraishi, M. A.; Thakum, V. R.; Dhawan, S. N. *Indian J. Chem. Sect. B* **1989**, *28B*, 891.
9. Okazaki, S.; Asao, T.; Wakida, M.; Jshida, K.; Washinosu, M.; Utsugi, T.; Yamada, Y. *Jp. Appl.* **94/107**, 190, **1994**; *Chem. Abstr.* **1996**, *124*, 232268a.
10. Brooks, J. R.; Berman, C.; Hichens, M.; Primka, R. L.; Reynolds, G. F.; Rasmusson, G. H. *Proc. Soc. Exp. Biol. Med.* **1982**, *169*, 67.
11. a) Weinreb, S. M. *Comprehensive Organic Synthesis*, Trost, B. M.; Fleming, I., Ed., Pergamon; Oxford, **1991**, Vol.5, p401. b) Boger, D. L.; Weinreb, S. M. *Hetero Diels-Alder Methodology in Organic Synthesis*, Academic; San Diego, **1987**, Chaps. 2 and 9.
12. a) Babu, G.; Perumal, P. T. *Tetrahedron Lett.* **1997**, *38*, 5025. b) Babu, G.; Perumal, P. T. *Tetrahedron* (In press).
13. a) Makioka, Y.; Shindo, T.; Taniguchi, Y.; Takaki, K.; Fujiwara, Y. *Synthesis* **1995**, 801. b) Gilchrist, T. L.; Stannard, A. M. *Tetrahedron Lett.* **1988**, *29*, 3585. c) Cabral, D.; Laszlo, P.; Montaufier, M. T. *Tetrahedron Lett.* **1988**, *29*, 547. d) Cabral, J.; Laszlo, P. *Tetrahedron Lett.* **1989**, *30*, 7237. e) Kametani, T.; Takeda, H.; Suzuki, Y.; Honda, T. *Synth. Commun.* **1985**, *15*, 499. f) Borriane, E.; Prato, M.; Scorrano, G.; Stivanello, M. *J. Heterocycl. Chem.* **1988**, *25*, 1831. g) Borriane, E.; Prato, M.; Scorrano, G.; Stivanello, M. *J. Chem. Soc. Perkin Trans. 1* **1989**, 2245.
14. Selected spectral data for compound **10** : ^1H NMR (300 MHz, CDCl_3) δ 7.80-7.08(m, 15H), 4.85(br s, 1H), 4.75(d, 1H, $J = 7.2\text{Hz}$), 4.71(br s, 1H, NH), 3.47(m, 2H), 2.50(m, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 146.2, 142.9, 142.7, 139.4, 132.6, 128.6, 128.5, 127.5, 127.3, 126.9, 126.7, 126.1, 125.2, 125.1, 125.0, 124.9, 119.6, 118.5, 117.6, 57.7, 48.1, 46.8, 30.9; MS(m/e): 347(M^+); IR (KBr) : 3414, 3065, 1577 cm^{-1} .