

A New and Simple ‘LEGO’ System for the Synthesis of 2,6-Oligopyridines

Gunther R. Pabst and Jürgen Sauer*,¹

Institut für Organische Chemie der Universität Regensburg, D-93040 Regensburg, Germany

Received 5 June 1998; accepted 6 July 1998

Abstract

The condensation of α -arylglyoxals with carboxamidrazones **1** - **2** is the best method for the synthesis of aryl or hetaryl substituted 1,2,4-triazines **3** - **4**. These 1,2,4-triazines can be easily transformed to pyridines by [4+2] cycloaddition with bicyclo[2.2.1]hepta-2,5-diene followed by [4+2] cycloreversions of nitrogen and cyclopentadiene. This reaction sequence offers a new, simple and general access to 2,6-oligopyridines **8** - **11**.

© 1998 Elsevier Science Ltd. All rights reserved.

Keywords: Cycloadditions; Oligomers; Pyridines; Triazines

Terpyridines and higher oligopyridines have been demonstrated to be useful building blocks in metallosupramolecular chemistry [1,2,3,4]. In order to investigate the coordinating behaviour of these polydentate ligands it is necessary to develop direct high-yield synthetic methods for easy access. Up to now there are two different approaches for the synthesis of oligopyridine ligands: the coupling of pyridine units [5,6,7,8] or halopyridine nuclei [9] and the construction of central pyridine rings [10,11]. Here we report on a new and simple ‘LEGO’ system for the synthesis of linear 2,6-oligopyridines **8** - **11** and hetaryl substituted 1,2,4-triazines **3** - **4** as their precursors.

3,5-Disubstituted 1,2,4-triazines are easily prepared by heating of carboxamidrazones with α -aryl-glyoxals in ethanol under reflux for 4-6 hours [12,13]. But these reaction conditions fail for 3,5-di-(2-pyridyl)-1,2,4-triazines because of the thermolabile 2-pyridylglyoxal, which is only available in aqueous solution [14]. Therefore, we have treated 2-pyridylglyoxal with the corresponding carboxamidrazone in ethanol-water (4:1 v/v) for 3-9 days at ambient temperature with eventually heating of the isolated precipitate in *N,N*-dimethylformamide at 100°C for 0.5 hours to complete condensation (Table 1, Scheme 1).

¹ E-mail: elisabeth.liebl@chemie.uni-regensburg.de


The concentration of 2-pyridylglyoxal was determined by ^1H NMR spectroscopy of its aqueous solution with 3-trimethylsilyl-(2,2,3,3- d_4)-propionic-acid-sodium salt as internal and integral standard.

Table 1. Mono-, bi- and bis-[1,2,4]-triazines synthesized according to Scheme 1.

Carboximidrazone [Ref.]	Triazine	Reaction Times and Conditions	Yield [%]	M.P. [°C]
1a [15]		3a 	1 d, r.t.	49 162-165
1b [16]		3b 	2 d, r.t.	85 220-222
2a [17]		4a 	3 d, r.t.; 2 h 100°C in DMF	85 336-339
1c analogue to [16]		3c 	2 d, r.t.	72 246-248
2b [18]		4b 	3 d, r.t.	67 293-298
2c analogue to [18]		4c 	8 d, r.t. 2 h 100°C in DMF	60 303-308
2d [19], analogue to [16]		4d 	9 d, r.t.; 2 h 100°C in DMF	71 395-398

Typical procedure for the preparation of 1,2,4-triazines **3 - 4**: **1b** (1.58 g, 7.40 mmol) and 2-pyridylglyoxal (1.00 g, 7.40 mmol, $c = 33.3 \mu\text{mol/l}$) in 222 ml ethanol-water (4:1 v/v) were stirred at ambient temperature. After 3 days the resulting yellow precipitate (**3b**) was collected by suction filtration, washed several times with water and ethanol and dried for 16 hours at 80°C/0.01 Torr. No further purification was necessary. Analytical data for **3b**: IR (KBr): $\tilde{\nu} = 3080, 3060, 3000, 1570, 1550, 1525, 1505, 1460, 1420, 1350, 1235, 1030, 980, 755, 725 \text{ cm}^{-1}$. ^1H NMR (400 MHz, CDCl_3): $\delta = 7.37$ (ddd, 1 H, $J = 7.5 \text{ Hz}$, $J = 4.8 \text{ Hz}$, $J = 1.2 \text{ Hz}$), 7.53 (ddd, 1 H, $J = 7.6 \text{ Hz}$, $J = 4.7 \text{ Hz}$, $J = 1.2 \text{ Hz}$), 7.90 (ddd, 1 H, $J = 8.0 \text{ Hz}$, $J = 7.5 \text{ Hz}$, $J = 1.8 \text{ Hz}$), 8.00 (ddd, 1 H, $J = 7.8 \text{ Hz}$, $J = 7.7 \text{ Hz}$, $J = 1.8 \text{ Hz}$), 8.09 (dd, 1 H, $J = 7.8 \text{ Hz}$, $J = 7.8 \text{ Hz}$), 8.66 (dd, 1 H, $J = 7.9 \text{ Hz}$, $J = 1.1 \text{ Hz}$), 8.70 (dd, 1 H, $J = 7.8 \text{ Hz}$, $J = 1.0 \text{ Hz}$), 8.73 (ddd, 1 H, $J = 4.8 \text{ Hz}$, $J = 1.8 \text{ Hz}$, $J = 0.9 \text{ Hz}$), 8.77 (ddd, 1 H, $J = 8.0 \text{ Hz}$, $J = 1.2 \text{ Hz}$, $J = 0.9 \text{ Hz}$), 8.81 (ddd, 1 H, $J = 7.8 \text{ Hz}$, $J = 1.2 \text{ Hz}$, $J = 0.9 \text{ Hz}$), 8.83 (ddd, 1 H, $J = 4.7 \text{ Hz}$, $J = 1.8 \text{ Hz}$, $J = 0.9 \text{ Hz}$), 10.34 (s, 1 H) ppm; EI-MS (70eV): m/z (%): 312 (35) [M^+], 284 (78) [$\text{M}^+ - \text{N}_2$], 236 (4) [$\text{M}^+ - \text{C}_5\text{H}_2\text{N}$], 206 (5) [$\text{M}^+ - \text{C}_5\text{H}_2\text{N} - \text{N}_2$], 181 (100) [$\text{M}^+ - \text{C}_7\text{H}_4\text{N} - \text{N}_2$], 155 [$\text{C}_{10}\text{H}_7\text{N}_2$], 128 (11) [$\text{C}_7\text{H}_6\text{N}$], 103 (95) [$\text{C}_7\text{H}_4\text{N}$], 78 (20) [$\text{C}_5\text{H}_4\text{N}$]; $\text{C}_{18}\text{H}_{12}\text{N}_6$ (312.3): calcd. C 69.22, H 3.87, N 26.91; found C 69.13, H 4.01, N 26.80. All other 1,2,4-triazines were characterized by the same analytical methods.






1,2,4-Triazines are known to participate as electron poor dienes in inverse-type Diels-Alder reactions with electron rich and angle strained dienophiles to yield dihydropyridine and pyridine derivatives after extrusion of molecular nitrogen [13,20].



Scheme 1. Reaction sequence for the synthesis of pyridines via 1,2,4-triazines

Acetylene as dienophile is too unreactive, unpracticable and dangerous. Therefore, we used bicyclo[2.2.1]hepta-2,5-diene (10 fold excess) in refluxing 1,2-dichlorobenzene as a synthetic equivalent for acetylene (Scheme 1).

Table 2. Synthesis of 2,6-oligopyridines according to Scheme 1.

Triazine	Oligopyridine	Reaction Conditions and Times	Yield [%]	M.P. [°C]	Reference	
3a	8		140°C, 1h	50	87-88	[7]
4a	9a		140°C, 1h	79	208-210	[8]
4b	10a		140°C, 3h	60	261-263	[8]
4c	10b		ΔT, 14 h	77	219-221	-
4d	11		ΔT, 168 h	78	326-329	[8]

Typical procedure for the preparation of oligopyridines **8 - 11**: **4a** (236 mg, 750 μmol) and bicyclo[2.2.1]hepta-2,5-diene (1.38 g, 15.0 mmol) were heated at 140°C under an inert atmosphere in 15 ml 1,2-dichlorobenzene for 1 hour. The reaction mixture was cooled and the resulting precipitate was collected by suction filtration, washed with 2 ml 1,2-dichlorobenzene and 40 ml petroleum ether 40/60. Recrystallization from benzene yielded **9a** as colorless crystals. Analytical data for **9a**: IR (KBr): ν = 3040, 3000, 1575, 1560, 1550, 1455, 1430, 1410, 1250, 1095, 1060, 755 cm⁻¹. ¹H NMR (400 MHz, CDCl₃): δ = 7.35 (ddd, 2 H, J = 7.5 Hz, J = 4.8 Hz, J = 1.3 Hz), 7.87 (ddd, 2 H, J = 8.0 Hz, J = 7.5 Hz, J = 1.8 Hz), 8.01 (dd, 2 H, J = 7.9 Hz, J = 7.8 Hz), 8.48 (dd, 2 H, J = 7.8 Hz, J = 1.1 Hz), 8.67 (ddd, 2 H, J = 8.0 Hz, J = 1.3 Hz, J = 0.9 Hz), 8.68 (dd, 2 H, J = 7.8 Hz, J = 1.1 Hz), 8.72 (ddd, 2 H, J = 4.8 Hz, J = 1.8 Hz, J = 0.9 Hz) ppm; All other oligopyridines were characterized in the same way.

The NMR spectra of 1,2,4-triazines **3** and **4** exhibit in each case one singlet for the expected triazine-H⁶ and the corresponding pyridines are confirmed by the expected coupling constants for 2,6-oligopyridines (footnote Table 2).

Recent investigations have shown that 1,2,4-triazines undergo [4+2] cycloaddition with ethynyltributyltin to furnish 4-tributylstannyl-pyridines [21,22], which lead to a variety of 4-functionalized pyridines. Further work on this topic with 1,2,4-triazines **3** - **4** is in progress.

Acknowledgements

We are grateful to the Deutsche Forschungsgemeinschaft (DFG) and BASF AG for financial support of this research.

References

- [1] Constable EC. Progress in Inorganic Chemistry, Vol. 42, Karlin KD, editor, New York: John Wiley & Sons, 1994; 42: 67-138.
- [2] Constable EC. Metals and Ligands, Weinheim: VCH, 1996.
- [3] Sauvage JP, Collin JP, Chambron JC, Guillerez S, Condret C, Balzani V, Barigelli F, DeCola L, Flamingi L. Chem. Rev. 1994; 94: 993-1020.
- [4] Lehn JM. Supramolecular Chemistry - Concepts and Perspectives, Weinheim: VCH, 1995.
- [5] Morgan G, Burstall FH. J. Chem. Soc. 1937: 1649.
- [6] Badger GM, Sasse WHF. J. Chem. Soc. 1956: 616.
- [7] Burstall FH. J. Chem. Soc. 1938: 1662-1673.
- [8] Cardenas DJ, Sauvage JP. Synlett 1996: 916-918.
- [9] Tiecco M, Testaferri L, Tingoli M, Chianelli D, Montanucci M. Synthesis 1984: 736.
- [10] Kröhnke F. Synthesis 1976; 1: 1-24.
- [11] Negoro T, Oae S. Rev. Heteroatom Chem. 1995; 13: 235.
- [12] Culbertson BM, Parr GR. J. Heterocyclic Chem. 1967; 4: 422-424.
- [13] Neunhoeffer H. 1,2,4-Triazines and their Benzo Derivatives, Comprehensive Heterocyclic Chemistry II, Katritzky AR, Rees CW, Scriven EFV, editors, Oxford: Pergamon Press, 1996; 6: 507-574.
- [14] Schank K. Chem. Ber. 1969; 102: 383-387.
- [15] Hage R, Prins R, Haasnot JG, Reedijk J, Vos JG. J. Chem. Soc. Dalton Trans 1987: 1389-1396.
- [16] Case FH. J. Org. Chem. 1966; 31: 2398-2400, but prolonged reaction times (7-14 days).
- [17] Dedichen G. Avhandl. Norske Videnskaps-Akad. Oslo, I, Mat.-Naturv. Klasse 1936, No. 5.
- [18] Case FH. J. Heterocyclic Chem. 1971; 8: 1043-1046.
- [19] Stanek J, Caravatti G, Capraro HG, Furet P, Mett H, Schneider P, Regenass U. J. Med. Chem. 1993; 36 (1): 46-54.
- [20] Sauer J. 1,2,4,5-Tetrazines, Comprehensive Heterocyclic Chemistry II, Katritzky AR, Rees CW, Scriven EFV, editors, Oxford: Pergamon Press, 1996; 6: 901-957.
- [21] Sauer J, Heldmann DK. Tetrahedron Lett. 1998; 39: 2549-2552.
- [22] Heldmann DK, Pabst GR, Sauer J. *submitted to Eur. J. Org. Chem.* 1998.