

A New and Simple ‘LEGO’ System for the Synthesis of Branched Oligopyridines

Gunther R. Pabst, Konrad Schmid and Jürgen Sauer*,¹

Institut für Organische Chemie der Universität Regensburg, D-93040 Regensburg, Germany

Received 5 June 1998; accepted 6 July 1998

Abstract

The condensation of carboxamidrazones **1** with 1,2-dicarbonyl compounds **2** is the best method for the synthesis of alkyl, aryl or hetaryl substituted 1,2,4-triazines **3** - **7**. These 1,2,4-triazines can be easily transformed to pyridines **8** - **12** by [4+2] cycloaddition with bicyclo[2.2.1]hepta-2,5-diene followed by [4+2] cycloreversions of nitrogen and cyclopentadiene. This reaction sequence offers a new, simple and general access to branched oligopyridines. © 1998 Elsevier Science Ltd. All rights reserved.

Keywords: Cycloadditions; Oligomers; Pyridines; Triazines

A number of reports deal with the synthesis and reactions of polydentate oligopyridines [1,2,3,4,5]. In this communication we extend our new and simple ‘LEGO’ system [6] to the synthesis of branched oligopyridines and 3,5,6-trisubstituted 1,2,4-triazines as their precursors.

1,2,4-Triazines are easily prepared by heating of carboxamidrazones **1a** - **1e** (Table 1) with 1,2-dicarbonyl compounds **2a** - **2c** (Table 1) in ethanol under reflux for 4-6 hours (Table 2) [7].


Table 1. Starting compounds for synthesis of 1,2,4-triazines: Amidrazones and 1,2-dicarbonyl compounds

Compound	Ref.	Compound	Ref.
1a pyridine-2-carboxamidrazone	[8]	1e benzene-1,4-dicarboxbisamidrazone	[11]
1b dicarboxbisamidrazone	[9]	2a 1,2-bis-(2-pyridyl)-ethane-1,2-dione	Fluka
1c pyridine-2,6-dicarboxbisamidrazone	[10]	2b 1,2-bis-(2,2'-bipyridin-6-yl)-ethane-1,2-dione	[12]
1d 2,2'-bipyridine-6,6'-dicarboxbisamidrazone	[6]	2c 1,2-bis-(2,2',6'-terpyridin-6-yl)-ethane-1,2-dione	[12]

1,2,4-Triazines are known to participate as electron poor dienes in inverse-type Diels-Alder reactions with electron rich and angle strained dienophiles to yield dihydropyridine and pyridine derivatives after extrusion of molecular nitrogen [6, 13, 14].

¹ E-mail: elisabeth.liebl@chemie.uni-regensburg.de

We used bicyclo[2.2.1]hepta-2,5-diene (10 fold excess) in refluxing toluene or p-xylene as a synthetic equivalent for acetylene (Scheme 1).



Scheme 1. Reaction sequence for the synthesis of pyridines via 1,2,4-triazines











For 1,2,4-triazines with higher molecular weights reaction times were too long and yields too low. In these cases the reactions were performed in 1,2-dichlorobenzene as solvent at 140°C leading to shorter reaction times and higher yields. The results are collected in Table 3.

Table 2. Mono-, bi- and bis-[1,2,4]-triazines synthesized according to Scheme 1.

Carbox-amidrazone	Dicarbonyl compound	Triazine [Reference]	R	Yield [%]	M.P. [°C]
1a	2a	3 [6]		-	80 160-162
1b	2a	4a [6]	H	85	276-278
1b	2b	4b	pyridin-2-yl	88	>280
1b	2c	4c	2,2'-bipyridin-6-yl	80	175-180
1c	2a	5a	H	94	254-256
1c	2b	5b	pyridin-2-yl	97	282-283
1d	2a	6	H	83	331-333
1e	2a	7a	H	64	>280
1e	2b	7b	pyridin-2-yl	90	283-285
1e	2c	7c	2,2'-bipyridin-6-yl	80	269-270

Analytical data for **5b**: IR (KBr): $\tilde{\nu}$ = 3060, 3020, 1575, 1560, 1510, 1495, 1470, 1450, 1425, 1405, 1370, 1305, 1250, 1145, 1090, 1075, 1055, 1035, 1015, 985, 825, 775, 735, 695 cm^{-1} . ^1H NMR (250 MHz, CDCl_3): δ = 7.12-7.21 (m, 4 H), 7.31 (ddd, 2 H, J = 8.0 Hz, J = 1.2 Hz, J = 1.0 Hz), 7.28-7.34 (m, 6 H), 8.09 (dd, 2 H, J = 7.9 Hz, J = 7.8 Hz), 8.11 (dd, 2 H, J = 7.9 Hz, J = 7.8 Hz), 8.29 (dd, 1 H, J = 8.0 Hz, J = 7.7 Hz), 8.40 (dd, 2 H, J = 7.8 Hz, J = 1.1 Hz), 8.41 (dd, 2 H, J = 7.9 Hz, J = 1.2 Hz), 8.42 (ddd, 2 H, J = 8.0 Hz, J = 1.3 Hz, J = 1.0 Hz), 8.52 (ddd, 2 H, J = 4.8 Hz, J = 1.8 Hz, J = 0.9 Hz), 8.54 (ddd, 2 H, J = 4.8 Hz, J = 1.8 Hz, J = 0.9 Hz), 8.61 (dd, 2 H, J = 7.7 Hz, J = 1.1 Hz), 9.02 (d, 2 H, J = 7.9 Hz) ppm; EI-MS (70eV); m/z (%): 853 (3) [M^+], 825 (7) [$\text{M}^+ + \text{H} - \text{N}_2$], 491 (23) [$\text{M}^+ - \text{C}_{22}\text{H}_{14}\text{N}_4$], 334 (51) [$\text{C}_{22}\text{H}_{14}\text{N}_4$], 333 (51) [$\text{C}_{22}\text{H}_{14}\text{N}_4 - \text{H}$], 180 (100) [$\text{C}_{12}\text{H}_8\text{N}_2$], 156 (16) [$\text{C}_{10}\text{H}_7\text{N}_2$]; $\text{C}_{51}\text{H}_{31}\text{N}_{15}$ (853.9); calcd. C 71.73, H 3.66, N 24.62; found C 71.26, H 3.66, N 24.39. All other 1,2,4-triazines were characterized by the same analytical methods.

Table 3. Synthesis of branched oligopyridines according to Scheme 1.

Triazine	Pyridine		Reaction conditions and times	Yield [%]	M.P. [°C]
3	8		p-xylene, ΔT, 3 d	45	122-123
4a	9a		toluene, ΔT, 3 d	88	273
4b	9b		p-xylene, ΔT, 15 h	88	260-262
4c	9c		toluene, ΔT, 24 h	82	169-170
5a	10a		1,2-dichlorobenzene, 140°C, 1 h	85	243-244
5b	10b		1,2-dichlorobenzene, 140°C, 6 h	94	250-252
6	11		1,2-dichlorobenzene, 140°C, 1 d	81	303-305
7a	12a		toluene, ΔT, 6 d	49	266-268
7b	12b		p-xylene, ΔT, 3 d	63	254
7c	12c		toluene, ΔT, 5 d	80	>280

Typical procedure for the preparation of pyridines **8** - **12**: **5b** (250 mg, 293 μmol) and norborna-2,5-diene (0.54 g, 5.86 mmol) were heated at 140°C under an inert atmosphere in 15 ml 1,2-dichlorobenzene for 6 hours. The reaction mixture was cooled and treated with petroleum ether 40/60 until it became cloudy. After standing in a refrigerator for several hours the precipitate was collected by suction filtration, washed with petroleum ether 40/60 and recrystallized from acetonitrile to furnish **10b** as colorless crystals.

Analytical data for **10b**: IR (KBr): $\tilde{\nu}$ = 3040, 3000, 1565, 1550, 1460, 1445, 1425, 1410, 1365, 1250, 1140, 1105, 1085, 1070, 1035, 1015, 980, 820, 805, 765, 735 cm^{-1} . ^1H NMR (400 MHz, CDCl_3): δ = 7.09-7.14 (m, 2 H), 7.28 (ddd, 2 H, J = 7.5 Hz, J = 4.8 Hz, J = 1.2 Hz), 7.30-7.34 (m, 4 H), 7.36 (dd, 2 H, J = 7.8 Hz, J = 1.7 Hz), 7.71 (dd, 2 H, J = 7.8 Hz, J = 7.8 Hz), 7.72 (ddd, 2 H, J = 7.8 Hz, J = 7.5 Hz, J = 1.8 Hz), 8.01 (dd, 2 H, J = 7.9 Hz, J = 7.7 Hz), 8.05 (t, 1 H, J = 7.8 Hz), 8.19 (ddd, 2 H, J = 8.0 Hz, J = 1.1 Hz, J = 1.1 Hz), 8.27 (dd, 2 H, J = 7.8 Hz, J = 1.0 Hz), 8.32 (d, 2 H, J = 8.1 Hz), 8.30-8.35 (m, 4 H), 8.52 (ddd, 2 H, J = 4.8 Hz, J = 1.7 Hz, J = 1.0 Hz), 8.66 (ddd, 2 H, J = 4.8 Hz, J = 1.8 Hz, J = 0.9 Hz), 8.75 (d, 2 H, J = 7.8 Hz), 8.89 (d, 2 H, J = 8.1 Hz) ppm; ^{13}C -NMR (100 MHz, CDCl_3 , DEPT): δ = 118.86 (2 C, +), 119.84 (2 C, +), 120.41 (2 C, +), 120.98 (2 C, +), 121.27 (2 C, +), 121.65 (2 C, +), 123.52 (2 C, +), 123.67 (2 C, +), 124.06 (2 C, +), 124.15 (2 C, +), 136.20 (2 C, 0), 136.48 (2 C, +), 136.76 (2 C, +), 136.80 (2 C, +), 137.59 (1 C, +), 137.83 (2 C, +), 140.08 (2 C, +), 148.68 (2 C, +), 148.97 (2 C, +), 154.31 (2 C, 0), 155.00 (2 C, 0), 155.09 (2 C, 0), 155.11 (2 C, 0), 155.67 (2 C, 0), 155.74 (2 C, 0), 156.14 (2 C, 0), 157.25 (2 C, 0), 158.54 (2 C, 0) ppm. EI-MS (70 eV): m/z (%): 850 (50) $[(\text{M} + \text{H})^+]$, 849 (100) $[\text{M}^+]$, 848 (87) $[(\text{M} - \text{H})^+]$, 821 (2) $[\text{M}^+ - \text{N}_2]$, 772.4 (41) $[\text{M}^+ - \text{C}_5\text{H}_3\text{N}]$, 771 (71) $[\text{M}^+ - \text{C}_5\text{H}_4\text{N}]$, 744 (17) $[\text{M}^+ - \text{C}_5\text{H}_3\text{N} - \text{N}_2]$, 425 (16) $[\text{M}^{2+}]$, 385 (12) $[(\text{M} - \text{C}_5\text{H}_4\text{N})^+]$, 57 (8), 55 (12), 44 (40), 40 (55): $\text{C}_{55}\text{H}_{35}\text{N}_{11}$ (850.0): calcd. C 77.72, H 4.15, N 18.13; found C 76.97, H 4.44, N 17.79. All other oligopyridines were characterized by the same analytical methods.

Further investigations are in progress to examine the variability of this new synthesis with respect to different carboxamidrazones and 1,2-dicarbonyl compounds and the capability of these oligopyridines to act as ligands in metal complexes.

Acknowledgements

We are grateful to the Deutsche Forschungsgemeinschaft (DFG) and BASF AG for financial support of this research.

References

- [1] Lawrence DS, Jiang T, Levett M. *Chem. Rev.* 1995; 95: 2229-2260.
- [2] Lehn JM. *Supramolecular Chemistry*, Weinheim: VCH, 1995.
- [3] Constable EC. *Progress in Inorganic Chemistry*, Vol. 42, Karlin KD, editor, New York: John Wiley & Sons, 1994; 42: 67-138.
- [4] Constable EC. *Metals and Ligands*, Weinheim: VCH, 1996.
- [5] Sauvage JP, Collin JP, Chambron JC, Guillerez S, Condret C, Balzani V, Barigelli F, DeCola L, Flammingi L. *Chem. Rev.* 1994; 94: 993-1020.
- [6] Pabst GR, Sauer J. *Tetrahedron Lett.* 1998; preceding publication.
- [7] *For a typical procedure see:* Case FH. *J. Org. Chem.* 1965; 30: 931-933; but prolonged heating up to 6 hours.
- [8] Hage R, Prins JG, Reedijk J, Vos JG. *J. Chem. Soc. Dalton Trans.* 1987; 1389-1396.
- [9] Dedichen G. *Avhandl. Norske Videnskaps-Akad. Oslo, I, Mat.-Naturv. Klasse* 1936, No.5.
- [10] Case FH. *J. Heterocyclic Chem.* 1971; 8: 1043-1046.
- [11] Ried W, Schomann P. *Liebigs Ann. Chem.* 1968; 714: 128-139.
- [12] Schmid K. *Reaktionen mehrfunktioneller Heterocyclen - Aufbau polyheterofunktioneller Heterocyclenkette*, University of Regensburg 1997.
- [13] Neunhoeffer H. 1,2,4-Triazines and their Benzo Derivatives, *Comprehensive Heterocyclic Chemistry II*, Katritzky AR, Rees CW, Scriven EFV, editors, Oxford: Pergamon Press, 1996; 6: 507-574.
- [14] Sauer J. 1,2,4,5-Tetrazines, *Comprehensive Heterocyclic Chemistry II*, Katritzky AR, Rees CW, Scriven EFV, editors, Oxford: Pergamon Press, 1996; 6: 901-957.