

A Short Synthesis of the Bicyclic Core of the Zaragozic Acids

Pierre Fraisse, Issam Hanna,* Jean-Yves Lallemand

Laboratoire de Synthèse Organique associé au CNRS, Ecole Polytechnique, F-91128 Palaiseau, France


Received 31 July 1998; accepted 17 August 1998

Abstract


A synthesis of the highly functionalized bicyclic core of the zaragozic acids is described, in which the key step involves the construction of the C4 quaternary carbon center by a CeCl_3 -promoted aldol reaction between the α -D-xylofuranuronic acid derivative **2** and D-(R)-glyceraldehyde acetonide. © 1998 Published by Elsevier Science Ltd. All rights reserved.

Keywords: Zaragozic Acids, aldol reaction, (R)-glyceraldehyde acetonide, α -D-xylofuranuronic acid.

The zaragozic acids (squalostatins), a family of naturally occurring fungal metabolites isolated in 1992, have been found to be potent inhibitors of the enzyme squalene synthase and thus of potential therapeutic use for the treatment of hypercholesterolaemia. Structurally, these compounds share a common densely functionalized 2,8-dioxabicyclo[3.2.1]octane skeleton **1**. The complexity of this structure combined with the biological activity of these natural products has served to stimulate a large amount of synthetic activity in a number of laboratories,¹ and culminated in the total syntheses of zaragozic acids A² and C.³




Zaragozic acids




In a new synthetic approach, we envisaged the construction of the bicyclic core of the zaragozic acids starting from the readily available α -D-xylofuranuronic acid derivative **2**⁴ in which the configuration at C2 and C3 correlates with the configuration at C6 and C7 of the target. As shown in the retrosynthetic analysis (Scheme 1), our plan involves the introduction of the three-carbon unit at C4 by aldol condensation of the enolate generated from **2** and D-(R)-glyceraldehyde acetonide **3**⁶. Assessment of the feasibility of quaternization at C4⁷ as well as internal ketalization would lead to a short alternative synthesis of **1**. The success of such an approach depends on

two crucial requirements: (a) The generation of an enolate from **2** and its reaction with a (D)-glyceraldehyde derivative bearing in mind that such a nucleophilic center which carries an adjacent oxygen is prone to epimerisation and β -elimination,⁸ and (b) the stereoselective introduction at C4 of the new C-C bond from the β -face of the enolate. We believed that the facial selectivity in the attack upon the aldehyde could be induced by the benzyloxy group at C3 α .




Scheme 1

Initial attempts to effect the aldol condensation of ester **2** with aldehyde **3** via the lithium, sodium or potassium enolates (using LDA, LiHMDS, NaHMDS or KHMDS) in THF at -100°C were disappointing. Under these conditions, the desired product **4** was isolated in only 0-10% yield. At higher temperature, β -elimination occurred followed by the degradation of the resulting product. With the zinc enolate, formed from the potassium enolate by metal exchange at -100°C , a higher yield of **4** was obtained (20-30%). After considerable experimentation,⁹ we found that addition of a precooled 1:1 mixture of cerium chloride and aldehyde **3** in THF to the solution of the potassium enolate in THF at -100°C gave **4** reproducibly in 65 to 70% yield. ^1H NMR analysis showed that **4** is a mixture of *three* out of the four possible diastereomers in 2.5:1.5:1 ratio. Dess-Martin oxidation¹⁰ of the mixture afforded *two* readily separable β -ketoesters **5**, $[\alpha]_{\text{D}} +10.8$ (c 1.45, CH_2Cl_2), and **6**¹¹ in a 4:1 ratio and in a combined yield of 88%. This established that the major diastereomers obtained from the aldol reaction possessed the same configuration at C4.




Scheme 2

Having introduced the C4 side chain onto the furanose ring, we next examined the ketalisation step. To this end, lithium aluminum hydride reduction of **4** afforded the expected diol which was protected as the dibenzyl ether **7** as a separable mixture of diastereomers. The major isomer was subjected to Dowex resin in refluxing acetonitrile for 20 h, and the crude product was treated with acetic anhydride in the presence of dimethylaminopyridine in dichloromethane to give a mixture of the desired ketals **8**¹¹ (57%) and its isomer **9** (19%). The structure and configuration of **8** were proven by extensive NMR experiments. In particular, the observed coupling constant between H6 and H7 ($J=2.6$ Hz) along with important cross-peaks in the NOESY spectrum between H3 and H6 provide strong arguments for the configuration depicted. Moreover, the observed J H3-H4 value of 10 Hz is in support with the *trans*-diaxial configuration in the chair conformation of the six membered ring.



As it has been already reported for such a system,¹² the ketalization step can lead to the undesired bicyclic ketal if the hydroxyl group at C4 (Squalestatins numbering) is unprotected. In fact, when compound **10**, readily prepared by reduction of **4** and subsequent stereoselective protection of the primary hydroxyl group using the stannylene procedure,¹³ was heated with HCl in MeOH-THF at 50°C for 24 h, the ketal **11**, mp 129-130°C; $[\alpha]_D^{25} +62$ (c 1.10, CHCl_3), was formed in 43% yield and 23% of the starting material was recovered. Like for compound **8**, the structure of ketal **11** was deduced on the basis of ^1H and ^{13}C NMR experiments.



In summary, we have achieved a short synthesis of the bicyclic core of zaragozic acids which features as the key step an aldol condensation of the readily available ester **2** with (*R*)-glyceraldehyde acetonide.

Aknowledgement: We thank Hoechst Marion Roussel and the CNRS for a grant (BDI to P. F.).

References and Notes

1. For a review of the chemistry and biology of the zaragozic acids (squalostatins) see: Nadin, A.; Nicolaou, K. C. *Angew. chem. Int. Ed. Engl.* **1996**, *35*, 1622-1656. For more recent approaches see: Hegde, S. G.; Myles, D. C. *Tetrahedron Let.* **1997**, *38*, 4329-4332. Xu, Y.; Johnson, C. R. *Tetrahedron Let.* **1997**, *38*, 1117-1120. Paterson, I.; Feßner, K.; Finaly, M. R. V.; Jacobs, M. F. *Tetrahedron Let.* **1997**, *38*, 4301-4304. Ito, H.; Matsumoto, M.; Yoshizawa, T.; Takao, K.; Kobayashi, S. *Tetrahedron Let.* **1997**, *38*, 9009-9012. Brogan, J. B.; Zercher, C. K. *Tetrahedron Let.* **1998**, *39*, 1691-1694. Kataoka, O.; Kitagaki, S.; Watanabe, N.; Kobayashi, J.; Nakamura, S.; Shiro, M.; Hashimoto, S. *Tetrahedron Let.* **1998**, *39*, 2371-2374.
2. Nicolaou, K. C.; Nadin, A.; Leresche, J. E.; Yue, E. W.; La Greca, S. *Angew. chem. Int. Ed. Engl.* **1994**, *33*, 2190-2191. Stoermer, D.; Caron, S.; Heathcock, C. H. *J. Org. Chem.* **1996**, *61*, 9115-9125. Caron, S.; Stoermer, D.; Mapp, A. K.; Heathcock, C. H. *J. Org. Chem.* **1996**, *61*, 9126-9134.
3. Evans, D. A.; Barrow, J. C.; Leighton, J. L.; Robichaud, A. J.; Sefkow, M. *J. Am. Chem. Soc.* **1994**, *116*, 12111-12112. Carreira, E. M.; Du Bois, J. *J. Am. Chem. Soc.* **1995**, *117*, 8106-8125. Sato, H.; Nakamura, S.; Watanabe, N.; Hashimoto, S. *Synlett*, **1997**, 451-454. Armstrong, A.; Jones, L. H.; Barsanti, P. A. *Tetrahedron Let.* **1998**, *39*, 3337-3340.
4. Compound **2** was prepared as a separable α : β (3:1) mixture of epimers at the anomeric center by refluxing the known 1,2-O-(1-methylethylidene)-3-O-(phenylmethyl)- α -D-xylofuranuronic acid⁵ in methanol with a catalytic amount of Amberlite IR ion exchange resin for 24 h, followed by the protection of the secondary alcohol as its methoxymethyl ether.
5. Dhavale, D. D.; Tagliavini, E.; Trombini, C.; Umani-Ronchi, A. *J. Org. Chem.* **1989**, *54*, 4100-4105.
6. Prepared from D-mannitol according to the literature procedure: Schmid, C.; Bryant, J. D. *Org. Synth.* **1995**, *72*, 6-13.
7. For the formation of carbon-carbon bonds at C4 of the furanose ring bearing an alkoxy group at C3 see: a) from an aldehyde at C4 *via* enamine and enol acetate derivatives: Secrist, J. A.; Winter, W. J. *J. Am. Chem. Soc.* **1978**, *100*, 2554-2555; by Cannizzaro reaction with formaldehyde: Jones, G. H.; Tanguchi, D. T.; Moffat, J. G. *J. Org. Chem.* **1979**, *44*, 1309-1317. Gurjar, M. K.; Das, S. K.; Sadalapure, K. S. *Tetrahedron Let.* **1995**, *36*, 1933-1936. b) from ester at C4 *via* the Claisen rearrangement of the enolate: Ireland, R. E.; Norbeck, D. W. *J. Am. Chem. Soc.* **1985**, *107*, 3279-3285. McVinish, L. M.; Rizzacasa, M. A. *Tetrahedron Let.* **1994**, *35*, 923-926.
8. Handa, S.; Tsang, R.; McPhail, A. T.; Fraser-Reid, B. *J. Org. Chem.* **1987**, *52*, 3489-3491 and references cited therein.
9. Attempts to achieve the aldol reaction *via* enol acetate, enamine or silyl ketene acetal derivatives of **2** were unsuccessful.
10. Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277-7287. Ireland, R. E.; Liu, L. *J. Org. Chem.* **1993**, *58*, 2899.
11. All new compounds are fully characterized by their spectroscopic and analytical data. **9**: $[\alpha]_D^{+32}$ (c 0.9, CHCl₃), IR ν_{\max} 2930, 2869, 1740, 1454, 1371, 1227, 1099, 1025 cm⁻¹. ¹H (CDCl₃) δ 7.40-7.15 (15 H, m, Harom), 5.57 (1 H, d, J = 4.5, H_I), 5.10 (1 H, dd, J = 2.6, J = 4.5, H₇), 4.78 (1 H, d, J = 11.8), 4.65 (1 H, d, J = 6.8), 4.62 (1 H, d, J = 7.6), 4.56 (1 H, d, J = 11.8), 4.50 (1 H, d, J = 7.6), 4.47 (1 H, d, J = 6.8), 4.27 (1 H, d, J = 2.6, H₆), 4.22 (1 H, dd, J = 2.2, J = 12.0, H_{3'a}), 4.02 (1 H, dd, J = 5.3, J = 12.0, H_{3'b}), 3.97 (1 H, d, J = 10.0, H₄), 3.92 (1 H, d, J = 11.9, H_{5'a}), 3.87 (1 H, d, J = 11.9, H_{5'b}), 3.78 (1 H, ddd, J = 2.2, J = 5.3, J = 10.0, H₃), 2.07 (3 H, s), 2.03 (3 H, s) ppm. ¹³C (CDCl₃) δ 170.9, 169.8 (C, C=O acetates), 138.3, 137.6, 137.9 (C, Bn), 128.4, 128.1, 128.0, 127.9, 127, 8, 127.6 (CH, Bn), 96.2 (CH, C_I), 85.7 (C, C₅), 81.0 et 80.7 (CH, C₆ et C₇), 74.0 et 73.8 (CH₂, C_{5'} et C_{3'}), 72.1 (CH), 71.8 (CH₂ Bn), 69.9 (CH), 68.1 and 63.7 (CH₂ Bn), 20.8 and 20.5 (CH₃, acetates) ppm.
12. See for examples: Paterson, I.; Feßner, K.; Finaly, M. R. V.; Jacobs, M. F. *Tetrahedron Let.* **1996**, *37*, 8803-8806. Xu, Y.; Johnson, C. R. *Tetrahedron Let.* **1997**, *38*, 1117-1120. Hegde, S. G.; Myles, D. C. *Tetrahedron* **1997**, *53*, 11179-11190. Armstrong, A.; Jones, L. H.; Barsanti, P. A. *Tetrahedron Let.* **1998**, *39*, 3337-3340.
13. For a review see: David, S.; Hanessian, S. *Tetrahedron*, **1985**, *41*, 643-663.