



Highly enantioselective synthesis of 1,3-mercapto alcohols from α,β -unsaturated ketones: asymmetric bifunctional group exchange reaction

Hiroaki Shiraki, Kiyoharu Nishide and Manabu Node *

Kyoto Pharmaceutical University, Misasagi, Yamashina, Kyoto 607-8414, Japan

Received 14 February 2000; accepted 03 March 2000


Abstract

Optically active 1,3-mercapto alcohols were synthesized from α,β -unsaturated ketones using a chiral reagent **B** and dimethylaluminum chloride in two steps. The transformation involved a tandem Michael addition–MPV reduction and a base-catalyzed elimination. The two newly created chiral carbons in *trans*-chalcone derivatives were enantioselectively controlled to a high degree. Using the above transformation, an asymmetric bifunctional group exchange reaction between the substrate and chiral reagent was developed. © 2000 Elsevier Science Ltd. All rights reserved.

The Meerwein–Ponndorf–Verley (MPV) reduction is a useful method for the chemoselective reduction of a ketone moiety among several types of carbonyl compounds.¹ Although many researchers² have tried to develop an asymmetric MPV reduction, high enantiomeric excess has been observed only in the intramolecular MPV reaction,³ with the exception of the catalytic reaction by Evans and co-workers.⁴ We have recently developed a tandem Michael addition–MPV reduction of α,β -unsaturated ketones using chiral mercapto alcohol **A** and Lewis acid. The resulting product **2A** was utilized in the asymmetric synthesis of secondary alcohols and allylic alcohols, as shown in equation (1) in Scheme 1.⁵ The stereochemistry of the two newly created chiral carbons in the sulfide **2A** is completely controlled by dynamic kinetic resolution via a reversible Michael addition. In order to utilize the two chiral carbons effectively, we designed a new method for asymmetric synthesis of 1,3-mercapto alcohols **3** from α,β -unsaturated ketones **1** using a new chiral reagent **B**, as shown equation (2) in Scheme 1.

Chiral reagent **A** is not adequate for the strategy of the above asymmetric synthesis, because the transformation from the sulfide **2A** into the thiol **3** is not easy to carry out. Therefore, we designed a new type of chiral reagent which has a hydrogen atom on the carbon atom between the two carbon atoms bearing the thiol and hydroxyl groups, as shown in the structure **B**. The sulfides **2B** utilizing a chiral reagent **B** should be easily converted with the aid of an amine base into the 1,3-mercapto alcohols **3** and


* Corresponding author.



Scheme 1.

the α,β -unsaturated ketone **4**, which is the starting material used in the preparation of the reagent **B**. The product **3** should have a high de and ee if the sulfide **2B** is produced under dynamic kinetic resolution conditions.⁵ The above transformation corresponds to a novel exchange reaction of bifunctional groups between the substrate and the reagent.

The exchange reaction using methyl vinyl ketone and a known chiral reagent **B₁**⁶ prepared from (–)-pulegone was carried out first (Scheme 2). The sulfide **2B₁** obtained by the tandem reaction was subjected to the elimination reaction using DBU in refluxing toluene for 24 h to give the desired 1,3-mercapto alcohol **3a** (90% ee) and (–)-pulegone (**4B₁**) in 70 and 79% yields, respectively. Because of the low chemical yield (38%) of **2B₁**, we further investigated the exchange reaction using a new chiral reagent **B₂**⁷ (98% ee), which was prepared from (+)-camphor. The tandem reactions of mesityl oxide **1b** and *trans*-chalcone **1c** with the reagent **B₂** afforded the sulfides **2b** and **2c** in 94 and 60% yields, respectively. These sulfides were next converted by base-catalyzed elimination to the 1,3-mercapto alcohols **3b** (74% yield, 82% ee) and **3c** (72% yield, 96% ee) together with the α,β -unsaturated ketone **4B₂** ($\leq 76\%$ yield). The mercapto alcohol **3c**, having two chiral carbons, showed an especially high optical purity and high *syn-anti* selectivity.⁸ Reagent **B₂** was more effective compared to reagent **B₁**, although the chemical yield was lower than that in the reaction using **A**, even given the fact that 2 equivalents of the substrate were used compared to 1 equivalent of the chiral reagent.



Scheme 2.

Although a novel asymmetric exchange reaction of bifunctional groups was demonstrated by the above reactions, nevertheless we attempted to improve the chemical yield in the base-catalyzed elimination. Since the elimination reaction is an equilibrium reaction, it is expected that the addition of a thiol would shift the equilibrium to the right. Actually, the thiol exchange reaction of **2c** proceeded smoothly using benzyl mercaptan in refluxing toluene to give the 1,3-mercapto alcohol **3c** and the sulfide **5** in 97 and 91% yields, respectively. Generalization of the asymmetric synthesis of 1,3-mercapto alcohols under the latter elimination conditions was tried. The results of the tandem reaction of α,β -unsaturated ketones using the chiral reagent **B**₂ and the base-catalyzed elimination of the reaction product are summarized in Table 1.

Table 1
Asymmetric synthesis of 1,3-mercaptoalcohol from α,β -unsaturated ketone⁹

Reaction scheme showing the conversion of an α,β -unsaturated ketone **1** to a sulfide **2** using B_2 (98% ee) and Me_2AlCl (1.2 eq.) in benzene at room temperature (r.t.). Sulfide **2** is then converted to a mixture of *anti*-**3** and *syn*-**3** using DBU (2 eq.) and BnSH in toluene under reflux.

Entry	Substrate 1				2		3			
		R ¹	R ²	R ³	time(h)	yield (%) ^a	time(h)	yield (%) ^a	anti / syn	ee (%) ^b
1	a	H	H	CH ₃	76	50	24	92 ^c	—	98 ^d
2	b	CH ₃	CH ₃	CH ₃	21	94	14	94	—	82 ^d
3	c	Ph	H	Ph	48	60	18	97	100/0	96 ^e
4	d	<i>p</i> -ClPh	H	Ph	43	56	22	93	100/0	93 ^e
5	e	<i>p</i> -MeOPh	H	Ph	48	56	18	99	100/0	94 ^e
6	f	<i>p</i> -MePh	H	Ph	44	63	18	92	100/0	96 ^e
7	g	Ph	H	CH ₃	21	74	21	96	36/64	95 ^e
8	h	Ph	H	<i>n</i> -C ₆ H ₁₇	20	50	17	97	47/53	98 ^e

a) Isolated yield b) Enantiomeric excesses of the alcoholic carbon; determined by HPLC analyses using Daicel CHIRALCEL OB, OD or CHIRALPAK AS.


c) The elimination was carried out after benzylation of the hydroxyl group of **2**.

d) Determined after desulfurization of the benzoate of **2** with Raney Ni (W2) in EtOH.

e) Determined after desulfurization of **2** with Raney Ni (W2)-NaPH₂O₂-acetate buffer-EtOH.¹⁰

Although the tandem reaction of the α,β -unsaturated ketone **1** to give the sulfide **2** proceeded in moderate yield, the base-catalyzed reaction of **2** with benzyl mercaptan resulted in high yields to give the 1,3-mercapto alcohols **3** with excellent ees,¹¹ except for **3b**. The relatively low ee of **3b** compared with **3a** might be attributed to the restriction in the transition state of the MPV reduction due to the bulkiness of the two methyl substituents on the β -carbon. In the reaction of the *trans*-chalcone derivatives (**1c–f**), two chiral carbons were created to give optically active 1,3-mercapto alcohols **3c–f** as a single diastereomer. The controlling factor for this exceedingly high diastereoselectivity⁸ for *anti* to *syn* must be dynamic kinetic resolution. On the other hand, the tandem reactions of benzalacetone **1g** and its derivative **1h** having an alkyl group at R³ afforded a diastereomeric mixture of the sulfides **2**, which were converted to the optically active alcohols **3g** and **3h** as *anti* and *syn* mixtures. This low diastereoselectivity suggests that the tandem reaction of **1g** and **1h** proceeded without control by dynamic kinetic resolution. It is presumed that the above difference between *trans*-chalcone and benzalacetone is due to the magnitude of a 1,3-diaxial like interaction of R¹ and R² toward R³ in the transition states of MPV reduction, as shown in Scheme 3. The MPV reduction of the diastereomer having a large steric interaction between the two phenyl groups does not take place and the other diastereomer is subject to the MPV reduction because of

a small steric interaction, i.e. that the reaction undergoes control by dynamic kinetic resolution. There are many papers on asymmetric Michael addition of a thiol to an α,β -unsaturated system;¹² however, there is no precedent for a synthetic method corresponding to the asymmetric Michael addition of hydrogen sulfide like the one observed in the reaction of the *trans*-chalcones.



Scheme 3. Plausible transition states for sulfide **2**

In conclusion, we have developed the first method for an asymmetric synthesis of the 1,3-mercapto alcohols **3** from the α,β -unsaturated ketones **1** through the tandem Michael addition–MPV reduction followed by elimination. An important feature of this method is that the α,β -unsaturated ketone **4** or the sulfide **5** obtained together with **3** is able to be recycled for the preparation of the chiral reagent **B**.⁷ This is characteristic of an asymmetric bifunctional group exchange reaction.

Acknowledgements

Partial financial support (Grant-in-Aid No. 09470489) for this research by the Ministry of Education, Science and Culture, Japan, is gratefully acknowledged.

References

- For reviews, see: (a) Wilds, A. L. *Org. React.* **1944**, 2, 178–223. (b) de Graauw, C. F.; Peters, J. A.; van Bakkum, H.; Huskens, J. *Synthesis* **1994**, 1007–1017. (c) Takahashi, K.; Shibagaki, M.; Kuno, H.; Matsushita, H. *Shokubai* **1995**, 37, 23–27, and references cited therein.
- Chiral Grignard reagent: (a) Streitwieser Jr., A.; Wolfe, J. R.; Schaeffer, W. D. *Tetrahedron* **1959**, 6, 338–344. (b) Foley, W. M.; Welch, F. J.; La Combe, E. M.; Mosher, H. S. *J. Am. Chem. Soc.* **1959**, 81, 2779–2784. (c) MacLeod, R.; Welch, F. J.; Mosher, H. S. *J. Am. Chem. Soc.* **1960**, 82, 876–880. (d) Burrows, E. P.; Welch, F. J.; Mosher, H. S. *J. Am. Chem. Soc.* **1960**, 82, 880–885. (e) Birtwistle, J. S.; Lee, K.; Morrison, J. D.; Sanderson, W. A.; Mosher, H. S. *J. Org. Chem.* **1964**, 29, 37–40. Chiral Aluminum alkoxide: (f) Doering, W. von E.; Young, R. W. *J. Am. Chem. Soc.* **1950**, 72, 631. (g) Jackman, L. M.; Mills, J. A.; Shannon, J. S. *J. Am. Chem. Soc.* **1950**, 72, 4814–4815. (h) Newman, P.; Rutkin, P.; Mislow, K. *J. Am. Chem. Soc.* **1958**, 80, 465–473. (i) Nasipuri, D.; Sarker, G. *J. Indian Chem. Soc.* **1967**, 44, 165–166. (j) Nasipuri, D.; Sarker, G.; Ghosh C. K. *Tetrahedron Lett.* **1967**, 5189–5192. (k) Fles, D.; Majhofer, B.; Kovac, M. *Tetrahedron* **1968**, 24, 3053–3057.
- (a) Zhou, W. S.; Zhou, X. M.; Ni, Y. *Acta Chimica Sinica* **1984**, 42, 706–709. (b) Zhou, W. S.; Wang, Z. Q.; Zhang, H.; Fei, R. Y.; Zhuang, Z. P. *Acta Chimica Sinica* **1985**, 43, 168–173. (c) Ishihara, K.; Hanaki, N.; Yamamoto, H. *J. Am. Chem. Soc.* **1991**, 113, 7074–7075. (d) Ishihara, K.; Hanaki, N.; Yamamoto, H. *Synlett* **1993**, 127–129. (e) Molander, G. A.; McKie, J. A. *J. Am. Chem. Soc.* **1993**, 115, 5821–5822. (f) Baramée, A.; Chaichit, N.; Intawee, P.; Thebtaranonth, C.; Thebtaranonth, Y. *J. Chem. Soc., Chem. Commun.* **1991**, 1016–1017. (g) Fujita, M.; Takarada, Y.; Sugimura, T.; Tai, A. *J. Chem. Soc., Chem. Commun.* **1997**, 1631–1632.
- Evans, D. A.; Nelson, S. G.; Gagne, M. R.; Muci, A. R. *J. Am. Chem. Soc.* **1993**, 115, 9800–9801.
- (a) Nishide, K.; Shigeta, Y.; Obata, K.; Node, M. *J. Am. Chem. Soc.* **1996**, 118, 13103–13104. (b) Node, M.; Nishide, K.; Shigeta, Y.; Shiraki, H.; Obata, K. *ibid.* **2000**, 122, 1927–1936.
- (a) Eliel, E. L.; Lynch, J. E. *Tetrahedron Lett.* **1981**, 22, 2855–2858. (b) Lynch, J. E.; Eliel, E. L. *J. Am. Chem. Soc.* **1984**, 106, 2943–2948. (c) Eliel, E. L.; Lynch, J. E.; Kume, F.; Frye, S. V. *Org. Synth.* **1987**, 65, 214–223.

7. Chiral reagent **B**₂ was prepared as follows: (1) The *exo* methylenation of (+)-camphor with phenyllithium and paraformaldehyde gave **4B**₂ in 89% yield; (2) the Michael addition of benzyl mercaptan of **4B**₂ gave **5** in 82% yield; (3) debenzylation with sodium metal in liquid ammonia after sodium borohydride reduction afforded **B**₂ in 17% yield, along with the other diastereomers.
8. The absolute configurations of the thiolic carbon in **3c–f** were determined by NOE experiments, because a 1,3-mercapto alcohol formed a chair conformation of a six-membered ring by intramolecular hydrogen bonding between the oxygen atom and the hydrogen atom of the thiol.
9. A typical procedure: (Entry 3 in Table 1) To a benzene solution (4 ml) of the mercapto alcohol **B**₂ (37 mg, 0.18 mmol) was added dropwise dimethylaluminum chloride (1.0 M hexane solution, 0.21 ml, 0.22 mmol). After the mixture was stirred for 15 min at room temperature, a benzene solution (2 ml) of the α,β -unsaturated ketone **1c** (75 mg, 0.36 mmol) was added, and the resulting mixture was stirred for 48 h at the same temperature. The reaction mixture was quenched with a 1N-HCl solution, and the aqueous layer extracted with chloroform. After the usual work-up, the product **2c** (36 mg, 60%) was isolated by silica gel chromatography. A mixture of **2c** (28 mg, 0.07 mmol), DBU (0.047 ml, 0.34 mmol), and benzyl mercaptan (15 μ l, 0.13 mmol) in toluene (8 ml) was refluxed for 18 h. The usual work-up and purification by silica gel chromatography gave the 1,3-mercapto alcohol **3c** (16 mg, 97% yield), along with **5** (18 mg, 91% yield).
10. (a) Nishide, K.; Shigeta, Y.; Obata, K.; Inoue, T.; Node, M. *Tetrahedron Lett.* **1996**, 37, 2271–2274. (b) Node, M.; Nishide, K.; Shigeta, Y.; Obata, K.; Shiraki, H.; Kunishige, H. *Tetrahedron* **1997**, 53, 12883–12894.
11. The absolute configuration of the alcoholic carbon in **3** was determined by comparison of the sign of the specific rotation with that in the literature (Holt, D. A. et al. *J. Am. Chem. Soc.* **1993**, 115, 9925–9938, and Ref. 5).
12. Asymmetric Michael addition of a thiol using an auxiliary: (a) Nagao, Y.; Kumagai, T.; Yamada, S.; Fujita, E.; Inoue, Y.; Nagase, Y.; Aoyagi, S.; Abe, T. *J. Chem. Soc., Perkin Trans. 1* **1985**, 2361–2367. (b) Wu, M.-J.; Wu, C.-C.; Tseng, T.-C.; Pridgen, L. N. *J. Org. Chem.* **1994**, 59, 7188–7189. (c) Tomioka, K.; Muraoka, A.; Kanai, M. *J. Org. Chem.* **1995**, 60, 6188–6190. Catalytic asymmetric Michael addition of a thiol: (d) Hiemstra, H.; Wynberg, H. *J. Am. Chem. Soc.* **1981**, 103, 417–430. (e) Yamashita, H.; Mukaiyama, T. *Chem. Lett.* **1985**, 363–366. (f) Sera, A.; Takagi, K.; Katayama, H.; Yamada, H.; Matsumoto, K. *J. Org. Chem.* **1988**, 53, 1157–1161. (g) Nishimura, K.; Ono, M.; Nagaoka, Y.; Tomioka, K. *J. Am. Chem. Soc.* **1997**, 119, 12974–12975. (h) Tomioka, K.; Okuda, M.; Nishimura, K.; Manabe, S.; Kanai, M.; Nagaoka, Y.; Koga, K. *Tetrahedron Lett.* **1998**, 39, 2141–2144. (i) Kanemasa, S.; Oderaotoshi, Y.; Wada, E. *J. Am. Chem. Soc.* **1999**, 121, 8675–8676. (j) Emori, E.; Arai, T.; Sasai, H.; Shibasaki, M. *J. Am. Chem. Soc.* **1998**, 120, 4043–4044.