



Corrigendum

**Corrigendum to “A stopperless tetrathiafulvalene based
[2]pseudorotaxane molecular shuttle”
[Tetrahedron Letters 42 (2001) 4223–4226][†]**

Martin R. Bryce,^a Graeme Cooke,^{b,*} Wayne Devonport,^a Florence M. A. Duclairoir^b and
Vincent M. Rotello^c

^aDepartment of Chemistry, University of Durham, South Road, Durham DH1 3LE, UK

^bDepartment of Chemistry, Heriot-Watt University, Riccarton, Edinburgh EH14 4AS, UK

^cDepartment of Chemistry, University of Massachusetts at Amherst, Amherst, MA 01002, USA

In this article we stated that we had produced a pseudorotaxane based molecular shuttle (Fig. 1a), in which the electron deficient macrocycle cyclobis-(paraquat-*p*-phenylene) moves intramolecularly between two degenerate tetrathiafulvalene units of a stopperless axle. After discussions with Professors Leigh and Stoddart, it seems more likely that the macrocycle passes from one end of the axle to the other by first dethreading from the pseudorotaxane, followed by either an intermolecular recomplexation with a vacant TTF unit (Fig. 1b). Therefore, as the interconversion between occupied and unoccupied stations probably occurs principally via an

intermolecular rather than an intramolecular route, the term shuttle is inappropriate.

It is the authors' intention to clarify to what extent shuttling and/or a decomplexation/recomplexation is occurring by obtaining thermodynamic and kinetic data for our original and analogous systems, and the results of this investigation will be investigated in due course.

We thank Professors David Leigh and Fraser Stoddart for bringing this misinterpretation to our attention.




Figure 1. Shuttling versus a decomplexation/recomplexation process.

* Corresponding author. Tel.: 0131 451 8184; fax: 0131 451 3180; e-mail: g.cooke@hw.ac.uk

[†] PII of original article: S0040-4039(01)00627-X