



Corrigendum

Corrigendum to “Asymmetric allylation of aldehydes with allyltrichlorosilane promoted by chiral sulfoxides” [Tetrahedron Lett. 44 (2003) 7179][☆]

Antonio Massa,^{a,*} Andrei V. Malkov,^b Pavel Kočovský^b and Arrigo Scettri^a

^a*Dipartimento di Chimica, Università di Salerno, Via S. Allende 84081 Baronissi, Salerno, Italy*

^b*Department of Chemistry, University of Glasgow, Glasgow G12 8QQ, UK*

Table 2 on page 7180 of the above paper contained errors. The corrected version is printed below.

Table 2. Allylation of aldehydes **1** using enantiopure (*R*)-methyl-*p*-tolylsulfoxide **4a**

Entry	1 (R ¹)	4a (equiv.)	Time (h)	Yield (%) ^a	% ee of (<i>S</i>)- 3 ^b
1 ^c	1a (Ph)	1.5	65	70	51 ^d
2 ^c	1a (Ph)	3.0	7	62	55 ^d
3 ^c	1b (2-furyl)	3.0	7	88	42 ^f
4 ^c	1c (PhCH=CH)	3.0	7	79	46 ^d

^a See note ‘a’ in Table 1.

^b The absolute configuration of **3** was found to be (*S*) according to the optical rotation reported in the Refs. 5 and 6.

^c The sulfoxide was recovered in 85% yield and 85% e.e.

^d Determined by HPLC analysis employing a Daicel Chiralcel OB column.

^e The sulfoxide was recovered in >95% yield and 97% e.e.

^f Determined by HPLC analysis of the corresponding 3,5-dinitrobenzoate (Daicel Chiralcel AD column).

[☆] DOI of original article: 10.1016/S0040-4039(03)01801-X

* Corresponding author. Tel.: +39 089 965370; fax: +39 089 965296; e-mail: amassa@unisa.it