

hypochlorite and N-chlorosuccinimide. Whilst the latter gave a quantitative yield of 2,2'-dichlorodiphenylamine-4,4'-dicarboxylic acid (VII) and by subsequent decarboxylation the known, 2,2'-dichlorodiphenylamine (I, 80% yield),³ compound IV yielded 95% of VII and 5% of IV which could not be separated; but their decarboxylation products I and VI were separated by column chromatography on alumina.

The structure of compound I was confirmed by its NMR spectrum which is typical of an *ortho*-disubstituted benzene.

A second route to the system VIII which lends itself to large-scale preparations, starts from the easily accessible⁴ 4,4'-dibromo-N-methyldiphenylamine. Its chlorination with *t*-butyl hypochlorite gave 4,4'-dibromo-2,2'-dichloro-N-methyldiphenylamine (IX), which could be selectively debrominated to VIII either with triphenylstannan⁵ or by magnesium in the presence of ethylene dibromide.⁶ The latter method gives a more easily purifiable product.

The NMR spectrum of VIII is sufficiently similar to that of I to make formula VIII unequivocal.

EXPERIMENTAL

Dimethyl diphenylamine-4,4'-dicarboxylate (II, R = CH₃)

Diphenylamine-4,4'-dicarboxylic acid¹ was stirred with excess SOCl₂ for 4 hr at room temp and for 2 hr at the boiling temp of SOCl₂. Excess of the latter was distilled off *in vacuo* and the residue dissolved in cold MeOH and refluxed for 2 hr. The reaction mixture was poured into water and the solid product filtered and recrystallized from aqueous EtOH as colorless crystals, m.p. 177° (lit.² 177°); yield, 95%.

2,2'-Dibromodiphenylamine-4,4'-dicarboxylic acid (III, R = H)

Bromine (5 g) in 50 ml AcOH was added with stirring at 5° to a soln of 8 g of II (R = H) in 100 ml of the same solvent. The mixture was kept at this temp for 2 hr and poured into water. The solid product was dissolved in 10% NaHCO₃ aq, refluxed with active carbon, filtered and isolated by acidification, giving from aqueous alcohol, m.p. above 300° a quantitative yield. (Found: C, 41.0; H, 2.6; N, 3.4; Br, 38.2. Calc. for C₁₄H₉Br₂NO₄: C, 40.5; H, 2.2; N, 3.4; Br, 38.6%).

The *dimethyl ester* (III, R = CH₃) was obtained analogously from II (R = CH₃) in 89% yield, from aqueous alcohol, m.p. 219°. (Found: C, 43.6; H, 3.1; N, 3.0; Br, 35.7. Calc. for C₁₆H₁₃Br₂NO₄: C, 43.3; H, 3.1; N, 3.2; Br, 36.1%).

N 2,2',6,6'-Pentachlorodiphenylamine-4,4'-dicarboxylic acid (IV)

At a temp not exceeding 10°, a current of chlorine was passed through a soln of 5 g of II (R = H) in glacial AcOH until the absorption of the gas ceased. After 12 hr at room temp, the soln was poured into water and the ppt filtered off and purified from aqueous alcohol, m.p. > 300°; yield, 90%. (Found: C, 38.8; H, 1.8; N, 3.2; Cl, 40.5. Calc. for C₁₄H₆Cl₅NO₄: C, 38.8; H, 1.8; N, 3.2; Cl, 40.9%).

2,2',6,6'-Tetrachlorodiphenylamine-4,4'-dicarboxylic acid (V)

A mixture of 4.5 g of IV, 2 g NaI, 100 ml AcOH and 60 ml benzene was kept at room temp for 24 hr, poured into water and treated with NaHSO₃ aq. The solid product, resulting from the evaporation of the benzene layer, was precipitated from bicarbonate soln with acid and melted above 300°, yield, 90%. To obtain an analytically pure product, trituration with aqueous alcohol proved sufficient. (Found: C, 41.9; H, 2.0; N, 3.5; Cl, 36.0. Calc. for C₁₄H₇Cl₄NO₄: C, 42.5; H, 1.8; N, 3.5; Cl, 35.9%).

2,2',6,6'-Tetrachlorodiphenylamine (VI)

A soln of 4 g of IV or V in 100 ml freshly distilled quinoline was refluxed with 0.5 g copper bronze until the evolution of CO₂ ceased. The mixture was poured into water, the biphasic liquid filtered from the copper, and the aqueous layer extracted with benzene. The benzene extract was then washed twice with 5% HCl, dried and concentrated *in vacuo*, and the residue dissolved in a little fresh benzene and chromatographed on alumina. A mixture of hexane-benzene (2:1) served as eluent. Yield, 70–75%; m.p. 94°. (Found: C, 46.6; H, 2.5; N, 4.3; Cl, 47.0. Calc. for C₁₂H₇Cl₄N: C, 46.9; H, 2.3; N, 4.5; Cl, 46.3%).

2,2'-Dichlorodiphenylamine-4,4'-dicarboxylic acid (VII)

(a) A soln of 4.5 g t-butyl hypochlorite in 50 ml glacial AcOH⁷ was added, at 5° with stirring, to 5 g of II (R = H), dissolved in 100 ml of the same solvent. After a further 6 hr at 5°, the mixture was poured into water, and the product was filtered, yield, 100%. The analysis showed that the product contained a small amount of V.

(b) Similarly, 5 g of II (R = H) was treated with 5 g N-chlorosuccinimide. The mixture was poured into water, heated at 50° and filtered. The product was recrystallized repeatedly from aqueous alcohol (active carbon) and melted above 300°, yield, quantitative. (Found: C, 51.0; H, 2.6; N, 4.0; Cl, 22.5. Calc. for C₁₄H₉Cl₂NO₄: C, 51.5; H, 2.7; N, 4.3; Cl, 22.0%).

2,2'-Dichlorodiphenylamine (I)

Decarboxylation of VII in the manner described above gave an oil which was chromatographed on alumina and eluted with hexane, m.p. 32–33° (lit.³: 32°); yield, 80%.

4,4'-Dibromo-N-methyldiphenylamine

In the manner described for diphenylamine,³ N-methyldiphenylamine was brominated and the product (yield, quantitative), recrystallized from EtOH; m.p. 120°. (Found: C, 45.2; H, 3.4; N, 4.1; Br, 48.0. Calc. for C₁₃H₁₁Br₂N: C, 45.7; H, 3.2; N, 4.1; Br, 47.0%).

4,4'-Dibromo-2,2'-dichloro-N-methyldiphenylamine (IX)

At 5°, a soln of 4.5 g t-butyl hypochlorite in 50 ml AcOH was added slowly to a soln of 7 g of the foregoing compound in 150 ml of the same solvent. After 12 hr at room temp, the mixture was diluted with water, and the solid which precipitated was recrystallized from alcohol; m.p. 103°; yield, 90%. (Found: C, 37.9; H, 2.2; N, 3.0. Calc. for C₁₃H₉Cl₂Br₂N: C, 38.0; H, 2.2; N, 3.1%).

2,2'-Dichloro-N-methyldiphenylamine (VIII)

(a) When a mixture of 8 g of IX and 10 g triphenylstannan was heated slowly to 140°, an exothermic reaction caused the temp to rise to 190°. The heating was interrupted, and the temp maintained at 150° for 1.5 hr. The mass was refluxed with 50 ml MeOH for 30 min and after addition of 6 g MeI for another 10 min. Then 5 g KOH was added and the heating continued for 10 min. The product was diluted with water and extracted with benzene. The residue of this extract was a mixture of solid and a liquid which was taken up in MeOH. Distillation under 0.5 mm press gave at 170–175° an oil (yield, 70%) which was separated from some triphenylstannol by chromatography on alumina and eventually melted at 40°. (Found: C, 61.9; H, 3.8; N, 5.5; Cl, 28.0. Calc. for C₁₃H₁₁Cl₂N: C, 62.0; H, 4.4; N, 5.6; Cl, 28.1%).

(b) To a suspension of 11.0 g Mg in 200 ml ether, 4 g of 1,2-dibromoethane was added at such a rate that the heat of reaction kept the ether boiling. In the same way a soln of 8 g of IX and 4 g dibromoethane in 200 ml ether was added gradually. The reaction mixture was refluxed for 1 hr and decomposed with 10% HCl, and the ethereal layer was dried and concentrated, b.p. 170–180° (0.5 mm). The product, m.p. 42°, was purified by chromatography, as indicated above.

REFERENCES

- ¹ A. I. Kizber, *Chem. Abstr.* **50**, 207 (1956).
- ² A. I. Kizber and V. A. Puchov, *Chem. Abstr.* **52**, 6131 (1958).
- ³ A. W. Chapman and C. H. Perrott, *J. Chem. Soc.* 1777 (1932).
- ⁴ K. Fries, *Liebigs Ann.* **346**, 213 (1906).
- ⁵ L. A. Rothman and E. I. Becker, *J. Org. Chem.* **25**, 2203 (1960).
- ⁶ D. E. Pearson, D. Cowan and J. D. Beckler, *Ibid.* **24**, 504 (1959).
- ⁷ ^a B. F. Clark, *Chem. News* **143**, 265 (1931);
^b D. R. Harvey and R. O. C. Norman, *J. Chem. Soc.* 3604 (1961).