



Organotin Perchlorates as Gentle Lewis Acid Catalysts in Mukaiyama Reaction

Jian-xie Chen and Junzo Otera*

Department of Applied Chemistry, Okayama University of Science, Ridai-cho, Okayama 700, Japan

Abstract: Organotin perchlorates catalyze Mukaiyama reaction of ketene silyl acetal in highly chemoselective but uncommon manners. The competition reaction between aldehyde and acetal leads to exclusive formation of the aldehyde aldols leaving the acetal counterpart intact, an unusual outcome in terms of reaction under acidic conditions. α -Enals react with ketene silyl acetal in preference to the corresponding alkanal. In the competition between electronically different aldehydes, an electron-donating group increases the reactivity of aldehyde while the reverse is true with an electron-withdrawing group. These are opposite to the reactivity order in nucleophilic addition to free carbonyls. In contrast to ketene silyl acetal, enol silyl ethers derived from ketones are not activated by organotin perchlorates. Thus, these two enol silyl ethers can be discriminated from each other. A disilyl enol ether derived from a keto ester undergoes the electrophilic attack by aldehyde and α -enone exclusively on the ester function. The catalytic activities of TBSClO₄ which should be formed if organotin perchlorates underwent transmetallation with ketene silyl acetal or silyl ether of the aldolate, are totally different from those of organotin perchlorates, indicating that the organotin species work as real active species. The reaction is interpreted in terms of the S_N2 mechanism where the initial coordination of carbonyl group with organotin perchlorates plays a key role. The remarkable selectivities are ascribed to the weak acidity of the catalysts. The reactivities of the complexed carbonyls are completely different from those of free carbonyls. © 1997 Elsevier Science Ltd.


INTRODUCTION

Organotin compounds exhibit Lewis acidity¹ but, in general, it is not strong enough to trigger synthetically useful reactions.² However, if their acidity is increased they would serve as versatile catalysts which work under mild conditions due to their stability in the open atmosphere and easiness to handle. We have been working on organotin chemistry along this line. One of such strategies is to make use of associated organotin templates involving multi-reaction sites.³ A substrate and a reagent are activated by coordination with different tin atoms. These tin atoms are located closely to each other so that the substrate and reagent could interact directly on the tin template to facilitate the reaction through entropy gain. Another way for this purpose is to attach electronegative groups on the tin atom. We disclosed that incorporation of the triflate groups on tin led to a unique Lewis acid which enabled various types of differentiation between carbonyls and acetals.⁴ More recently, attachment of the perfluorophenyl group also was found to successfully increase the acidity.⁵ The present study stemmed from the expectation that the highly electron-withdrawing perchlorate group would create new organotin Lewis acid catalysts of synthetic promise. This is indeed the case. We report here that organotin perchlorates induce uncommon chemoselectivities in aldol and Michael reactions of enol silyl ethers (Mukaiyama reaction).


RESULTS AND DISCUSSION

Mukaiyama-Aldol Reaction of Ketene Silyl Acetal. We employed two organotin perchlorates, $\text{Bu}_3\text{SnClO}_4$ (**1a**) and $\text{Bu}_2\text{Sn}(\text{ClO}_4)_2$ (**1b**), in this study. These compounds were most conveniently prepared according to Lambert's method.⁶ Bu_3SnH or Bu_2SnH_2 was treated with 1 equiv. or 2 equiv. of Ph_3CClO_4 in dichloromethane and the resulting solution was used in situ for the Mukaiyama reaction. The solution contains

Table 1. Organotin Perchlorate-Catalyzed Mukaiyama-Aldol Reaction of Ketene Silyl Acetal.



entry	1	2	3	yield (%) of 4
1	1a	2a	3a	4aa 89
2	1b	2a	3a	4aa 92
3	1a	2a	3b	4ab 80
4	1b	2a	3b	4ab 89
5	1a	2a	3c	4ac 82
6	1a	2a	3d	4ad 86
7	1a	2b	3a	4ba 45
8	1a	2c	3a	4ca 52
9	1a	2d	3a	4da 90
10	1b	2d	3a	4da 91
11	1a	2d	3b	4db 81
12	1b	2d	3b	4db 87
13	1a	2e	3a	4ea 85
14	1b	2e	3a	4ea 86
15	1a	2e	3b	4eb 69
16	1b	2e	3b	4eb 73
17	1a	2e	3c	4ec 83
18	1a	2e	3d	4ed 60
19	1a	2f	3a	4fa 88




PhCHO (**3a**) $n\text{-C}_7\text{H}_{15}\text{CHO}$ (**3b**) $4\text{-MeOC}_6\text{H}_4\text{CHO}$ (**3c**) $4\text{-NCC}_6\text{H}_4\text{CHO}$ (**3d**)

Ph₃CH but this has no influence on the reaction. This is the safest way to arrive at **1** because the perchlorates are always manipulated in solution. We have no data on physical and chemical properties of the solid organotin perchlorates although metal perchlorates are explosive on some occasions. Anyway, we have encountered no troubles so far in the present method.




As expected, both **1a** and **1b** catalyzed reaction of ketene silyl acetals **2** with aldehydes **3** (Table 1); no significant difference was observed between these two catalysts. Unsubstituted, monosubstituted, and disubstituted ketene silyl acetals reacted quite smoothly, and both aromatic and aliphatic aldehydes were employable.

When α -enals were used as substrates, 1,2- and 1,4-additions competed (Table 2). Unsubstituted ketene silyl acetal **2a** favored the 1,2-addition (entries 1-3) while the 1,4-addition predominated with disubstituted one **2e** (entries 11,12). Virtually, no selectivity was observed with monosubstituted reagent **2d** (entries 6-8). By contrast, α -enone **6** always underwent exclusive 1,4-addition irrespective of nucleophiles (entries 4,5,9,10,13).

Table 2. Organotin Perchlorate-Catalyzed Mukaiyama Reaction of α -Enal and -Enone.



entry	1	2	5 or 6	Yield (%)		
				7	8	7:8
1	1a	2a	5a	7aa 81	8aa 6	93:7
2	1b	2a	5a	7aa 84	8aa 10	89:11
3	1a	2a	5b	7ab 83	0	100:0
4	1a	2a	6	0	8ac 78	0:100
5	1b	2a	6	0	8ac 66	0:100
6	1a	2d	5a	7da 38	8da 47	45:55
7	1b	2d	5a	7da 36	8da 47	43:57
8	1a	2d	5b	7db 45	8db 48	48:52
9	1a	2d	6	0	8dc 77	0:100
10	1b	2d	6	0	8dc 77	0:100
11	1a	2e	5a	7ea 17	8ea 73	19:81
12	1a	2e	5b	7eb 19	8eb 70	21:79
13	1a	2e	6	0	8ec 40	0:100



Competition between Aldehyde and Acetal. Although it is rather difficult to differentiate between the carbonyl functions and their acetals in Lewis acid-promoted reactions, there appeared some selective reactions. Trimethylsilyl triflate induced the Mukaiyama-aldol reaction of enol silyl ethers with acetals but not with carbonyls⁷ and alkyltitanium chlorides reacted with an acetal in the presence of a ketone.⁸ These results are easy to explain in terms of the facile formation of the oxocarbenium intermediate from acetal under acidic conditions. The reversal of this chemoselectivity is of synthetic interest. Previously, we disclosed unique chemoselectivities in Mukaiyama-aldol reaction by dibutyltin bis(triflate) (DBTT).^{4a} Smooth reaction occurred with aldehydes and ketals while ketones and acetals of aldehydes failed to react under the same reaction conditions. DBTT also differentiated various carbonyls and acetals in dithiane and dithiolane synthesis from thiostannanes.⁹ Mukaiyama et al. also reported the reversed selectivity in Mukaiyama-aldol reaction catalyzed by TBSCl-InCl₃.¹⁰ Now, we have found that **1** give rise to the perfect preference for aldehydes over acetals in Mukaiyama-aldol reaction of ketene silyl acetals (Table 3).¹¹ In the competition reaction between aldehyde and acetal, no products were formed at all from acetals **9** which are prone to generate the carbocation. The control data with conventional Lewis acids are given in entries 3-7 and 12. Except DBTT, all failed to give the high chemoselectivity. Notably, TiCl₄ having the strongest acidity preferred the acetal and the upon decreasing the acidity, the aldehyde is getting more favored. Apparently, organotin perchlorates are not acidic enough to generate the oxocarbenium and the reaction proceeds via S_N2 mechanism. This is consistent with the fact that **1** cannot induce reaction with less nucleophilic enol silyl ethers derived from ketone (*vide infra*).

Table 3. Competition between Aldehyde and Acetal in Mukaiyama Reaction of Ketene Silyl Acetal.

$$\begin{array}{c}
 \text{RCHO} + \text{R}'\text{R}''\text{C}(\text{OMe})_2 + \text{2} \xrightarrow[\text{CH}_2\text{Cl}_2, -78^\circ\text{C}]{\text{L.A.}} \begin{array}{c} \text{OSiR}^4_3 \\ | \\ \text{R} \\ | \\ \text{R}' \\ | \\ \text{R}'' \end{array} \text{COOR}^3 + \begin{array}{c} \text{OMe} \\ | \\ \text{R}' \\ | \\ \text{R}'' \end{array} \text{COOR}^3 \\
 \text{3} \qquad \qquad \text{9} \qquad \qquad \qquad \qquad \text{4} \qquad \qquad \qquad \qquad \text{10}
 \end{array}$$


entry	L.A.	2	3	9	L.A.:2:3:9	Yield (%)		
						4	10	4:10
1	1a	2a	3a	9a	0.1:1.5:1.0:1.0	94	0	100:0
2	1b	2a	3a	9a	0.1:1.5:1.0:1.0	84	0	100:0
3	Bu ₂ Sn(OTf) ₂	2a	3a	9a	0.1:1.0:1.0:1.0	62	0	100:0
4	TiCl ₄	2a	3a	9a	1.0:1.0:1.0:1.0	12	26	32:68
5	SnCl ₄	2a	3a	9a	0.1:1.0:1.0:1.0	34	5	87:13
6	Ph ₃ CClO ₄	2a	3a	9a	0.1:1.0:1.0:1.0	44	8	85:15
7	TMSOTf	2a	3a	9a	0.1:1.0:1.0:1.0	70	9	89:11
8	1a	2a	3a	9b	0.1:1.5:1.0:1.0	89	0	100:0
9	1b	2a	3a	9b	0.1:1.5:1.0:1.0	82	0	100:0
10	1a	2a	3b	9a	0.1:1.5:1.0:1.0	78	0	100:0
11	1a	2a	3b	9c	0.1:1.0:1.0:1.0	64	0	100:0
12	TMSOTf	2a	3b	9c	0.1:1.0:1.0:1.0	48	23	68:32
13	1a	2d	3a	9a	0.1:1.0:1.0:1.0	87	0	100:0
14	1a	2d	3b	9a	0.1:1.5:1.0:1.0	67	0	100:0
15	1a	2e	3a	9a	0.1:1.5:1.0:1.0	85	0	100:0

PhCH(OMe)₂ (**9a**) PhC(OMe)₂CH₃ (**9b**) *n*-C₇H₁₅C(OMe)₂CH₃ (**9c**)

Competition between Aldehyde and α -Enal. Because of the prevalence of nucleophilic addition towards carbonyls in organic synthesis, differentiation between different carbonyl functions is highly desired.⁹ In general, incorporation of electron-donating group on carbonyl decreases the electrophilicity of the carbonyl carbon. An α,β -unsaturated group also reduces the reactivity of the carbonyl group through the delocalization of π -electrons and, hence, α -enones and -enals are less reactive than the corresponding alkanones and alkanals. The reversal of the reactivity between these carbonyl groups is difficult and finds only a few precedent examples. Reetz disclosed that $\text{CH}_3\text{Ti}(\text{OiPr})_3$ reacted with α -enone and -enal faster than the corresponding aliphatic ketone and aldehyde.¹² He explained this selectivity in terms of the more increased steric hindrance of aliphatic carbonyls. Markó et al. reported the analogous selectivity for the addition of thallium ate complexes to a mixture of enone and ketone; α -enones and acetophenones reacted preferentially over aliphatic ketones (selectivity 5:1~75:1).¹³ In this case, the initial electron transfer from the ate complex to the carbonyls was suggested. Nakai et al. put forth the Lewis acid promoted protocol.¹⁴ In $\text{Eu}(\text{fod})_3$ -catalyzed Mukaiyama reaction of acetals, α -enones were much more reactive than the corresponding saturated ketone. This was interpreted in terms of stronger coordinating ability and, hence, more effective activation of the α -enones. It seems that such differentiation is more difficult between aldehyde pairs because of their higher reactivities. Herein, we describe the high preference of α -enals over aldehyde by virtue of organotin perchlorate catalyst.¹⁵


As shown in Table 4, exposure of an equimolar mixture of **3** and **5** to **2** in the presence of catalytic **1a** resulted in the preferential or exclusive formation of **7** and **8** derived from **5** except one case (entry 4). Aliphatic aldehyde **3b** was most inactive and thus led to the exclusive formation of **7** (and **8**) (entries 2, 6). Benzaldehyde was a little more reactive to result in the decreased selectivities (entries 1, 5). An electron-withdrawing group on the aromatic ring reduced the reactivity of the benzaldehyde derivative leading to the perfect selectivity (entry 3) while an electron-donating group increased the reactivity giving rise to the outcome opposite to the others (entry 4). Dimethyl-substituted ketene silyl acetal **2e** also exhibited the preference for α -enal but gave the Michael adduct predominantly (entry 7).

Table 4. Competition between Aldehyde and α -Enal in Mukaiyama Reaction of Ketene Silyl Acetal.



entry	2	3	5	Yield (%)			4:(7 + 8)
				4	7	8	
1	2a	3a	5b	7	77	0	8:92
2	2a	3b	5b	0	81	0	0:100
3	2a	3d	5b	0	78	0	0:100
4	2a	3c	5b	52	33	0	61:39
5	2a	3a	5a	16	62	4	20:80
6	2a	3b	5a	0	69	4	0:100
7	2e	3a	5b	3	12	64	4:96

transmetallation processes was more unambiguously ruled out by the reaction with TBSClO₄ that was supposed to be formed if the transmetallation had occurred. As depicted in Scheme 1, the TBSClO₄ catalyst exhibited totally different features from **1**. It smoothly catalyzed aldol reaction of enol silyl ether derived from pinacolone **11** with aldehyde and acetal in contrast to the complete ineffectiveness of the organotin perchlorates. On the other hand, octanal which can be activated by **1a** reacted sluggishly with **2a** in the presence of TBSClO₄. Reaction of acetal **9a** with **2a** was also effected by TBSClO₄. It follows from these results that TBSClO₄ did not play any role in the organotin perchlorate-catalyzed reaction.



^aYields in the **1a**-catalyzed reaction are given in parentheses.

Scheme 1.

The reactivities revealed in this study is totally in accord with the SN₂ mechanism. The preference of aldehyde over acetal is the typical example. Organotin perchlorates are not acidic enough to generate the carbocation and the coordination with carbonyl is the crucial step. The importance of this step is reflected on the preferred reaction of α -enals to saturated aldehydes. The SN₂ mechanism also enables differentiation between enol silyl ethers derived from esters and ketones. The former ethers are more nucleophilic so that the reaction is triggered by **1** while the latter ones are not nucleophilic enough to be activated by **1**. Detection of the subtle differences requires to invoke the weak catalytic activities. Hence, the usefulness of gentle Lewis acids for designing highly chemoselective reactions will receive more attention.

EXPERIMENTAL SECTION

Reaction of 2 with 3 (Typical Procedure). To a CH₂Cl₂ solution (1 mL) of Ph₃CClO₄ (34.2 mg, 0.1 mmol) was added Bu₃SnH (30.6 mg, 0.105 mmol) at room temperature. In the meantime, the yellow color disappeared. The solution was stirred for 1 h to form **1a**. To the resulting solution were added a CH₂Cl₂

solution (2 mL) of **3a** (106 mg, 1 mmol) and then **2a** (242 mg, 1.2 mmol) in CH_2Cl_2 (2 mL) at -78°C . After 2 h, the reaction mixture was subjected to aqueous workup and column chromatography on silica gel (1:3 CH_2Cl_2 -hexane) of the crude product afforded **4aa**¹⁹ (274 mg, 89%): $^1\text{H NMR}$ (CDCl_3) δ -0.13 (s, 3H, CH_3), 0.06 (s, 3H, CH_3), 0.89 (s, 9H, t-Bu), 1.25 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.53, 2.71 (ABX, 2H, $J_{\text{AB}}=14.6$, $J_{\text{AX}}=4.0$, $J_{\text{BX}}=9.4\text{Hz}$, CH_2), 4.17 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.17-5.21 (m, 1H, CH), 7.29-7.40 (m, 5H_{arom}). The other reactions were carried out analogously.

4ab: $^1\text{H NMR}$ (CDCl_3) δ 0.03 (s, 3H, CH_3), 0.06 (s, 3H, CH_3), 0.86 (s, 9H, t-Bu), 0.88 (t, 3H, $J=6.5\text{Hz}$, CH_3), 1.25 (t, 3H, $J=7.1\text{Hz}$, CH_3), 1.27-1.49 (m, 12H, 6CH_2), 2.40-2.43 (m, 2H, CH_2), 4.08-4.15 (m, 3H, CH, CH_2). This compound was confirmed by desilylation to give the known alcohol.²⁰

4ac: $^1\text{H NMR}$ (CDCl_3) δ -0.19 (s, 3H, CH_3), -0.01 (s, 3H, CH_3), 0.83 (s, 9H, t-Bu), 1.25 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.50, 2.73 (ABX, 2H, $J_{\text{AB}}=14.4$, $J_{\text{AX}}=4.3$, $J_{\text{BX}}=9.3\text{Hz}$, CH_2), 3.79 (s, 3H, OCH_3), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.08-5.11 (m, 1H, CH), 6.84 (m, 2H_{arom}), 7.25 (m, 2H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²¹

4ad: $^1\text{H NMR}$ (CDCl_3) δ -0.16 (s, 3H, CH_3), -0.01 (s, 3H, CH_3), 0.84 (s, 9H, t-Bu), 1.24 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.52, 2.69 (ABX, 2H, $J_{\text{AB}}=14.8$, $J_{\text{AX}}=4.5$, $J_{\text{BX}}=8.7\text{Hz}$, CH_2), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.16-5.20 (m, 1H, CH), 7.41 (m, 2H_{arom}), 7.62 (m, 2H_{arom}); HRMS calcd for $\text{C}_{18}\text{H}_{28}\text{NO}_3\text{Si}$ ($\text{M}^+ + 1$) 334.1838, found 334.1855.

4ba: $^1\text{H NMR}$ (CDCl_3) δ -0.19 (s, 3H, CH_3), 0.04 (s, 3H, CH_3), 0.85 (s, 9H, t-Bu), 1.42 (s, 9H, t-Bu), 2.46, 2.66 (ABX, 2H, $J_{\text{AB}}=14.9$, $J_{\text{AX}}=4.3$, $J_{\text{BX}}=8.7\text{Hz}$, CH_2), 5.08-5.13 (m, 1H, CH), 7.24-7.35 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²²

4ca: $^1\text{H NMR}$ (CDCl_3) δ 0.47 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.85 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 1.24 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.55, 2.74 (ABX, 2H, $J_{\text{AB}}=14.7$, $J_{\text{AX}}=4.4$, $J_{\text{BX}}=9.9\text{Hz}$, CH_2), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.14-5.18 (m, 1H, CH), 7.25-7.38 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²³

4da: $^1\text{H NMR}$ (CDCl_3) δ -0.30 (s, 3H, CH_3), -0.02 (s, 3H, CH_3), 0.80 (s, 9H, t-Bu), 0.84 (d, 3H, $J=7.1\text{Hz}$, CH_3), 2.61-2.68 (m, 1H, CH), 3.57 (s, 3H, OCH_3), 4.69 (d, 1H, $J=9.3\text{Hz}$, CH), 7.23-7.31 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²⁴

4db: $^1\text{H NMR}$ (CDCl_3) δ 0.03 (s, 3H, CH_3), 0.05 (s, 3H, CH_3), 0.86 (s, 9H, t-Bu), 0.88 (t, 3H, $J=6.5\text{Hz}$, CH_3), 1.10 (d, 3H, $J=7.1\text{Hz}$, CH_3), 1.26-1.47 (m, 12H, 6CH_2), 2.61-2.66 (m, 1H, CH), 3.65 (s, 3H, OCH_3), 4.00-4.02 (m, 1H, CH); HRMS calcd for $\text{C}_{18}\text{H}_{39}\text{O}_3\text{Si}$ ($\text{M}^+ + 1$) 331.2668, found 331.2633.

4ea: $^1\text{H NMR}$ (CDCl_3) δ 0.45 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.83 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 0.96 (s, 3H, CH_3), 1.16 (s, 3H, CH_3), 1.24 (t, 3H, $J=7.1\text{Hz}$, CH_3), 4.13 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.00 (s, 1H, CH), 7.23-7.28 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²⁵

4eb: $^1\text{H NMR}$ (CDCl_3) δ 0.59 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.86 (t, 3H, $J=6.5\text{Hz}$, CH_3), 0.95 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 1.05 (s, 3H, CH_3), 1.14 (s, 3H, CH_3), 1.17-1.42 (m, 15H, 6CH_2 , CH_3), 3.85-3.89 (m, 1H, CH), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2); HRMS calcd for $\text{C}_{20}\text{H}_{43}\text{O}_3\text{Si}$ ($\text{M}^+ + 1$) 359.2981, found 359.3024.

4ec: $^1\text{H NMR}$ (CDCl_3) δ 0.45 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.83 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 0.95 (s, 3H, CH_3), 1.14 (s, 3H, CH_3), 1.26 (t, 3H, $J=7.1\text{Hz}$, CH_3), 3.80 (s, 3H, OCH_3), 4.09 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.95 (s, 1H, CH), 6.81 (m, 2H_{arom}), 7.20 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²⁶

4ed: $^1\text{H NMR}$ (CDCl_3) δ 0.45 (q, 6H, $J=7.8\text{ Hz}$, 3CH_2), 0.84 (t, 9H, $J=7.8$, 3 CH_3), 0.96 (s, 3H, CH_3), 1.16 (s, 3H, CH_3), 1.25 (t, 3H, $J=7.1\text{ Hz}$, CH_3), 4.11 (q, 2H, $J=7.1\text{ Hz}$, CH_2), 5.05 (s, 1H, CH), 7.41 (m, 2H_{arom}), 7.60 (m, 2H_{arom}); HRMS calcd for $\text{C}_{20}\text{H}_{32}\text{NO}_3\text{Si}$ ($\text{M}^+ + 1$) 362.2155, found 362.2188.

4fa: $^1\text{H NMR}$ (CDCl_3) δ -0.35 (s, 3H, CH_3), -0.01 (s, 3H, CH_3), 0.87 (s, 9H, t-Bu), 0.98 (s, 3H, CH_3), 1.16 (s, 3H, CH_3), 3.67 (s, 3H, OCH_3), 4.96 (s, 1H, CH), 7.27 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.²⁷

Reaction of 2 with 5 or 6 (Typical Procedure). To a CH_2Cl_2 solution (0.5 mL) of **1a** (0.1 mmol) were added a CH_2Cl_2 solution (2 mL) of **5a** (70 mg, 1 mmol) and subsequently **2a** (263 mg, 1.3 mmol) in CH_2Cl_2 (2 mL) at -78°C . After 2 h, the reaction mixture was subjected to aqueous workup and column chromatography on silica gel (1:20 EtOAc-hexane) of the crude product afforded **7aa** (221 mg, 81%):²⁸ $^1\text{H-NMR}$ (CDCl_3) δ 0.01 (s, 3H, CH_3), 0.02 (s, 3H, CH_3), 0.84 (s, 9H, t-Bu), 1.24 (t, 3H, $J=7.1\text{Hz}$, CH_3), 1.65 (d, 3H, $J=5.8\text{Hz}$, CH_3), 2.38, 2.49 (ABX, 2H, $J_{\text{AB}}=14.3$, $J_{\text{AX}}=5.2$, $J_{\text{BX}}=8.2\text{Hz}$, CH_2), 4.12 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.51-4.53 (m, 1H, CH), 5.44-5.47 (m, 1H, CH), 5.58-5.65 (m, 1H, CH); **8aa** (17 mg, 6%):²⁸ $^1\text{H NMR}$ (CDCl_3) δ 0.11 (s, 6H, 2CH_3), 0.89 (s, 9H, t-Bu), 1.02 (d, 3H, $J=6.8\text{Hz}$, CH_3), 1.24 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.24 (d, 2H, $J=6.4\text{Hz}$, CH_2), 2.52-2.62 (m, 1H, CH), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.89 (dd, 1H, $J=12.1$, 8.8Hz, CH), 6.27 (d, 1H, $J=12.1\text{Hz}$, CH). The other reactions were conducted analogously.

7ab:²⁹ $^1\text{H NMR}$ (CDCl_3) δ 0.06 (s, 3H, CH_3), 0.08 (s, 3H, CH_3), 0.89 (s, 9H, t-Bu), 1.26 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.51, 2.62 (ABX, 2H, $J_{\text{AB}}=14.4$, $J_{\text{AX}}=5.2$, $J_{\text{BX}}=7.7\text{Hz}$, CH_2), 4.13 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.76-4.78 (m, 1H, CH), 6.19 (dd, 1H, $J=15.8$, 7.0Hz, CH), 6.57 (d, 1H, $J=15.8\text{Hz}$, CH), 7.24-7.38 (m, 5H_{arom}).

7da:²⁸ $^1\text{H NMR}$ (CDCl_3) δ -0.02 (s, 6H, 2CH_3), 0.83 (s, 9H, t-Bu), 1.00 (d, 3H, $J=7.0\text{Hz}$, CH_3), 1.66 (d, 3H, $J=6.9\text{Hz}$, CH_3), 2.46-2.52 (m, 1H, CH), 3.63 (s, 3H, OCH_3), 4.14-4.18 (m, 1H, CH), 5.25-5.35 (m, 1H, CH), 5.56-5.63 (m, 1H, CH).

7db:³⁰ $^1\text{H NMR}$ (CDCl_3) δ 0.23 (s, 6H, 2CH_3), 0.87 (s, 9H, t-Bu), 1.08 (d, 3H, $J=7.0\text{ Hz}$, CH_3), 2.60-2.67 (m, 1H, CH), 3.65 (s, 3H, OCH_3), 4.38-4.43 (m, 1H, CH), 6.06 (dd, 1H, $J=15.9$, 7.0 Hz, CH), 6.53 (d, 1H, $J=15.9\text{ Hz}$, CH), 7.30-7.40 (m, 5H_{arom});

7ea: $^1\text{H NMR}$ (CDCl_3) δ 0.53 (q, 6H, $J=7.8\text{ Hz}$, 3CH_2), 0.92 (t, 9H, $J=7.8\text{ Hz}$, 3CH_3), 1.03 (s, 3H, CH_3), 1.13 (s, 3H, CH_3), 1.24 (t, 3H, $J=7.1\text{ Hz}$, CH_3), 1.68 (d, 3H, $J=6.1\text{Hz}$, CH_3), 4.10 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.24 (d, 1H, $J=8.1\text{Hz}$, CH), 5.39 (dd, 1H, $J=15.3$, 8.1Hz, CH), 5.55-5.62 (m, 1H, CH); HRMS calcd for $\text{C}_{16}\text{H}_{33}\text{O}_3\text{Si}$ ($\text{M}^+ + 1$) 301.2199, found 301.2205.

7eb: $^1\text{H NMR}$ (CDCl_3) δ 0.58 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.93 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 1.10 (s, 3H, CH_3), 1.20 (s, 3H, CH_3), 1.26 (t, 3H, $J=7.1\text{Hz}$, CH_3), 4.15 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.49 (d, 1H, $J=7.9\text{Hz}$, CH), 6.14 (dd, 1H, $J=15.9$, 7.9Hz, CH), 6.50 (d, 1H, $J=15.9\text{Hz}$, CH), 7.25-7.39 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known alcohol.³¹

8ac:¹⁷ $^1\text{H NMR}$ (CDCl_3) δ -0.16 (s, 3H, CH_3), -0.14 (s, 3H, CH_3), 0.89 (s, 9H, t-Bu), 0.99 (d, 3H, $J=6.8\text{Hz}$, CH_3), 1.16 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.16-2.19 (m, 2H, CH_2), 3.05-3.16 (m, 1H, CH), 4.03 (q, 2H, $J=7.1\text{Hz}$, CH_2), 4.83 (d, 1H, $J=9.6\text{Hz}$, CH), 7.29-7.40 (m, 5H_{arom}).

8da:²⁸ $^1\text{H NMR}$ (CDCl_3) δ 0.11 (s, 6H, 2CH_3), 0.90 (s, 9H, t-Bu), 0.97 (d, 3H, $J=6.5\text{Hz}$, CH_3), 1.09 (d, 3H, $J=6.5\text{Hz}$, CH_3), 2.23-2.36 (m, 2H, 2CH), 3.63 (s, 3H, OCH_3), 4.74-4.92 (m, 1H, CH), 6.23 (d, 1H, $J=11.8\text{Hz}$, CH).

8db:³² $^1\text{H NMR}$ (CDCl_3) δ 0.96 (d, 3H, $J=7.0\text{Hz}$, CH_3), 2.68-2.89 (m, 3H, CH, CH_2), 3.39-3.45 (m, 1H, CH), 3.70 (s, 3H, OCH_3), 7.16-7.34 (m, 5H_{arom}), 9.54 (s, 1H, CHO).

8dc:³³ $^1\text{H NMR}$ (CDCl_3) δ -0.01 (s, 3H, CH_3), 0.08 (s, 3H, CH_3), 0.98 (s, 9H, t-Bu), 1.10 (d, 3H, $J=6.8\text{Hz}$, CH_3), 1.18 (d, 3H, $J=6.8\text{Hz}$, CH_3), 2.35-2.50 (m, 1H, CH), 2.67-2.76 (m, 1H, CH), 3.68 (s, 3H, OCH_3), 4.87 (d, 1H, $J=10.9\text{Hz}$, CH), 7.35-7.49 (m, 5H_{arom}).

8ea: $^1\text{H NMR}$ (CDCl_3) δ 0.66 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.92 (d, 3H, $J=7.2\text{Hz}$, CH_3), 0.98 (t, 9H, $J=7.7\text{Hz}$, 3CH_3), 1.06 (s, 3H, CH_3), 1.08 (s, 3H, CH_3), 1.25 (t, 3H, $J=7.1\text{Hz}$, CH_3), 2.34-2.40 (m, 1H, CH), 4.11 (q,

2H, $J=7.1\text{Hz}$, CH_2), 4.87 (dd, 1H, $J=11.8, 9.8\text{Hz}$, CH), 6.27(d, 1H, $J=11.8\text{Hz}$, CH); HRMS calcd for $\text{C}_{16}\text{H}_{33}\text{O}_3\text{Si}$ ($M^+ + 1$) 301.2199, found 301.2197.

8eb: ^1H NMR (CDCl_3) δ 0.64 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.95 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 1.07 (s, 3H, CH_3), 1.14 (s, 3H, CH_3), 1.20 (t, 3H, $J=7.1\text{Hz}$, CH_2), 3.50 (d, 1H, $J=10.7\text{Hz}$, CH), 4.04 (q, 2H, $J=7.1\text{Hz}$, CH_2), 5.43 (dd, 1H, $J=11.7, 10.7\text{Hz}$, CH), 6.32 (d, 1H, $J=11.7\text{Hz}$, CH), 7.14-7.28 (m, 5H_{arom}). This compound was confirmed by desilylation to give the known aldehyde.³⁴

8ec: ^{17}H NMR (CDCl_3) δ 0.62 (q, 6H, $J=7.8\text{Hz}$, 3CH_2), 0.92 (t, 9H, $J=7.8\text{Hz}$, 3CH_3), 0.96 (d, 3H, $J=6.7\text{Hz}$, CH_3), 1.06 (s, 3H, CH_3), 1.08 (s, 3H, CH_3), 1.15 (t, 3H, $J=7.1\text{Hz}$, CH_2), 2.67-2.73 (m, 1H, CH), 3.95-4.10 (m, 2H, CH_2), 4.93 (d, 1H, $J=11.2\text{Hz}$, CH), 7.26-7.39 (m, 5H_{arom}).

Competition reaction (General Procedure). To a CH_2Cl_2 solution (1 mL) of **1a** (0.1 mmol) were added a mixture of electrophiles (1.0 mmol each) in CH_2Cl_2 (3 mL) and subsequently **2** (1.5 mmol) in CH_2Cl_2 (2 mL) at -78°C . After 2 h, aqueous workup and evaporation afforded a crude product that was analyzed by GLC.

Competition Reaction of Enol Silyl Ethers Derived from Ester and Ketone (Typical Procedure). To a CH_2Cl_2 solution (1 mL) of **1a** (0.1 mmol) were added a CH_2Cl_2 solution (2 mL) of **3a** (106 mg, 1.0 mmol) at -78°C . Then, **2a** (202 mg, 1.0 mmol) in CH_2Cl_2 (3 mL) and **11** (174 mg, 1.0 mmol) in CH_2Cl_2 (3 mL) were added. The solution was stirred for 3 h. Aqueous workup and evaporation afforded a crude product that was analyzed by GLC.

Preparation of 13 (two stereoisomers). To a THF solution (90 mL) of diisopropylamine (11.2 g, 110 mmol) was added BuLi (1.6 M hexane solution, 69 mL, 110 mmol) at 0°C . The solution was cooled to -78°C and methyl β -acetylpropionate (6.5 g, 50 mmol) was added. The reaction mixture was stirred at -78°C for 1 h and trimethylsilyl chloride (15.1 g, 140 mmol) was added rapidly. After stirring for 1 h at room temperature, the reaction mixture was filtered and then THF was evaporated. The residue was dissolved in 150 mL of pentane and the salt precipitated was removed by filtration. After evaporation, the residue was distilled under reduced pressure ($69\text{-}73^\circ\text{C}/3\text{ mmHg}$) to give **13** (4.80 g, 35%, 7:3 *E/Z* mixture). ^1H NMR (CDCl_3) δ 0.19 (major) and 0.20 (minor) (each s, 9H, 3CH_3), 0.21 (major) and 0.22 (minor) (each, s, 9H, 3CH_3), 2.68 (d, 2H, $J=7.1\text{Hz}$, CH_2), 3.51 (minor) and 3.52 (major) (each s, 3H, OCH_3), 3.56 (t, 1H, $J=7.1\text{Hz}$, CH), 4.03 (minor) and 4.05 (major) (each s, 1H, CH), 4.11 (minor) and 4.13 (major) (each s, 1H, CH).

Reaction of 13 with Aldehyde. To a CH_2Cl_2 solution (1 mL) of **1a** (0.1 mmol) were added a CH_2Cl_2 solution (2 mL) of **3a** (106 mg, 1.0 mmol) and then **13** (329 mg, 1.2 mmol) in CH_2Cl_2 (2 mL) at -78°C . After 2 h, 2-(hydroxymethyl)pyridine (33 mg, 0.3 mmol) and pyridine (0.1 mL) were added. The resulting mixture was diluted with hexane (60 mL) and washed with aqueous NaHCO_3 . The organic layer was dried (Na_2SO_4) and evaporated. The residue was chromatographed on silica gel (1:8 EtOAc-hexane) to give **14a** (266 mg, 86%). ^1H NMR (CDCl_3) δ -0.02 (s, 9H, 3CH_3), 2.03 (s, 3H, CH_3), 2.32, 2.69 (ABX, 2H, $J_{\text{AB}}=17.8$, $J_{\text{AX}}=3.6$, $J_{\text{BX}}=10.6\text{Hz}$, CH_2), 3.21-3.28 (m, 1H, CH), 3.70 (s, 3H, OCH_3), 4.84 (d, 1H, $J=7.7\text{Hz}$, CH), 7.24-7.34 (m, 5H_{arom}); HRMS calcd for $\text{C}_{16}\text{H}_{25}\text{O}_4\text{Si}$ ($M^+ + 1$) 309.1522, found 309.1500. The other reaction was carried out analogously to afford **14b** which was confirmed as the corresponding alcohol after desilylation: ^1H NMR (CDCl_3) δ 0.87 (t, 3H, $J=6.2\text{Hz}$, CH_3), 1.24-1.80 (m, 12H, 6CH_2), 2.18 (s, 3H, CH_3), 2.75-2.83 (m, 1H, CH), 2.92-3.08 (m, 2H, CH_2), 3.70 (s, 3H, OCH_3), 3.97-4.05 (m, 1H, CH); HRMS calcd for $\text{C}_{14}\text{H}_{25}\text{O}_3$ ($M^+ - \text{OH}$) 241.1804, found 241.1804.

Reaction of 13 with 6. The reaction was carried out analogously as described above employing **6** (146 mg, 1.0 mmol) in place of **13** to give **15** (216 mg, 62%). This compound was confirmed as follows: **15** was treated with Bu_4NF (178 mg, 0.68 mmol) and THF (3 mL) at room temperature for 1 h. Aqueous NaHCO_3 was added to this solution, and the mixture was extracted with ethyl acetate. The organic layer was dried and

evaporated. The residue was chromatographed on silica gel (1:6 EtOAc-hexane) to give 6-benzoyl-4-methoxycarbonyl-5-methyl-2-hexanone: $^1\text{H NMR}$ (CDCl_3) δ 0.99 (d, 3H, $J=6.8\text{Hz}$, CH_3), 2.18 (s, 3H, CH_3), 2.49-3.14 (m, 6H, 2CH, 2 CH_2), 3.68 (s, 3H, OCH_3), 7.28-7.60 (m, 5 H_{arom}): HRMS calcd for $\text{C}_{16}\text{H}_{21}\text{O}_4$ ($M^+ + 1$) 277.1440, found 277.1487.

Mukaiyama reaction catalyzed by TBSClO_4 was carried out in the same way as organotin perchlorate-catalyzed reaction.

REFERENCES

1. Davies, A. G.; Smith, P. J. In *Comprehensive Organometallic Chemistry*; Wilkinson, G. Ed.; Pergamon Press: Oxford, 1982; Chapter 11.
2. Pereyre, M.; Quintard, J.-P.; Rahm, A. *Tin in Organic Synthesis*; Butterworths: London, 1987.
3. Otera, J. In *Advanced in Detailed Reaction Mechanisms*; Coxon, J. M. Ed.; JAI Press Inc.: London, 1994, Vol. 3, p. 167. Otera, J. *Chem. Rev.* **1993**, *93*, 1449.
4. (a) Sato, T.; Otera, J.; Nozaki, H. *J. Am. Chem. Soc.* **1990**, *112*, 901. (b) Sato, T.; Wakahara, Y.; Otera, J.; Nozaki, H. *Tetrahedron Lett.* **1990**, *31*, 1581. (c) Sato, T.; Wakahara, Y.; Otera, J.; Nozaki, H. *Tetrahedron* **1991**, *47*, 9773.
5. Chen, J.; Sakamoto, K.; Orita, A.; Otera, J. *Synlett* **1996**, 877.
6. Lambert, J. B.; Kuhlmann, B. *J. Chem. Soc., Chem. Commun.* **1992**, 931.
7. Murata, S.; Suzuki, M.; Noyori, R. *Tetrahedron* **1988**, *44*, 4259.
8. Mori, A.; Maruoka, K.; Yamamoto, H. *Tetrahedron Lett.* **1984**, *25*, 4421.
9. Sato, J.; Otera, J.; Nozaki, H. *J. Org. Chem.* **1993**, *58*, 4971, and references cited therein.
10. Mukaiyama, T.; Ohno, T.; Han, J. S.; Kobayashi, S. *Chem. Lett.* **1991**, 949.
11. For a preliminary report: Otera, J.; Chen, J. *Synlett* **1996**, 321.
12. Reetz, M. T. *Organotitanium Reagents in Organic Synthesis*; Springer-Verlag: Berlin, 1986, Chapter 3.
13. Markó, I. E.; Leung, C. W. *J. Am. Chem. Soc.* **1994**, *116*, 371.
14. Hanyuda, K.; Hirai, K.; Nakai, T. *Synlett* **1997**, 31.
15. For a preliminary report: Chen, J. Otera, J. *Synlett* **1997**, 29.
16. Denmark, S. E.; Almstead, N. G. *J. Am. Chem. Soc.* **1993**, *115*, 3133.
17. Inaba, S.-I.; Ojima, I. *Tetrahedron Lett.* **1977**, 2009. Hirai, K.; Ojima, I. *ibid.* **1983**, *24*, 785. Nakamura, E.; Kuwajima, I. *ibid.* **1983**, *24*, 3347. Otera, J.; Fujita, Y.; Sakuta, N.; Fujita, M.; Fukuzumi, S. *J. Org. Chem.* **1996**, *61*, 2951.
18. Carreira, E. M.; Singer, R. A. *Tetrahedron Lett.* **1994**, *35*, 4323.
19. Mukaiyama, T.; Takashima, T.; Kusaka, H.; Shimpuku, T. *Chem. Lett.* **1990**, 1777.
20. Han, B. H.; Boudjouk, P. *J. Org. Chem.* **1982**, *47*, 5030.
21. Huang, Y. Z.; Liao, Y. *J. Org. Chem.* **1991**, *56*, 1381.
22. Burkhardt, E. R.; Bergman, R. G.; Heathcock, C. H. *Organometallics*, **1990**, *9*, 30.
23. Mukaiyama, T.; Kobayashi, S.; Sano, T. *Tetrahedron*, **1990**, *46*, 4653.
24. Heathcock, C. H.; Buse, C. T.; Kleschick, W. A.; Pirrung, M. C.; Sohn, J. E.; Lampe, J. *J. Org. Chem.* **1980**, *45*, 1066.
25. Parmee, E. R.; Tempkin, O.; Masamune, S. *J. Am. Chem. Soc.* **1991**, *113*, 9365.
26. Unterhalt, B.; Weyrich, K. *Arch. Pharm.* **1980**, *313*, 795.
27. Reetz, M. T.; Vougioukas, A. E. *Tetrahedron Lett.* **1987**, *28*, 793.
28. Mikami, K.; Terada, M.; Nakai, T. *J. Org. Chem.* **1991**, *56*, 5456.
29. Hara, R.; Mukaiyama, T. *Chem. Lett.* **1989**, 1909.
30. Isayama, S.; Mukaiyama, T. *Chem. Lett.* **1989**, 2005.
31. Kiyooka, S.; Kaneko, Y.; Komura, M.; Matsuo, H.; Nakano, M. *J. Org. Chem.* **1991**, *56*, 2276.
32. Shimada, S.; Sagio, K.; Abe, M.; Sudo, A.; Hasegawa, M. *Chem. Lett.* **1992**, 1445.
33. Mukaiyama, T.; Tamura, M.; Kobayashi, S. *Chem. Lett.* **1986**, 1817.
34. Moreau, J. L.; Gaudemar, M. *C. R. Acad. Sci. Ser. 2*, **1985**, *300*, 399.