

Total synthesis of (±)-camphorataimides and (±)-himanimides by NaBH₄/Ni(OAc)₂ or Zn/AcOH stereoselective reduction

Chia-Fu Cheng, Zhen-Chang Lai, Yean-Jang Lee*

Department of Chemistry, National Changhua University of Education, Changhua 50058, Taiwan

Received 22 January 2008; received in revised form 19 February 2008; accepted 22 February 2008

Available online 4 March 2008

Abstract

Maleic anhydride **1** of *Antrodia camphorate*, which can be isolated from Chinese herbal medicine, is achieved in which the longest linear sequence is only five steps, in 40% overall yield from commercially available succinic anhydride. The crucial antrodinimides **3** and **2** can be readily transformed by the chemoselective reduction with Zn/AcOH and NaBH₄/Ni(OAc)₂·4H₂O to afford the naturally occurring camphorataimides, **4** and **5**, in high yields as well, respectively. This synthetic strategy can also be modified to give access to a variety of different maleic acid derivatives, himanimides **6–8**.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: *Antrodia camphorate*; Himanimides; NaBH₄/Ni(OAc)₂; Zn/AcOH; Stereoselective reduction

1. Introduction

As part of our research aims to uncover from natural products new compounds with improved biological activities, including antioxidant, anti-human immunodeficiency virus (HIV), and tumor growth inhibition activities, we attended to the coumarin and benzofuran family. Recently, several structurally interesting compounds with substituted maleic anhydride and maleimide moieties have been isolated as bioactive natural products.¹ Studies on the constituents of the mycelium of *Antrodia camphorate* (Polyporaceae, Aphyllophorales),^{1b} which is used as a traditional Chinese medicine, showed them to contain a number of pharmacological interests.² Maleic anhydride **1**, maleimide **2** and **3** together with succinic acids **4** and **5** were isolated from the chloroform-soluble fraction of the mycelium of *A. camphorate* and repeatedly chromatographed on silica gel with *n*-hexane/acetone and CHCl₃/MeOH. In addition, maleimides **2** and **3** showed appreciable cytotoxic activity against Lewis lung carcinoma cell lines. Recently, during the syntheses of the phytotoxins,

carpacin³ and ailanthoidol,⁴ we became aware of the facile conversion of *p*-alkoxyphenyl bromide to aryl stannanes and the proceeding Stille⁵ coupling of bromomaleic acid derivatives with stannanes to substituted maleate skeleton. Therefore, **1–3** are ideal synthetic target for the application of these reactions. Although **1–3** have been synthesized by Argade^{6a} and Stewart,^{6b} they were achieved with very time-consuming and complicated synthetic approaches. To date, no **4** and **5** have been synthesized and deficiently examined the biological activity of the molecules, we undertook the first total synthesis of (±)-camphorataimides, **4** and **5**, the structural novelty of this family (Fig. 1).

Perhaps the easiest route to **1** would have been to sequentially couple the known dibromomaleic anhydride **10**, *i*-BuMgBr, and the known⁷ bromo(prenyloxy)benzene. In previous projects⁸ we have achieved such coupling by treating a mixture of two aryl bromides with a bis(trialkyltin) in the presence of a palladium catalyst. The reaction proceeds via in situ conversion of one aryl bromide to the corresponding stannane followed by coupling of the latter with the other aryl bromide.^{8a} In addition, although Stewart et al. have achieved cross-coupling by the Negishi and Suzuki reactions, we would still like to investigate the versatility of Stille cross-

* Corresponding author. Tel.: +886 4 7232105x3511; fax: +886 4 7211190.
E-mail address: leeyj@cc.ncue.edu.tw (Y.-J. Lee).




Figure 1. Maleic anhydride and maleimide natural products of *Antrodia camphorate* and himanimide.

coupling reactions to provide **1**, hence synthesizing the target (±)-camphorataimides **4** and **5**. Herein, we report the first synthesis of the naturally occurring (±)-*cis*-camphorataimide **D** and applied the Stille reaction and stereoselective reduction with Zn/AcOH to achieve (±)-*cis*-himanimide **D**.

2. Results and discussion


In the beginning of our synthesis, although **10** has been synthesized by several groups,⁹ it was achieved with very time-consuming and complicated synthetic approaches. To overcome these technical difficulties, we report our studies on the synthesis of **10** from the commercially available succinic anhydride with Br₂ in a PTFE-vessel reactor in very good yield (89%). In the case of **10** and 4-bromo(prenyloxy)benzene, however, the palladium-catalyzed reaction with (Me₃Sn)₂ to give **1** was unsuccessful. The major product was biphenyl, the product from the homocoupling of 4-bromo(prenyloxy)benzene.

With the aim of developing a successful route to **1**, an alternative method of construction was examined (Scheme 1). Coupling of dibromide **11** with the commercially available isobutyl magnesium bromide proceeded in dioxane by Pd(0)-catalysis to give the desired **12** in very poor 12% yield. The isolated yield might be low because Grignard reagent and

pyrroledione **11** appeared a competitive reactant in the 1,2-addition of carbonyl group. Fortunately, the expected **12** can be afforded with CuI instead of Pd-catalyst in moderate 61% yield. Similarly, bromomaleimide **12** and the corresponding stannane **13** could also undergo the palladium-catalyzed cross-coupling to give the key product **14** in an excellent yield (94%). The resultant benzyl maleimide **14** was then transformed into the target compound **1** under reflux in KOH/H₂O/THF/MeOH. Finally, the anhydride **1** was readily converted by heating at 85 °C in AcONH₄ or NH₂OH·HCl/pyridine to afford the natural products, maleimides **2** and **3**, respectively. ¹H and ¹³C NMR spectra of the synthetic products are in agreement with those reported for the naturally derived materials.¹

A further insight into the nature of NaBH₄ with metallic salts,¹⁰ it was envisioned that a chemoselective reduction of **3** could prove legible for the formation of the thermodynamically more stable (±)-*trans*-camphorataimide **E** (**5**). With the crucial intermediate **14** in hand, the chemoselective reduction was smoothly testified with NaBH₄/Ni(OAc)₂·4H₂O in mixed solvent (THF/MeOH 1:1) to provide the expected (±)-*trans*-succinimide **15** only in good yield 65%. Unfortunately, the resultant **15** wasn't then transformed into the corresponding anhydride under reflux in KOH/H₂O/THF/MeOH. Subsequently, the task turned to extend the original system with a functioning prototype so that the variety of substrates was evaluated by 1,4-reduction with the Luche's reagent. For instance, compound **2** was employed by the Luche's method to give the desired *cis/trans*-**17** (1:2 ratio) in high yield 78%. It is worth noting that the stereoselectivity preferentially provided the *trans*-isomer with the Luche's reagent, but the mechanistic issues uncovered in the stereoselective process as well as the scope and generality of this area will be probed in detail in our continuing studies. In addition, the N–O bond of **3** and **16** was first broken to provide **2**, and then efficiently reduced to afford the precedent **17** in high yield. Fortunately, and to our greater satisfaction, we succeeded in obtaining (±)-*trans*-isomer **18** only from **17**, which epimerized the C-3 position to give the thermodynamically more stable *trans* diastereomer as the major product in an excellent 94% yield, suitable for further transformation to achieve the desired **5** (Scheme 2).

With the aim of developing a successful route to (±)-*cis*-**4**, an alternative method of construction, which was recently



Scheme 1.

4.1.1. 2,3-Dibromomaleic anhydride (**10**)

A mixture of succinic anhydride (1.01 g, 10.1 mmol), and bromine (3.70 g, 23.1 mmol) was stirred in a small autoclave (PTFE-lined vessel)¹³ at 210 °C (sand bath) for 3 h (*Caution*: Don't run over 2.00 g each time or raise temperature rapidly, otherwise, it might appear explosion.). After cooling down to room temperature, the mixture was dissolved in ethyl ether. The solvent was then removed and the residue was purified by chromatography (SiO₂, 1:4 hexane/diethyl ether) to give 2,3-dibromomaleic anhydride (2.30 g, 89%) as a white solid, mp 116–117 °C (lit.^{9g} mp 113–114 °C). ¹³C NMR (CDCl₃) δ 125.0, 163.0. Anal. Calcd for C₄O₃Br₂: C, 18.78; O, 18.76. Found: C, 18.96; O, 18.90.

4.1.2. *N*-Benzyl-3,4-dibromomaleimide (**11**)

This compound was prepared as a white solid from anhydride **10** according to the procedure of Viaud-Massuard.^{9g} To a solution of **10** (4.62 g, 18.0 mmol) in acetic acid (20.0 mL) was added benzylamine (2.0 mL, 18.0 mmol) dropwise at room temperature. The mixture was stirred for 10 min and then heated at reflux for 2 h. After cooling down to room temperature, the mixture was extracted with EtOAc (3×30.0 mL). The solvent was then removed and the residue was purified by chromatography (SiO₂, 3:1 hexane/CH₂Cl₂) to give **11** (5.32 g, 86%) as a white solid, mp 112–113 °C (lit.¹⁴ mp 117–117.5 °C). ¹H NMR (CDCl₃) δ 4.75 (s, 2H), 7.31–7.35 (m, 5H); ¹³C NMR (CDCl₃) δ 43.1, 128.2, 128.7, 128.8, 129.4, 135.1, 163.5. Anal. Calcd for C₁₁H₇O₂NBr₂: C, 38.30; H, 2.05; N, 4.06; O, 9.28. Found: C, 38.59; H, 1.74; N, 4.22; O, 9.23.

4.1.3. *N*-Benzyl-3-bromo-4-isobutylmaleimide (**12**)

A solution of benzylidibromomaleimide **11** (1.04 g, 3.0 mmol) and CuI (0.02 g, 0.1 mmol) in anhydrous THF (30.0 mL) was cooled down to 0 °C under N₂ pressure for 20 min. The commercially available Grignard reagent (isobutyl magnesium bromide; 2 M, 2.2 mL) was added dropwise at 0 °C, and allowed to warm to room temperature. After stirring for 8 h at room temperature, the reaction mixture was quenched with saturated aqueous NH₄Cl (1.2 mL) and added MgSO₄ for the removal of water. The resulting solution was filtered and the filtrate evaporated. The product was isolated by flash column chromatography (SiO₂, 9:1 cyclohexane/CH₂Cl₂) to give **12** (0.59 g, 61%) as an oil. ¹H NMR (CDCl₃) δ 0.95 (d, *J*=7.2 Hz, 6H), 2.08 (sep, *J*=7.2 Hz, 1H), 2.35 (d, *J*=7.2 Hz, 2H), 4.69 (s, 2H), 7.27–7.36 (m, 5H); ¹³C NMR (CDCl₃) δ 22.6, 27.8, 34.1, 42.2, 125.5, 127.9, 128.4, 128.6, 135.8, 145.1, 165.1, 168.9; HRMS (EI) calcd for C₁₅H₁₆O₂NBr (M⁺) 321.0368, found 321.0363.

4.1.4. 4-[(3-Methyl-2-butenyloxy)phenyl]bromide

This compound was prepared as a colorless oil from 4-bromophenol according to the procedure of Bernard.⁷ To a mixture of 4-bromophenol (10.0 g, 57.8 mmol), KI (0.58 g, 3.5 mmol), and K₂CO₃ (19.94 g, 144.5 mmol) in acetone (100.0 mL) was added 3-methyl-2-butenyl bromide (11.20 g, 75.1 mmol) at room temperature, and then refluxed at 80 °C

for 12 h. After cooling, the solution was evaporated in vacuo and the brown solid was subjected to flash chromatography (SiO₂, 5:1 hexane/diethyl ether) to obtain the known⁷ 4-[(3-methyl-2-butenyloxy)phenyl]bromide (13.88 g, 57.8 mmol) in 100% yield. ¹H NMR (CDCl₃) δ 1.73 (s, 3H), 1.79 (s, 6H), 4.47 (d, *J*=6.6 Hz, 2H), 5.46 (br t, *J*=6.6 Hz, 1H), 6.78 (d, *J*=9.0 Hz, 2H), 7.35 (d, *J*=9.0 Hz, 2H); ¹³C NMR (CDCl₃) δ 18.2, 25.8, 65.0, 112.7, 116.5, 119.3, 132.2, 138.5, 157.9.

4.1.5. 4-[(3-Methyl-2-butenyloxy)phenyl]tributylstannane (**13**)

The stannane **13** was prepared, using the previous procedure,^{8f} from 4-[(3-methyl-2-butenyloxy)phenyl]bromide. A solution of 4-[(3-methyl-2-butenyloxy)phenyl]bromide (1.02 g, 4.2 mmol) in 60.0 mL of dry THF was placed in a dry, two-necked, 100-mL, round-bottomed flask under a nitrogen atmosphere with a small magnetic stir bar. The flask was then immersed in an acetone/dry ice bath. *n*-Butyllithium solution (2.9 mL, 1.6 M in hexanes, 4.6 mmol) was introduced slowly (1 drop/2 s) during which time the solution changed color from pale yellow to dark orange. After the solution was stirred for 15 min at –78 °C, the reaction was treated at this temperature with neat tributyltin chloride (1.51 g, 4.6 mmol). The dry ice bath was removed and the solution was allowed to reach room temperature. The reaction was quenched with 0.5 mL of H₂O and the solution concentrated in vacuo. After removal of the solvent, the residue was purified by chromatography (SiO₂, 3:1 hexane/CH₂Cl₂) to give **13** (1.82 g, 4.0 mmol) in 96% yield as colorless oil. ¹H NMR (CDCl₃) δ 0.84–1.05 (m, 15H), 1.27–1.69 (m, 12H), 1.73 (s, 3H), 1.79 (s, 3H), 4.50 (d, *J*=6.6 Hz, 2H), 5.50 (br t, *J*=6.6 Hz, 1H), 6.91 (d, *J*=8.3 Hz, 2H), 7.35 (d, *J*=8.3 Hz, 2H); ¹³C NMR (CDCl₃) δ 9.6, 13.7, 18.1, 25.8, 27.4, 29.1, 64.4, 114.6, 119.8, 129.4, 137.4, 138.0, 159.0; HRMS (EI) calcd for C₂₃H₄₀OSn (M⁺) 452.2101, found 452.2107.

4.1.6. *N*-Benzyl-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]maleimide (**14**)

Bromide **12** (0.52 g, 1.6 mmol), tetrakis(triphenylphosphine)palladium(0) (0.18 g, 10 mol %), and stannane **13** (0.73 g, 1.6 mmol) were introduced into a sealable tube containing anhydrous dioxane (10.0 mL), and the reaction mixture degassed. The tube was sealed and heated for 5 h at 145 °C, after cooling, and the solution was filtered and evaporated in vacuo. The filtrate residue was subjected to flash chromatography (SiO₂, 1:3 CH₂Cl₂/diethyl ether) to provide the desired **14** (0.61 g, 94%) as a light green-yellow oil. ¹H NMR (CDCl₃) δ 0.89 (d, *J*=7.2 Hz, 6H), 1.87 (s, 3H), 1.88 (s, 3H), 2.07 (sep, *J*=7.2 Hz, 1H), 2.51 (d, *J*=7.2 Hz, 2H), 4.55 (d, *J*=6.6 Hz, 2H), 4.72 (s, 2H), 5.49 (br t, *J*=6.6 Hz, 1H), 6.96 (d, *J*=9.0 Hz, 2H), 7.49 (d, *J*=9.0 Hz, 2H), 7.24–7.40 (m, 5H); ¹³C NMR (CDCl₃) δ 18.2, 22.7, 25.8, 28.1, 32.9, 41.7, 64.8, 114.7, 119.2, 121.4, 127.7, 128.4, 128.6, 130.9, 136.7, 137.7, 138.1, 138.6, 159.9, 171.1, 171.9; HRMS (EI) calcd for C₂₆H₂₉O₃N (M⁺) 403.2147, found 403.2149.

4.1.7. 3-Isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]furan-2,5-dione (**1**)

A mixture of benzyl maleimide **14** (0.45 g, 1.1 mmol) in THF/MeOH (50.0/50.0 mL), water (50.0 mL), and KOH (0.43 g) was refluxed for 3 h. After it was poured into water, acidified with 1 N HCl, and then extracted with EtOAc. The organic phase was washed with saturated aqueous NaHCO₃ and brine, and then dried over MgSO₄. After removal of the solvent, the residue was purified by flash chromatography (SiO₂, 7:3 EtOAc/cyclohexane) to provide **1** (0.32 g, 92%) as an oil. ¹H NMR (CDCl₃) δ 0.94 (d, *J*=7.2 Hz, 6H), 1.77 (s, 3H), 1.82 (s, 3H), 2.13 (sep, *J*=7.2 Hz, 1H), 2.60 (d, *J*=7.2 Hz, 2H), 4.55 (d, *J*=6.6 Hz, 2H), 5.50 (br t, *J*=6.6 Hz, 1H), 7.02 (d, *J*=9.0 Hz, 2H), 7.62 (d, *J*=9.0 Hz, 2H); ¹³C NMR (CDCl₃) δ 18.2, 22.7, 25.8, 27.9, 33.6, 65.0, 115.1, 118.9, 119.9, 131.1, 139.0, 139.8, 140.2, 160.9, 165.4, 166.4; HRMS (EI) calcd for C₁₉H₂₂O₄ (M⁺) 314.1518, found 314.1513.

4.1.8. 3-Isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]-1H-pyrrol-2,5-dione (**2**)

A solution of furandione **1** (0.22 g, 0.7 mmol) and ammonium acetate (0.32 g, 4.2 mmol) in acetic acid (20.0 mL) was heated at reflux for 24 h. After cooling, the solvent was removed in vacuo and the resulting solid was subjected to column chromatography (SiO₂, 1:1 diethyl ether/CH₂Cl₂) to obtain **2** (0.19 g, 88%) as a white solid, mp 104.5–105 °C (lit.¹ mp 110–111 °C). ¹H NMR (CDCl₃) δ 0.90 (d, *J*=7.2 Hz, 6H), 1.76 (s, 3H), 1.81 (s, 3H), 2.06 (sep, *J*=7.2 Hz, 1H), 2.51 (d, *J*=7.2 Hz, 2H), 4.56 (d, *J*=6.9 Hz, 2H), 5.50 (br t, *J*=6.9 Hz, 1H), 6.99 (d, *J*=6.9 Hz, 2H), 7.51 (d, *J*=6.9 Hz, 2H), 7.69 (br s, NH); ¹³C NMR (CDCl₃) δ 18.2, 22.7, 25.8, 28.1, 32.8, 64.9, 114.8, 119.2, 121.1, 130.9, 138.6, 138.7, 139.1, 160.0, 171.3, 171.9; HRMS (EI) calcd for C₁₉H₂₃NO₃ (M⁺) 313.1678, found 313.1676.

4.1.9. 1-Hydroxy-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]-1H-pyrrol-2,5-dione (**3**)

A mixture of furandione **1** (0.19 g, 0.6 mmol) and hydroxylamine hydrochloride (0.10 g, 1.4 mmol) in pyridine (10.0 mL) was heated at reflux for 2 h. After cooling, the solvent was removed in vacuo and the resulting solid was subjected to column chromatography (SiO₂, 1:10 EtOAc/cyclohexane) to obtain **3** (0.16 g, 82%) as a colorless oil. ¹H NMR (CDCl₃) δ 0.89 (d, *J*=7.2 Hz, 6H), 1.76 (s, 3H), 1.81 (s, 3H), 2.04 (sep, *J*=7.2 Hz, 1H), 2.51 (d, *J*=7.2 Hz, 2H), 4.56 (d, *J*=6.6 Hz, 2H), 5.49 (br t, *J*=6.6 Hz, 1H), 6.98 (d, *J*=8.7 Hz, 2H), 7.51 (d, *J*=8.7 Hz, 2H), 8.53 (br s, 1H); ¹³C NMR (CDCl₃) δ 18.2, 22.7, 25.8, 28.1, 33.0, 64.9, 114.9, 119.0, 120.7, 131.0, 135.8, 135.9, 138.9, 160.2, 168.2, 168.9; HRMS (EI) calcd for C₁₉H₂₃NO₄ (M⁺) 329.1627, found 329.1619.

4.1.10. (±)-trans-1-Benzyl-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]pyrrolidine-2,5-dione (**15**)

A solution of **14** (0.16 g, 0.4 mmol) in THF (5.0 mL) was introduced into the pre-dissolved solution of Ni(OAc)₂·4H₂O (0.10 g, 0.4 mmol) in MeOH (5.0 mL) and stirred for 30 min at room temperature. To the resulting mixture was added one

portion of NaBH₄ (29 mg, 0.8 mmol) and stirred for 30 min, followed by the other portion of NaBH₄ (29 mg, 0.8 mmol) at room temperature. The suspension was allowed to stir for 4 h and then quenched with saturated aqueous NH₄Cl, which was extracted with CH₂Cl₂ (3×20.0 mL). After the removal of solvent, the yellow residue was subjected to column chromatography (SiO₂, 10:1:0.005 hexane/diethyl ether/AcOH) to provide **15** (0.10 g, 65%) as a light yellow oil. ¹H NMR (CDCl₃) δ 0.73 (d, *J*=6.0 Hz, 3H), 0.81 (d, *J*=6.0 Hz, 3H), 1.49–1.73 (m, 1H), 1.49–1.73 (m, 1H), 1.73 (s, 3H), 1.79 (s, 3H), 1.84 (sep, *J*=6.0 Hz, 1H), 2.91 (td, *J*=5.4, 4.8 Hz, 1H), 3.55 (d, *J*=4.8 Hz, 1H), 4.48 (d, *J*=6.9 Hz, 2H), 4.70 (ABq, *J*=14.4 Hz, 2H), 5.47 (br t, *J*=6.9 Hz, 1H), 6.86 (d, *J*=8.7 Hz, 2H), 7.03 (d, *J*=8.7 Hz, 2H), 7.27–7.39 (m, 5H); ¹³C NMR [(CD₃)₂CO] δ 17.4, 20.8, 22.6, 25.0, 25.3, 40.3, 42.0, 46.9, 52.3, 64.5, 114.8, 120.2, 127.5, 128.0, 128.5, 129.2, 129.9, 136.7, 136.8, 158.4, 176.8, 178.5; MS (EI) *m/z* 405 (M⁺, 4), 337 (16), 281 (100), 203 (27), 133 (99), 91 (56), 69 (27); HRMS (EI) calcd for C₂₆H₃₁NO₃ (M⁺) 405.2304, found 405.2300.

4.1.11. N-Benzoyloxy-3-isobutyl-4-[4-(3-methylbutenyloxy)phenyl]-1H-pyrrol-2,5-dione (**16**)

To a mixture of **3** (0.15 g, 0.5 mmol) and K₂CO₃ (0.19 g, 1.4 mmol) in acetone (5.0 mL) was added BnBr (60.0 μL, 0.5 mmol) at room temperature, and then refluxed at 65 °C for 12 h. After cooling, the solution was evaporated in vacuo, and the brown solid was subjected to flash chromatography (SiO₂, 1:15 EtOAc/hexane) to give **16** (0.19 g, 0.5 mmol) in 100% yield. ¹H NMR (CDCl₃) δ 0.88 (d, *J*=7.2 Hz, 6H), 1.75 (s, 3H), 1.81 (s, 3H), 2.00 (sep, *J*=7.2 Hz, 1H), 2.47 (d, *J*=7.2 Hz, 2H), 4.55 (d, *J*=6.6 Hz, 2H), 5.15 (s, 2H), 5.49 (br t, *J*=6.6 Hz, 1H), 6.97 (d, *J*=8.7 Hz, 2H), 7.48 (d, *J*=8.7 Hz, 2H), 7.36–7.53 (m, 5H); ¹³C NMR (CDCl₃) δ 18.2, 22.6, 25.8, 28.0, 32.9, 64.8, 79.5, 114.8, 119.1, 120.8, 128.5, 129.2, 129.8, 130.9, 134.0, 135.4, 135.7, 138.7, 160.1, 167.0, 167.7; HRMS (EI) calcd for C₂₆H₂₉NO₄ (M⁺) 419.2097, found 419.2099.

4.1.12. (±)-trans-3-Isobutyl-4-[4-(3-methylbutenyloxy)phenyl]pyrrolidine-2,5-dione (**17**)

A solution of **2** (0.25 g, 0.8 mmol) in THF (8.0 mL) was introduced into the pre-dissolved solution of Ni(OAc)₂·4H₂O (0.20 g, 0.8 mmol) in MeOH (8.0 mL) and stirred for 30 min at ambient temperature. To the resulting mixture was added one portion of NaBH₄ (58 mg, 1.6 mmol) and stirred for 30 min, followed by the other portion of NaBH₄ (58 mg, 1.6 mmol) at ambient temperature. The suspension was allowed to stir for 4 h and then quenched with saturated aqueous NH₄Cl, which was extracted with CH₂Cl₂ (3×20.0 mL). After the removal of solvent, the yellow residue was subjected to column chromatography (SiO₂, 1:3 EtOAc/cyclohexane) to provide **17** (0.19 g, 78%, *cis/trans* 1:2) as a light yellow oil. ¹H NMR (CDCl₃) δ 0.73 (d, *J*=6.3 Hz, 3H), 0.91 (d, *J*=6.3 Hz, 3H), 1.53–1.89 (m, 1H), 1.53–1.89 (m, 1H), 1.74 (s, 3H), 1.77 (s, 3H), 1.86 (sep, *J*=6.3 Hz, 1H), 2.98 (td, *J*=5.4, 4.8 Hz, 1H), 3.60 (d, *J*=4.8 Hz, 1H), 4.49 (d, *J*=6.9 Hz, 2H), 5.49 (br t, *J*=6.9 Hz, 1H), 6.90 (d, *J*=8.7 Hz, 2H), 7.12 (d, *J*=8.7 Hz, 2H), 8.86 (br s, 1H); ¹³C NMR (CDCl₃) δ 18.2, 21.3, 23.0,

25.5, 25.8, 40.5, 48.2, 53.7, 64.7, 115.3, 119.4, 128.5, 128.7, 138.4, 158.5, 178.1, 179.9; HRMS (EI) calcd for $C_{19}H_{25}NO_3$ (M^+) 315.1834, found 315.1837.

4.1.13. ($\pm$)-cis-3-Isobutyl-4-[4-(3-methylbutenyloxy)-phenyl]pyrrolidine-2,5-dione (17**)**

1H NMR ($CDCl_3$) δ 0.67 (d, $J=6.3$ Hz, 3H), 0.81 (d, $J=6.3$ Hz, 3H), 1.47–1.52 (m, 1H), 1.47–1.52 (m, 1H), 1.74 (s, 3H), 1.79 (s, 3H), 1.79 (sep, $J=6.3$ Hz, 1H), 3.13 (dt, $J=9.0$, 5.4 Hz, 1H), 4.11 (d, $J=9.0$ Hz, 1H), 4.49 (d, $J=6.6$ Hz, 2H), 5.48 (br t, $J=6.6$ Hz, 1H), 6.88 (d, $J=8.7$ Hz, 2H), 7.02 (d, $J=8.7$ Hz, 2H), 9.04 (br s, 1H); ^{13}C NMR ($CDCl_3$) δ 18.1, 21.7, 22.5, 25.3, 25.8, 35.2, 44.1, 51.4, 64.8, 115.0, 119.4, 125.8, 130.0, 138.3, 158.4, 178.6, 180.4; MS (EI) m/z 315 (M^+ , 1), 247 (26), 191 (100), 176 (14), 133 (84), 69 (30); HRMS (EI) calcd for $C_{19}H_{25}NO_3$ (M^+) 315.1834, found 315.1833.

4.1.14. ($\pm$)-trans-1-(tert-Butoxycarbonyl)-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]pyrrolidine-2,5-dione (18**)**

To a mixture of **17** (0.19 g, 0.6 mmol) and DMAP (7.3 mg, 0.06 mmol) in CH_3CN was added Boc_2O (0.14 g, 0.7 mmol) dropwise at $0^\circ C$. Stirring continued at $0^\circ C$ for 30 min and at room temperature for 1 h. Evaporation and chromatography (SiO_2 , 1:9 EtOAc/hexane) gave **18** (0.23 g, 94%) as a yellow oil. 1H NMR ($CDCl_3$) δ 0.74 (d, $J=6.3$ Hz, 3H), 0.90 (d, $J=6.3$ Hz, 3H), 1.48–1.65 (m, 1H), 1.58 (s, 9H), 1.74 (s, 3H), 1.77 (s, 3H), 1.80–1.91 (m, 1H), 1.80–1.91 (m, 1H), 2.99 (m, 1H), 3.60 (d, $J=6.0$ Hz, 1H), 4.50 (d, $J=6.6$ Hz, 2H), 5.48 (br t, $J=6.6$ Hz, 1H), 6.91 (d, $J=8.7$ Hz, 2H), 7.13 (d, $J=8.7$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 18.2, 21.4, 22.9, 25.5, 25.8, 27.7, 40.4, 47.0, 52.9, 64.8, 86.2, 115.3, 119.4, 127.9, 128.9, 138.4, 146.5, 158.6, 173.3, 175.2; HRMS (EI) calcd for $C_{24}H_{23}NO_5$ (M^+) 415.2359, found 415.2360.

4.1.15. ($\pm$)-trans-1-Hydroxy-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]pyrrolidine-2,5-dione (5**)**

To a solution of **18** (0.17 g, 0.4 mmol) in CH_3CN was added an excess (50 wt %) aqueous solution of NH_2OH (0.2 mL, 3.4 mmol) at ambient temperature. The mixture was allowed to stir for 1 h, and followed by removal of the solvent in vacuo and the resulting solid was subjected to column chromatography (SiO_2 , 6:1 CH_2Cl_2 /EtOAc) to obtain **5** (0.10 g, 77%) as a colorless oil. 1H NMR ($CDCl_3$) δ 0.69 (d, $J=6.0$ Hz, 3H), 0.89 (d, $J=6.0$ Hz, 3H), 1.51 (m, 1H), 1.73 (s, 3H), 1.79 (s, 3H), 1.72–1.84 (m, 1H), 1.72–1.84 (m, 1H), 2.87 (m, 1H), 3.52 (d, $J=4.2$ Hz, 1H), 4.47 (d, $J=6.6$ Hz, 2H), 5.47 (br t, $J=6.6$ Hz, 1H), 6.87 (d, $J=8.4$ Hz, 2H), 7.08 (d, $J=8.4$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 18.2, 21.3, 23.0, 25.3, 25.8, 40.3, 44.6, 49.8, 64.7, 115.3, 119.3, 128.0, 128.9, 138.4, 158.6, 173.4, 175.1; HRMS (EI) calcd for $C_{19}H_{25}NO_4$ (M^+) 331.1784, found 331.1792.

4.1.16. ($\pm$)-cis-1-Hydroxy-3-isobutyl-4-[4-(3-methyl-2-butenyloxy)phenyl]pyrrolidine-2,5-dione (4**)**

To a solution of **3** (0.13 g, 0.4 mmol) in 5.0 mL of acetic acid was added Zn dust (3.9 mmol) in one portion. The mixture was stirred at room temperature for 2 h, quenched by addition of

water (2.0 mL), and extracted with CH_2Cl_2 (3×5.0 mL). After removal of the solvent, the residue was subjected to column chromatography (SiO_2 , 1:2 EtOAc/cyclohexane) to provide **5** (0.035 g, 27%) and **4** (0.070 g, 53%) as a colorless oil. 1H NMR ($CDCl_3$) δ 0.67 (d, $J=6.0$ Hz, 3H), 0.81 (d, $J=6.0$ Hz, 3H), 1.03 (m, 1H), 1.44–1.48 (m, 1H), 1.44–1.48 (m, 1H), 1.73 (s, 3H), 1.79 (s, 3H), 3.09 (m, 1H), 4.08 (d, $J=8.4$ Hz, 1H), 4.48 (d, $J=6.6$ Hz, 2H), 5.47 (br t, $J=6.6$ Hz, 1H), 6.85 (d, $J=8.4$ Hz, 2H), 6.97 (d, $J=8.4$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 18.2, 21.7, 22.4, 25.2, 25.8, 35.3, 40.2, 47.4, 64.8, 115.0, 119.3, 125.3, 130.2, 138.3, 158.6, 174.1, 175.5; HRMS (EI) calcd for $C_{19}H_{25}NO_4$ (M^+) 331.1784, found 331.1791.

4.1.17. N-Benzyl-3-benzyl-4-[4-(3-methyl-2-butenyloxy)-phenyl]maleimide (19**)**

This compound was prepared as a colorless oil from **11** according to the procedure of Kelly.^{8f} 1H NMR ($CDCl_3$) δ 1.74 (s, 3H), 1.80 (s, 3H), 3.95 (s, 2H), 4.53 (d, $J=6.6$ Hz, 2H), 4.72 (s, 2H), 5.48 (br t, $J=6.6$ Hz, 1H), 6.94 (d, $J=9.0$ Hz, 2H), 7.53 (d, $J=9.0$ Hz, 2H), 7.14–7.41 (m, 5H), 7.14–7.41 (m, 5H); ^{13}C NMR ($CDCl_3$) δ 18.2, 25.8, 29.9, 41.8, 64.8, 114.8, 119.1, 121.0, 126.7, 127.7, 128.4, 128.5, 128.6, 128.8, 131.1, 135.9, 136.5, 137.3, 138.2, 138.7, 160.2, 171.0, 171.6; HRMS (EI) calcd for $C_{29}H_{27}NO_3$ (M^+) 437.1991, found 437.1997.

4.1.18. 3-Benzyl-4-[4-(3-methyl-2-butenyloxy)phenyl]furan-2,5-dione (6**)**

A mixture of benzyl maleimide **19** (0.21 g, 0.5 mmol) in THF/MeOH (20.0 mL/20.0 mL), water (20.0 mL), and KOH (0.23 g) was refluxed for 3 h. After it was poured into water, acidified with 1 N HCl, and then extracted with EtOAc, the organic phase was washed with saturated aqueous $NaHCO_3$ and brine and then was dried over $MgSO_4$. After removal of the solvent, the residue was purified by flash chromatography (SiO_2 , 1:1 CH_2Cl_2 /hexane) to provide **6** (0.14 g, 81%) as an oil. 1H NMR ($CDCl_3$) δ 1.75 (s, 3H), 1.81 (s, 3H), 4.03 (s, 2H), 4.57 (d, $J=6.9$ Hz, 2H), 5.48 (br t, $J=6.9$ Hz, 1H), 6.99 (d, $J=9.0$ Hz, 2H), 7.63 (d, $J=9.0$ Hz, 2H), 7.21 (d, $J=7.4$ Hz, 2H), 7.25–7.35 (m, 3H); ^{13}C NMR ($CDCl_3$) δ 18.2, 25.8, 30.5, 65.0, 115.2, 118.8, 119.5, 127.3, 128.3, 129.1, 131.3, 135.7, 137.4, 139.0, 140.6, 161.3, 165.3, 166.2; HRMS (EI) calcd for $C_{22}H_{20}O_4$ (M^+) 348.1362, found 348.1366.

4.1.19. 1-Hydroxy-3-benzyl-4-[4-(3-methyl-2-butenyloxy)-phenyl]-1H-pyrrol-2,5-dione (7**)**

A mixture of furandione **6** (0.16 g, 0.5 mmol) and hydroxylamine hydrochloride (0.13 g, 1.8 mmol) in pyridine (10.0 mL) was heated at reflux for 2 h. After cooling, the solvent was removed in vacuo and the resulting solid was subjected to column chromatography (SiO_2 , 1:3 EtOAc/hexane) to obtain **7** (0.14 g, 84%) as a colorless oil. 1H NMR ($CDCl_3$) δ 1.75 (s, 3H), 1.80 (s, 3H), 3.94 (s, 2H), 4.54 (d, $J=6.6$ Hz, 2H), 5.48 (br t, $J=6.6$ Hz, 1H), 6.96 (d, $J=9.0$ Hz, 2H), 7.54 (d, $J=9.0$ Hz, 2H), 7.16–7.30 (m, 5H), 8.15 (br s, 1H); ^{13}C NMR ($CDCl_3$) δ 18.2, 25.8, 29.9, 64.9, 115.0, 119.0, 120.3, 126.9, 128.3, 128.9, 131.1, 133.5, 136.2, 136.7, 138.8, 160.6, 167.9,

168.5; HRMS (EI) calcd for $C_{22}H_{21}NO_4$ (M^+) 363.1471, found 363.1469.

4.1.20. ($\pm$)-cis-1-Hydroxy-3-benzyl-4-[4-(3-methyl-2-butenyloxy)phenyl]pyrrolidine-2,5-dione (8**)**

To a solution of **7** (0.079 g, 0.2 mmol) in 4.0 mL of acetic acid was added Zn dust (1.6 mmol) in one portion. The mixture was stirred at room temperature for 2 h, quenched by addition of water (1.0 mL), and extracted with CH_2Cl_2 (3×5.0 mL). After removal of the solvent, the residue was subjected to column chromatography (SiO_2 , 1:2 EtOAc/cyclohexane) to provide the single diastereomer **8** (0.045 g, 57%) as a colorless oil. 1H NMR ($CDCl_3$) δ 1.59 (s, 3H), 1.62 (s, 3H), 2.43 (dd, $J=15.0$, 10.5 Hz, 1H), 3.18 (dd, $J=15.0$, 4.5 Hz, 1H), 3.48 (m, 1H), 4.02 (d, $J=8.7$ Hz, 1H), 4.47 (d, $J=6.9$ Hz, 2H), 5.47 (br t, $J=6.9$ Hz, 1H), 6.70–6.76 (br s, 2H), 6.84 (br s, 4H), 7.08–7.21 (br s, 3H); ^{13}C NMR ($CDCl_3$) δ 18.2, 25.8, 31.6, 43.7, 47.0, 64.8, 114.9, 119.3, 124.8, 126.4, 128.2, 128.4, 130.3, 137.4, 138.4, 158.7, 173.4, 174.1; HRMS (EI) calcd for $C_{22}H_{23}NO_4$ (M^+) 365.1627, found 365.1626.

Acknowledgements

We thank the National Science Council of the Republic of China for financial support (Research Grants NSC 94-2113-M-018-003 and 95-2113-M-018-004). We also acknowledge with appreciation to the former High School research participants Mr. Xin-Quan Chen, Miss Pei-Xing Lai, Yi-Chang Lee, Lei-Ming Kuo, Yi-Xun Xie, and M.S. Shun-Yu Lin for pioneer studies.

Supplementary data

Supplementary data associated with this article can be found in the online version, at [doi:10.1016/j.tet.2008.02.077](https://doi.org/10.1016/j.tet.2008.02.077).

References and notes

- (a) Adlington, R. M.; Baldwin, J. E.; Cox, R. J.; Pritchard, G. J. *Synlett* **2002**, 820 and references therein; (b) Nakamura, N.; Hirakawa, A.; Gao, J. J.; Kakuda, H.; Shiro, M.; Komatsu, Y.; Sheu, C. C.; Hattori, M. *J. Nat. Prod.* **2004**, 67, 46.
- Tsai, Z. T.; Liaw, S. L. *The Use and the Effect of Ganoderma*; Sheng-Yun: Taichung, Taiwan, 1982; p 116.
- Tseng, T. H.; Tsheng, Y. M.; Lee, Y. J.; Hsu, H. L. *J. Chin. Chem. Soc.* **2000**, 47, 1165.
- Lin, S. Y.; Chen, C. L.; Lee, Y. J. *J. Org. Chem.* **2003**, 68, 2968.
- (a) Stille, J. K. *Angew. Chem., Int. Ed. Engl.* **1986**, 25, 508; (b) Stille, J. K.; Groh, B. L. *J. Am. Chem. Soc.* **1987**, 109, 813; (c) Echavarren, A. M.; Stille, J. K. *J. Am. Chem. Soc.* **1987**, 109, 5478.
- (a) Baag, M. M.; Argade, N. P. *Synthesis* **2006**, 1005; (b) Stewart, S. G.; Polomska, M. E.; Lim, R. W. *Tetrahedron Lett.* **2007**, 48, 2241.
- Bernard, A. M.; Cocco, M. T.; Onnis, V.; Piras, P. P. *Synthesis* **1998**, 3, 256.
- (a) Kelly, T. R.; Li, Q.; Bhushan, V. *Tetrahedron Lett.* **1990**, 31, 161; (b) Kelly, T. R.; Jagoe, C. T. *Tetrahedron Lett.* **1991**, 32, 4263; (c) Kelly, T. R.; Lang, F. *Tetrahedron Lett.* **1995**, 36, 5319; (d) Kelly, T. R.; Lang, F. *J. Org. Chem.* **1996**, 61, 4623; (e) Kelly, T. R.; Xu, W.; Ma, Z.; Li, Q.; Bhushan, V. *J. Am. Chem. Soc.* **1993**, 115, 5843; (f) Kelly, T. R.; Lee, Y. J.; Mears, R. J. *J. Org. Chem.* **1997**, 62, 2774.
- (a) Lutz, R. E. *J. Am. Chem. Soc.* **1930**, 52, 3423; (b) Brown, J. P. *Chem. Abstr.* **1966**, 65, 2131a; (c) Shoppee, C. W.; Wu, W. Y. *Aust. J. Chem.* **1987**, 40, 1137; (d) Horowska, B.; Mazerska, Z.; Ledochowski, A.; Cristalli, G.; Franchetti, P.; Martelli, S. *Eur. J. Med. Chem. Chim. Ther.* **1988**, 23, 91; (e) Choi, D. S.; Huang, S.; Huang, M.; Barnard, T. S.; Adams, R. D.; Seminario, J. M.; Tour, J. M. *J. Org. Chem.* **1998**, 63, 2646; (f) Carman, R. M.; Derbyshire, R. P. C.; Hansford, K. A.; Kadirvelraj, R.; Robinson, W. T. *Aust. J. Chem.* **2001**, 54, 117; (g) Dubernet, M.; Caubert, V.; Guillard, J.; Viaud-Massuard, M.-C. *Tetrahedron* **2005**, 61, 4585.
- Gemal, A. L.; Luche, J. L. *J. Am. Chem. Soc.* **1981**, 103, 5454.
- Comins, D. L.; Brooks, C. A.; Ingalls, C. L. *J. Org. Chem.* **2001**, 66, 2181 and references therein.
- (a) Selles, P. *Org. Lett.* **2005**, 7, 605; (b) Aqueveque, P.; Anke, T.; Sterner, O. *Z. Naturforsch.* **2002**, 57c, 257.
- Autoclave, a PTFE-lined vessel No. 4749, was purchased from Parr Instrument Company, USA.
- Joyce, R. P.; Gainor, J. A.; Weinreb, S. M. *J. Org. Chem.* **1987**, 52, 1177.