



Oxidative coupling of indoles using thallium(III) trifluoroacetate

Paul A. Keller^{a,*}, Nageshwar R. Yepuri^a, Michael J. Kelso^a, Michael Mariani^a,
Brian W. Skelton^b, Allan H. White^b

^aSchool of Chemistry, University of Wollongong, Wollongong, NSW 2522, Australia

^bSchool of Biomedical, Biomolecular and Chemical Sciences, University of Western Australia, Crawley WA 6009, Australia

ARTICLE INFO

Article history:

Received 22 March 2008

Received in revised form 12 May 2008

Accepted 30 May 2008

Available online 5 June 2008

ABSTRACT

The oxidative coupling of polysubstituted electron-rich indoles mediated by thallium trifluoroacetate was found to be a facile, clean, and high yielding reaction. Indolic coupling sites were determined by the nature of the substituents present, with dimerisation at the indole 2-position being the dominant outcome. Indoles bearing two potential reaction sites with similar reactivity were additionally found to undergo heterocoupling.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Numerous synthetic methods are available for introducing the biaryl linkage into organic compounds.^{1,2} The popular Suzuki reaction and its related organometallic cousins employ specific functionality to direct the sites of reaction and many examples have of course been reported.³ Also well known are oxidative coupling methods,⁴ where biaryl bonds are formed directly at unsubstituted aryl sites, which have been activated either by the aryl units themselves or by ring substituents. The clear advantage of oxidative coupling over organometallic methods, particularly for the formation of biaryl dimers is that it requires no prior functionalisation of the aromatic reaction site(s).¹

As part of our ongoing investigations into the biological activities of homo- and hetero-dimeric aromatic systems,⁵ we wished to identify simple methods for constructing focused libraries of symmetrical and non-symmetrical biindoles incorporating biaryl linkages at the indole 2 and 7-positions. While there have been many syntheses of 2,2'-biindoles,⁶ the procedures usually involve numerous steps and are impractical for preparing multiple analogues. In the case of 7,7'-biindoles, very few syntheses have been reported but one notable example⁷ describes the oxidative coupling of 4,6-dimethoxy-2,3-diphenylindole in the presence of quinone, chloranil or dichlorodicyanoquinone to yield the homodimeric 7,7'-coupled biindole in 100%, 70% and 60% yields, respectively.

Attracted by the simplicity of oxidative coupling methods, we sought to identify a suitable oxidant for preparing our biindole libraries. Thallium(III) trifluoroacetate has previously been used for

effecting oxidative dimerisations of electron-rich naphthyl compounds,⁸ 2-substituted indoles⁹ and indolocarbazoles.¹⁰ We now wish to report that thallium(III) trifluoroacetate, in combination with BF₃·Et₂O, is an effective reagent for the oxidative coupling of electron-rich polysubstituted indoles.

2. Results and discussion

2.1. Synthesis of 7,7'-biindoles


Our preliminary investigations focussed on the 7,7'-dimerisation of a model compound 2,3-dicarbomethoxy-4,6-dimethoxyindole **1**. This indole monomer had its most reactive positions (C2 and C3) blocked, allowing the two methoxy substituents to activate the C7-position to oxidative dimerisation. The use of thallium trifluoroacetate/BF₃·Et₂O as oxidant afforded the 7,7'-dimer **2** as the sole product in excellent yield (Scheme 1). This rapid and selective reaction was superior to the previously reported oxidation, which employed ICl and produced **2** in 34% yield along with 31% of the iodinated monomer **3**.⁷

2.2. Synthesis of 2,2'-biindoles


Formation of 2,2'-biindoles by oxidative dimerisation of 3-aryl-4,6-dimethoxyindoles using thallium trifluoroacetate/BF₃·Et₂O was explored next. These monomers had two potential dimerisation sites (C2 and C7) but initial experiments showed that indole N-protection limited dimerisation at the 7-position (data not shown). The substituted indole monomers **4**–**11** were NH protected using sodium hydride and benzenesulfonyl chloride to give **8**–**11** after recrystallisation from methanol (Scheme 2). Clean formation of the 2,2'-dimers was achieved by treating an acetonitrile solution of the protected indole monomers with a half-mole equivalent of

* Corresponding author. Tel.: +61 2 4221 4692; fax: +61 2 4221 4287.

E-mail address: keller@uow.edu.au (P.A. Keller).



Scheme 1.



X	Yield		
	monomer	dimer	deprotected dimer
<i>p</i> -MeOC ₆ H ₄	65% 8	70% 12	15% 16
Ph	70% 9	70% 13	15% 17
<i>p</i> -BrC ₆ H ₄	65% 10	74% 14	10% 18
<i>p</i> -NO ₂ C ₆ H ₄	65% 11	0% 15	N/A

Scheme 2.

thallium(III) trifluoroacetate, followed by addition of excess BF₃·Et₂O. After 1 h of stirring at room temperature, the dimers were isolated and recrystallised from methanol/dichloromethane. Compound **11** bearing the strongly electron-withdrawing *p*-NO₂C₆H₄ at the indole 3-position did not couple under these conditions, even after prolonged reaction.


As a representative example, the structure of **12** was supported by analysis of the mass spectrum (ES), which showed a peak at *m/z* 845 assigned as the [M+1] ion. Further support was provided by the ¹H NMR spectra, which showed the disappearance of a peak at 7.31 ppm assigned to H2 in **8**. The presence of only 'one set' of peaks in the ¹H NMR spectrum was consistent with the formation of a symmetrical dimer. Comparison of the ¹H NMR spectrum of **12** with that of the monomer **8** showed a shift in the aromatic peaks assigned to the benzenesulfonyl group from 7.89–7.56 to 7.53–7.30 ppm, and those assigned to the 3-aryl ring from 7.47–6.92 to 7.03–6.65 ppm. The structure of **12** was eventually confirmed by a single crystal X-ray study (Fig. 1), which indicated that both phenyl rings are aligned to one side of the molecule, accounting for the downfield shifts of the aromatic signals in the NMR spectra of the dimer **12** relative to the monomer **8**. The 2,2'-dimeric indoles


12–15 can exist as atropoisomers due to the presence of four *ortho* substituents. Presumably both of these isomers were formed in each of the products as racemic mixtures.

Based on a mechanism previously proposed for thallium(III) promoted oxidative dimerisations,⁸ we postulate that oxidative 2,2'-dimerisation of these indoles proceeds via an initial one-electron transfer from the electron-rich indole substrate to thallium(III) to form an indole radical cation (Scheme 3). Subsequent electrophilic substitution with a second indole substrate, followed by oxidative aromatisation, gives the symmetrical 2,2'-dimers.

Initial attempts to remove the benzenesulfonyl protecting group from the dimers using thionyl chloride at room temperature yielded complex mixtures. Deprotection was eventually achieved, albeit in disappointing yields of 10–15%, by heating the dimers at reflux for 4 h in a 20% solution of sodium hydroxide. Dimer **17** was spectroscopically identical to the reported compound that had previously been synthesised in only 1.5% yield.¹²

The poor outcome in the final deprotection step led us to investigate indole 2,2'-dimerisations using alternative protecting groups. A selection of 3-arylsubstituted indoles were *N*-protected with various aryl or alkyl sulfonates and then subjected to the dimerisation/deprotection sequence. These results are summarised in Scheme 4. All of the protection and dimerisation reactions proceeded smoothly and in good yield with the exception of *N*-4-nitrobenzenesulfonyl-3-(*p*-nitrophenyl)indole **21**, which produced no dimer and quantitatively returned the starting material even after prolonged reaction times. This was not unexpected given the failure of compound **11** to undergo dimerisation. Confirmation of


Figure 1. Molecular (ORTEP) projection of **12** down the quasi-two-fold axis.



Scheme 3.


the structure of **28** was provided from a single crystal X-ray study (Fig. 2).

Different reaction conditions were trialled for deprotection of the dimers using thiophenol, DMF and K_2CO_3 at room temperature. Only **29** gave the deprotected 2,2'-biindole **16** in 25% yield, a slight improvement over the 15% obtained from **12** by heating at reflux in



R	X	Yield		
		monomer	dimer	deprotected dimer
<i>p</i> -Me $C_6H_4SO_2$	<i>p</i> -Br C_6H_4	70% 19	70% 25	Complex mixture
<i>p</i> -NO ₂ $C_6H_4SO_2$	<i>p</i> -Br C_6H_4	65% 20	70% 26	Complex mixture
<i>p</i> -NO ₂ $C_6H_4SO_2$	<i>p</i> -NO ₂ C_6H_4	65% 21	NR	-
MeSO ₂	Ph	75% 22	70% 27	Complex mixture
MeSO ₂	<i>p</i> -Br C_6H_4	70% 23	75% 28	Complex mixture
MeSO ₂	<i>p</i> -OMe C_6H_4	75% 24	70% 29	25% 16

Scheme 4.

Figure 2. Molecular (ORTEP) projection of **28** down the crystallographic two-fold axis.

NaOH (Scheme 2). All other deprotection reactions again resulted in complex mixtures.


2.3. Oxidative dimerisation with *N*-acyl protected indoles

The effects of indole *N*-protection were further explored using acetyl, trifluoroacetyl and Boc groups. The results from the protection, dimerisation and deprotection reactions are summarised in Scheme 5.

Indole acetylation was achieved using tetrabutylammonium hexafluorophosphate with potassium hydroxide and CH_2Cl_2 under phase transfer conditions. Trifluoroacetyl protection of the indoles was carried out using a modified literature protocol.¹¹ Oxidative dimerisation was attempted on compounds **30–34** using the conditions described above. In stark contrast to monomers **8** and **24** (which contain 3-*p*-methoxyphenyl indolic substituents and *N*-phenylsulfonyl and *N*-methylsulfonyl groups, respectively) the analogous *N*-acetyl and *N*-trifluoroacetyl-protected indoles **30** and **32** yielded no 2,2'-biindoles. Inexplicably, the corresponding *N*-Boc derivative **34** produced a complex mixture of products. Furthermore, given that compounds **11** and **21** (which carry 3-*p*-nitrophenyl indolic substituents) failed to undergo dimerisation, it was very surprising to find that the *N*-acetyl protected 3-(*p*-nitrophenyl)indole **31** dimerised to **35** in 30% yield! What was perhaps even more surprising was the finding that the 3-(*p*-nitrophenyl)indole **33** gave no trace of the 2,2-dimer but instead produced the 2,7',7''-terindole **36** in 35% yield (Scheme 5). The remaining mass was identified as the starting indole **33** and its *N*-deprotected precursor.

Support for the structure of the trimer **36** was provided by its mass spectrum (ES), which showed a peak at 1179 corresponding to the $M+1$ ion. Analysis of the 1H NMR spectrum showed singlet peaks at 6.73, 6.66 and 6.44 ppm corresponding to the H5', H5 and H5'' protons. These three protons showed no *meta*-coupling to their respective H7 protons (usually observed), suggesting the molecule contains 2,7'- and 7,7''-linkages. Additionally, one H2 signal was absent from the 1H NMR spectrum while there were clearly two other singlets at 7.21 and 6.95 ppm corresponding to the H2' and H2'' protons. The proposed structure was fully supported by gHMBC, gCOSY and NOESY correlations.

The mechanism for the formation of **36** is not clear, although on the basis of current knowledge about TFA promoted oxidations and dimerisation of indoles,^{8,13} it may be due to the *p*-nitro group and trifluoroacetate group at the N1-position not activating the 2-position to the same level as occurs in the examples that underwent coupling. This would allow reaction at the C7-position to



R	X	Yield		
		monomer	dimer	trimer
COCH_3	<i>p</i> -OMePh	65% 30	NR	NR
COCH_3	<i>p</i> -NO ₂ Ph	57% 31	2,2'- dimer 30% 35	-
COCF_3	<i>p</i> -OMePh	60% 32	NR	NR
COCF_3	<i>p</i> -NO ₂ Ph	60% 33	-	2,7',7,7''- trimer 35% 36
Boc	<i>p</i> -OMePh	99% 34	-	-


Scheme 5.

become a significant process. The 2,7'-dimer could then undergo further oxidative coupling between the 7 and 7'-positions yielding the 2,7',7,7''-trimer **36**.

2.4. Thallium(III) trifluoroacetate mediated coupling of 4,6-dimethoxy-3-aryl indoles without N-protection

The oxidative dimerisation of indoles **4**, **6** and **7** using thallium(III) trifluoroacetate was attempted in the absence of NH protecting groups. After preliminary optimisation reactions, it was found that 3–5 equiv of the Lewis acid were best added at 0°C before continued stirring at room temperature for 3 h. The results are summarised in Scheme 6. Indole **4** gave the 2,2'-dimer **16** in 35% yield, the remainder being starting material and other baseline products. Indole **6** gave 20% of 2,2'-dimer **18** with 50% of the

starting material recovered. Interestingly, 4,6-dimethoxy-3-(*p*-nitrophenyl)indole **6** gave only the 2,7'-dimer **37** (20%) with no traces of formation of 2,2'-dimer.



X	Yield	
	2,2'-dimer	2,7'-dimer
<i>p</i> -OMePh	35% 16	-
<i>p</i> -BrPh	20% 18	-
<i>p</i> -NO ₂ Ph	-	15% 37

Scheme 6.

For all oxidative coupling reactions attempted without *N*-protecting groups the reactions took longer and returned significant quantities of starting material, suggesting that dimerisation is facilitated by *N*-protection. The results also suggest that the alkyl or arylsulfonyl groups may help in the regioselectivity of the 2,2'-dimerisation provided the 3-aryl substituent is mildly electron-withdrawing or electron-donating.

3. Conclusions

Oxidative coupling of indoles with thallium(III) trifluoroacetate provides good to excellent yields of 2,2'-dimers. Protecting groups such as benzenesulfonyl, tosyl or methanesulfonyl aid formation of the 2,2'-dimer, provided that the *para* substituent on the 3-aryl ring is mildly deactivating or an electron-donating group. When the 3-aryl ring bears a *para* substituent, which is a strong electron-withdrawing group, e.g., NO_2 , no reaction is observed unless an *N*-acetyl protecting group is present, which leads to 2,7'-dimerisation. Changing the indole *N*-protecting group to a trifluoroacetate group deactivates C2 and activates C7 such that trimeric-indoles are formed.

4. Experimental

4.1. General

Melting point determinations were carried out on a Gallenkamp melting point apparatus. Chemical ionisation (CI) and electron impact (EI) mass spectra were obtained on a Shimadzu QP-5000 mass spectrometer by a direct insertion technique with an electron beam energy of 70 eV. Electrospray (ESI) mass spectra were obtained on a VG Autospec spectrometer. High-resolution mass spectra (HRMS) were determined on a micromass QToF2 spectrometer using polyethylene glycol or polypropylene glycol as the internal standard. The m/z values are stated with their peak

intensity as a percentage in parentheses. Proton and carbon nuclear magnetic resonance (NMR) spectra were obtained as specified on a Varian Mercury 300 MHz or Varian Inova 500 MHz spectrometer. Spectra were recorded in the specified deuterated solvent, and referenced to the residual non-deuterated solvent signal. Chemical shifts (δ) in parts per million were measured relative to the internal standard. Analytical thin layer chromatography (TLC) was carried out on Merck silica gel 60 F₂₅₄ pre-coated aluminium plates with a thickness of 0.2 mm. All column chromatography was performed under 'flash' conditions on Merck silica gel 60 (230–400 mesh). Chromatography solvent mixtures were measured by volume. Sodium hydride was used as a dispersion in oil. Organic solvent extracts were dried with anhydrous magnesium sulfate. All compounds were judged to be of greater than 95% purity based upon ¹H NMR and TLC analysis.

4.1.1. 7,7'-Bi(4,6-dimethoxyindolyl)-2,2',3,3'-tetracarboxylate (**2**)

To a solution of dimethyl 4,6-dimethoxyindole-2,3-dicarboxylate **1** (460 mg, 1.57 mmol) and thallium trifluoroacetate (432 mg, 0.79 mmol) in dry acetonitrile (40 mL) was added BF₃·(CH₃CH₂)₂O (785 μ L, 0.79 mmol) and the reaction allowed to stir at room temperature for 40 min. The reaction was then extracted with water (100 mL) and CH₂Cl₂ (4×50 mL). The combined organic extracts were dried (MgSO₄) and concentrated and the residue subjected to flash silica gel column chromatography (5:1 CH₂Cl₂/EtOAc) to yield the dimer **2** (420 mg, 83%) as a pale yellow powder, which was spectroscopically identical to that reported.⁷

4.2. General procedure (A) for the synthesis of protected indole starting materials

To a stirred solution of 4,6-dimethoxy-3-phenylindole (1 mol) in CH₂Cl₂ under a nitrogen atmosphere was added sodium hydride (1 mol, 60% dispersion in oil) and the reaction then stirred for 25 min. The sulfonyl chloride (1 mol) was then added and the reaction left to stir at 25 °C for an additional 2 h. The solvent was then removed under reduced pressure and the resultant residue was suspended in water (50 mL) and extracted with CH₂Cl₂ (3×100 mL). The organic fractions were combined, dried (MgSO₄), concentrated under reduced pressure and the residue recrystallised from methanol yielding the corresponding protected indole.

4.2.1. *N*-Benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)-indole (**8**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-methoxyphenyl)indole **4** (2.00 g, 7.0 mmol), sodium hydride (0.28 g, 7.0 mmol) in 60% oil dispersion) and benzenesulfonyl chloride (1.23 g, 7.0 mmol) to give the indole **8** (2.11 g, 73%) as a white solid, mp 131–133 °C. ¹H NMR δ : 7.89 (2H, d, J =7.5 Hz, sulfonyl ArH2'' and 6''), 7.59–7.41 (3H, m, sulfonyl ArH3'', 4'' and 5''), 7.47 (2H, d, J =8.7 Hz, ArH2' and 6'), 7.31 (1H, s, H2), 7.20 (1H, d, J =1.8 Hz, H7), 6.92 (2H, d, J =8.7 Hz, ArH3' and 5'), 6.32 (1H, d, J =1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.84 (3H, s, C6 OCH₃), 3.71 (3H, s, ArC4 OCH₃). ¹³C NMR δ : 159.5 (ArC4'), 159.0 (C6), 155.0 (C4), 138.5 (C7a), 137.5 (sulfonyl ArC1''), 134.0 (sulfonyl ArC4''), 130.8 (sulfonyl ArC2'' and 6''), 129.4 (sulfonyl ArC3'' and 5''), 126.9 (C3), 124.4 (ArC1'), 121.1 (C2), 113.3 (ArC3' and 5'), 110.0 (C3a), 96.0 (C7), 90.0 (C5), 55.5 (C4 OCH₃), 54.5 (C6 OCH₃), 54.3 (ArC4 OCH₃). CIMS m/z : 424 ([M+1]⁺, 10%), 391 (15), 340 (10), 284 (30), 149 (20), 125 (15), 113 (50), 97 (70), 85 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₂₃H₂₂NO₅ 424.1219; found 424.1230.

4.2.2. *N*-Benzenesulfonyl-4,6-dimethoxy-3-phenylindole (**9**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-phenylindole **5** (2.00 g, 7.9 mmol), sodium hydride

(0.32 g, 8.0 mmol) and benzenesulfonyl chloride (1.40 g, 7.9 mmol) to give the indole **9** (2.14 g, 69%) as a white solid, mp 125–128 °C. ¹H NMR δ : 7.90–7.54 (5H, m, sulfonyl ArH''), 7.46–7.36 (5H, m, ArH'), 7.36 (1H, s, H2), 7.20 (1H, d, J =3.0 Hz, H7), 6.33 (1H, d, J =3.0 Hz, H5), 3.95 (1H, s, C4 OCH₃), 3.65 (1H, s, C6 OCH₃). ¹³C NMR δ : 159.5 (C6), 154.5 (C4), 139.0 (C7a), 138.0 (ArC4'), 134.0 (sulfonyl ArC4''), 129.5 (sulfonyl ArC2'' and 6''), 129.2 (sulfonyl ArC1''), 129.0 (sulfonyl ArC3'' and 5''), 128.0 (ArC4'), 127.5 (ArC2' and 6'), 127.0 (ArC3' and 5'), 124.7 (C3), 122.0 (C2), 112.5 (C3a), 95.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.7 (C6 OCH₃). ES-MS m/z : 394 ([M+1]⁺, 80%), 310 (25), 254 (100), 187 (50), 141 (20); ES-HRMS m/z : calcd for [M+1]⁺ C₂₂H₂₀NO₄S 394.1113; found 394.1108.

4.2.3. *N*-Benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole (**10**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.50 g, 7.5 mmol), sodium hydride (0.30 g, 7.5 mmol, 60% dispersion in oil) and benzenesulfonyl chloride (1.33 g, 7.5 mmol) to give the indole **10** (2.49 g, 70%) as a white solid, mp 137–139 °C. ¹H NMR δ : 7.96 (2H, d, J =8.7 Hz, sulfonyl ArH2'' and 6''), 7.56 (2H, m, sulfonyl ArH3'' and 5''), 7.49–7.54 (4H, m, ArH2', 3', 5' and 6'), 7.25–7.48 (1H, m, sulfonyl ArH4''), 7.34 (1H, s, H2), 7.19 (1H, d, J =1.8 Hz, H7), 6.33 (1H, d, J =1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.71 (3H, s, C6 OCH₃). ¹³C NMR δ : 160.0 (C6), 154.0 (C4), 138.3 (C7a), 138.0 (ArC4'), 134.0 (sulfonyl ArC1''), 133.0 (sulfonyl ArC4''), 131.3 (ArC2' and 6'), 130.9 (ArC3' and 5'), 129.5 (sulfonyl ArC2'' and 6''), 127.0 (sulfonyl ArC3'' and 5''), 123.7 (ArC1''), 121.7 (C2), 121.4 (C3), 112.9 (C3a), 96.0 (C7), 90.0 (C5), 58.5 (C4 OCH₃), 58.0 (C6 OCH₃). CIMS m/z : 474 ([M+1]⁺ ⁸¹Br 20%), 332 (15), 173 (30), 159 (75), 143 (100); EI-HRMS m/z : calcd for C₂₂H₁₈NO₄S⁷⁹Br 471.0139; found 471.0126.

4.2.4. *N*-Benzenesulfonyl-4,6-dimethoxy-3-(4-nitrophenyl)-indole (**11**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-nitrophenyl)indole **7** (2.00 g, 6.7 mmol), sodium hydride (0.27 g 6.7 mmol) and benzenesulfonyl chloride (1.18 g, 6.7 mmol) to give the indole **11** (1.78 g, 61%) as a white solid, mp 150–152 °C. ¹H NMR δ : 8.22 (2H, d, J =9.0 Hz, ArH2' and 6'), 7.93 (2H, d, J =9.0 Hz, ArH3' and 5'), 7.69 (2H, d, J =8.7 Hz, sulfonyl ArH2'' and 6''), 7.59–7.45 (2H, m, sulfonyl ArH3'', 4'' and 5''), 7.19 (1H, d, J =2.1 Hz, H7), 6.36 (1H, d, J =2.1 Hz, H5), 3.90 (3H, s, C4 OCH₃), 3.79 (3H, s, C6 OCH₃). ¹³C NMR δ : 159.8 (C6), 154.6 (C4), 146.9 (sulfonyl ArC1''), 141.1 (sulfonyl ArC1''), 138.1 (C7a), 137.7 (ArC4), 134.4 (sulfonyl ArC4''), 130.3 (ArC2' and 6'), 129.6 (sulfonyl ArC3' and 5'), 127.1 (ArC3' and 5'), 123.1 (sulfonyl ArC2'' and 6''), 122.7 (C2), 112.3 (C3a), 95.6 (C7), 89.9 (C5), 56.6 (C4 OCH₃), 55.4 (C6 OCH₃). ES-MS m/z : 439 ([M+1]⁺, 100%); EI-HRMS m/z : calcd for C₂₂H₁₈N₂O₆S 438.0885; found 438.0907.

4.2.5. 3-(4-Bromophenyl)-4,6-dimethoxy-*N*-(4'-methylphenyl)-sulfonylindole (**19**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.24 g, 6.0 mmol) and *p*-toluenesulfonyl chloride (0.19 g, 6.0 mmol) to give the indole **19** (2.20 g, 76%) as white solid, mp 138–140 °C. ¹H NMR δ : 7.78 (2H, d, J =7.8 Hz, ArH2' and H6'), 7.49 (2H, d, J =9.0 Hz, tosyl ArH3'' and 5''), 7.39 (2H, d, J =9.0 Hz, tosyl ArH2'' and 6''), 7.34 (1H, s, H2), 7.25 (2H, d, J =7.8 Hz, Ar3H' and 5'), 7.18 (1H, d, J =1.8 Hz, H7), 6.32 (1H, d, J =1.8 Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.70 (3H, s, C6 OCH₃), 2.35 (3H, s, CH₃). ¹³C NMR δ : 159.4 (C6), 154.6 (C4), 145.2 (tosyl C1''), 137.5 (C7a), 135.1 (ArC4'), 133.0 (ArC4'), 131.3 (ArC2' and 6'), 130.9 (ArC3' and 5'), 130.1 (tosyl ArC2'' and 6''), 127.0 (tosyl ArC3'' and 5''), 125.0 (tosyl ArC4''), 123.4 (C3), 121.3 (ArC1'), 121.7 (C2), 112.8 (C3a), 95.4 (C7), 90.0 (C5), 56.1 (C4

OCH₃), 55.5 (C6 OCH₃), 22.0 (CH₃). EI-HRMS *m/z*: calcd for C₂₃H₂₀NO₄S⁷⁹Br 485.0296; found 485.0293.

4.2.6. 3-(4-Bromophenyl)-4,6-dimethoxy-N-(*p*-nitrobenzenesulfonyl)indole (**20**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.29 g, 6.0 mmol, 50% dispersion in oil) and *p*-nitrobenzenesulfonyl chloride (1.33 g, 6.0 mmol) to give the indole **20** (2.25 g, 73%) as yellow solid, mp 220–222 °C. ¹H NMR δ: 8.29 (2H, d, *J*=8.7 Hz, Ar2H' and 6'), 8.05 (2H, d, *J*=8.7 Hz, ArH3' and 5'), 7.49 (2H, d, *J*=6.6 Hz, sulfonyl ArH3'' and 5''), 7.37 (2H, d, *J*=6.6 Hz, sulfonyl Ar2H'' and 6''), 7.3 (1H, s, H2), 7.17 (1H, d, *J*=1.8 Hz, H7), 6.36 (1H, d, *J*=1.8 Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.71 (3H, s, C6 OCH₃). ¹³C NMR δ: 160.0 (C6), 155.0 (C4), 151.5 (sulfonyl ArC1'), 143.5 (ArC4'), 138.0 (C7a), 133.0 (ArC1'), 131.3 (ArC2' and 6'), 131.0 (ArC3' and 5'), 128.2 (sulfonyl ArC2'' and 6''), 125.0 (sulfonyl ArC4''), 124.8 (sulfonyl ArC3'' and 5''), 122.0 (C3), 121.1 (C2), 113.4 (C3a), 95.9 (C7), 90.1 (C5), 56.1 (C4 OCH₃), 55.4 (C6 OCH₃). CIMS *m/z*: 519 ([M+1]⁺, ⁸¹Br, 100%), 455 (10), 332 (50), 236 (10); EI-HRMS *m/z*: calcd for C₂₂H₁₇BrN₂O₆S 515.9990; found 515.9982.

4.2.7. 4,6-Dimethoxy-N-(*p*-nitrobenzenesulfonyl)-3-(4-nitrophenyl)indole (**21**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-nitrophenyl)indole **7** (2.00 g, 6.7 mmol) sodium hydride (0.32 g, 6.7 mmol) and *p*-nitrobenzenesulfonyl chloride (1.48 g, 6.7 mmol) to give the indole **21** (2.17 g, 67%) as a pale yellow solid, mp 190–193 °C. ¹H NMR δ: 8.34 (2H, d, *J*=9.0 Hz, ArH2 and 6), 8.24 (2H, d, *J*=8.7 Hz, sulfonyl ArH3 and 5), 8.10 (2H, d, *J*=9.0 Hz, ArH3 and 5), 7.68 (2H, d, *J*=8.7 Hz, sulfonyl ArH2 and 6), 7.26 (1H, s, H2), 7.18 (1H, d, *J*=1.8 Hz, H7), 6.39 (1H, d, *J*=1.8 Hz, H5), 3.91 (3H, s, C4 OCH₃), 3.73 (3H, s, C6 OCH₃). ¹³C NMR δ: 160.1 (C6), 156.0 (C4), 146.1 (ArC4), 144.0 (ArC1), 141.0 (C7a), 148.0 (sulfonyl ArC4), 131.0 (ArC2 and 6), 128.1 (sulfonyl ArC2 and 6), 125.5 (ArC3 and 5), 125.0 (sulfonyl ArC1), 123.0 (C2), 122.0 (ArC2 and 6), 121.0 (C3), 113.0 (C3a), 96.0 (C7), 90.0 (C5), 58.0 (C4 OCH₃), 56.0 (C6 OCH₃). CIMS *m/z*: 484.0 ([M+1]⁺, 100%), 420 (10), 299 (30); EI-HRMS *m/z*: calcd for C₂₂H₁₇N₃O₈S 483.0736; found 483.0746.

4.2.8. N-Methanesulfonyl-4,6-dimethoxy-3-phenylindole (**22**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-phenylindole **5** (2.00 g, 7.9 mmol) sodium hydride (0.32 g, 8.0 mmol) and methanesulfonyl chloride (0.90 g, 7.9 mmol) to give the indole **23** (1.96 g, 75%) as white solid, mp 140–142 °C. ¹H NMR δ: 7.57 (2H, d, *J*=8.4 Hz, ArH2' and 6'), 7.38–7.34 (3H, m, ArH3', 4' and 5'), 7.23 (1H, s, H2), 7.10 (1H, d, *J*=2.1 Hz, H5), 6.41 (1H, d, *J*=2.1 Hz, H7), 3.89 (3H, s, C4 OCH₃), 3.76 (3H, s, C6 OCH₃), 3.11 (3H, s, CH₃). ¹³C NMR δ: 160.0 (C6), 155.0 (C4), 138.0 (C7a), 134.0 (ArC1'), 130.0 (ArC2' and 6'), 128.0 (ArC3' and 5'), 127.6 (ArC4'), 125.0 (C3), 122.0 (C2), 113.0 (C3a), 96.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.6 (C6 OCH₃), 40.0, CH₃. CIMS *m/z*: 332 ([M+1]⁺, 45%), 270 (15), 132 (15), 103 (20), 71 (100); ES-HRMS *m/z*: calcd for [M+1]⁺ C₁₇H₁₈NO₄S 332.0957; found; 332.0959.

4.2.9. 3-(4-Bromophenyl)-N-methanesulfonyl-4,6-dimethoxyindole (**23**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-bromophenyl)indole **6** (2.00 g, 6.0 mmol), sodium hydride (0.24 g, 6.0 mmol) and methanesulfonyl chloride (0.69 g, 6.0 mmol) to give the indole **25** (1.72 g, 70%) as a white solid, mp 147–150 °C. ¹H NMR δ: 7.51 (2H, d, *J*=9.0 Hz, ArH2' and 4'), 7.43 (2H, d, *J*=9.0 Hz, ArH3' and 5'), 7.22 (1H, s, H2), 7.02 (1H, d, *J*=2.1 Hz, H7), 6.40 (1H, d, *J*=2.1 Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.77 (3H, s, C6 OCH₃), 3.13 (s, Me). ¹³C NMR δ: 160.0 (C6), 156.5 (C4), 138.0 (C7a), 134.0 (ArC4'), 130.5 (ArC2' and 6'), 130.0 (ArC3' and 5'), 124.0

(ArC1'), 123.0 (C3), 122.0 (C2), 112.0 (C3a), 96.0 (C7), 90.0 (C5), 55.0 (C4 OCH₃), 54.5 (C6 OCH₃), 42.0 CH₃. CIMS *m/z*: 412 ([M+1]⁺, ⁸¹Br, 20%), 332 (15), 254 (10), 113 (25), 97 (30), 81 (100); ES-HRMS *m/z*: calcd for [M+1]⁺ C₁₇H₁₇NO₄S⁷⁹Br 410.0062; found 410.0061.

4.2.10. N-Methanesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)indole (**24**)

This was prepared by general procedure (A) using 4,6-dimethoxy-3-(4'-methoxyphenyl)indole **4** (2.00 g, 7.0 mmol), sodium hydride (0.28 g, 7.0 mmol) and methanesulfonyl chloride (0.81 g, 7.1 mmol) to give the indole **27** (1.99 g, 79%) as white solid, mp 145–148 °C. ¹H NMR δ: 7.50 (2H, d, *J*=9.0 Hz, ArH2' and 6'), 7.17 (1H, s, H2), 7.10 (1H, d, *J*=2.1 Hz, H5), 6.93 (2H, d, *J*=9.0 Hz, ArH3' and 5'), 6.40 (1H, d, *J*=2.1 Hz, H7), 3.88 (3H, s, C4 OCH₃), 3.85 (3H, s, C6 OCH₃), 3.77 (3H, s, ArC4 OCH₃), 3.09 (3H, s, CH₃). ¹³C NMR δ: 159.9 (ArC4'), 158.5 (C6), 155.0 (C4), 138.0 (C7a), 132.0 (ArC2' and 6'), 126.0 (ArC1'), 124.0 (C3), 121.0 (C2), 114.0 (ArC3' and 5'), 110.0 (C3a), 96.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.6 (C6 OCH₃), 55.7 (ArC4 OCH₃), 40.0 (CH₃). CIMS *m/z*: 362 ([M+1]⁺, 90%), 340 (15), 284 (30), 149 (15), 113 (20), 97 (25), 81 (100); EI-HRMS *m/z*: calcd for C₁₈H₁₉NO₅S 361.0983; found 361.0989.

4.2.11. N-Acetyl-4,6-dimethoxy-3-(4-methoxyphenyl)indole (**30**)

To a solution of 4,6-dimethoxy-3-(4-methoxyphenyl)indole **4** (0.50 g, 1.80 mmol), acetyl chloride (0.14 g, 1.80 mmol) and potassium hydroxide (0.10 g, 1.80 mmol) in CH₂Cl₂ (15 mL) was added a catalytic amount of the phase transfer catalyst tetrabutylammonium hexafluorophosphate (20 mol %) in CH₂Cl₂ (25 mL) under a nitrogen atmosphere. The reaction mixture was stirred for 12 h before water (30 mL) was added and the organic layer separated. The solvent was then removed under reduced pressure, and the oily liquid subjected to flash chromatography (ethylacetate/hexane 2:8) to yield the indole **31** (0.37 g, 65%) as a white solid, mp 145–158 °C. ¹H NMR δ: 7.77 (1H, d, *J*=1.8 Hz, H7), 7.51 (2H, d, *J*=9.0 Hz, Ar2' and 6'), 7.10 (1H, s, H2), 6.94 (2H, d, *J*=9.0 Hz, Ar3' and 5'), 6.40 (1H, d, *J*=1.8 Hz, H5), (3.89, 3.85 and 3.75, 3×OCH₃), 2.61 (3H, s, COCH₃). ¹³C NMR δ: 169.1 (C=O), 159.8 (ArC4'), 159.0 (C6), 154.4 (C4), 138.4 (C7a), 130.7 (ArC2' and 6'), 127.0 (ArC1'), 124.2 (C3), 120.4 (C2), 113.3 (ArC3' and 5'), 112.8 (C3a), 95.9 (C7), 93.1 (C5), (55.9, 55.5 and 55.4 3×OCH₃), 24.4 (COCH₃). CIMS *m/z*: 326 ([M+1]⁺, 100%), 283 (20), 268 (10). EI-HRMS *m/z*: calcd for C₁₉H₁₉NO₄ 325.1314; found 325.1398.

4.2.12. N-Acetyl-4,6-dimethoxy-3-(4'-nitrophenyl)indole (**31**)

To a solution of 4,6-dimethoxy-3-(4'-nitrophenyl)indole **7** (0.50 g, 1.70 mmol), acetyl chloride (0.13 g, 1.70 mmol) and potassium hydroxide (0.10 g, 1.8 mmol) in CH₂Cl₂ (15 mL) was added a catalytic amount of phase transfer catalyst tetrabutylammonium hexafluorophosphate (20 mol %) in CH₂Cl₂ (25 mL) under a nitrogen atmosphere. The reaction mixture was stirred for 12 h. To the reaction mixture water was added, and the organic layer separated. The solvent was removed under reduced pressure, and the oily liquid then purified by flash chromatography (ethylacetate/hexane, 2:8) to yield the indole **32** (0.32 g, 57%) as a white solid, mp 180–182 °C. ¹H NMR δ: 8.27 (2H, d, *J*=9.0 Hz, ArH2' and 6'), 7.87 (2H, d, *J*=9.0 Hz, ArH3' and 5'), (1H, s, H2), 7.70 (1H, d, *J*=1.8 Hz, H7), 6.53 (1H, d, *J*=1.8 Hz, H5), 3.84 (3H, s, C4 OCH₃), 3.79 (3H, s, C6 OCH₃), 2.68 (3H, s, acetyl CH₃). ¹³C NMR δ: 160.0 (C6), 154.0 (C4), 149.0 (C=O), 142.0 (ArC4'), 139.0 (C7a), 131.0 (ArC2' and 6'), 123.0 (ArC3' and 5'), 122.0 (C2), 121.8 (C3), 112.0 (C3a), 97.0 (C7), 93.0 (C5), 56.0 (C4 OCH₃), 55.0 (C6 OCH₃), 26.0 (COCH₃). CIMS *m/z*: 341 ([M+1]⁺, 100%); EI-HRMS *m/z*: calcd for C₁₈H₁₆N₂O₅ 340.1059; found 340.1072.

4.2.13. N-tert-Butylcarbamate-4,6-dimethoxy-3-(4-methoxyphenyl)indole (**34**)

To a solution of 4,6-dimethoxy-3-(4'-methoxyphenyl)indole **4** (0.50 g, 1.80 mmol) and di-tert-butylidicarbonate (0.37 g,

1.80 mmol) in CH_2Cl_2 (25 mL) was added a catalytic amount of DMAP and the reaction stirred under a nitrogen atmosphere for 30 min. To the reaction mixture was added water, the organic layer separated, the solvent removed under reduced pressure, and the solid then recrystallised from methanol to yield the indole **35** (0.68 g, 99%), as a white solid, mp 151–153 °C. ^1H NMR δ : 8.22 (1H, d, $J=1.8$ Hz, H7), 7.51 (2H, d, $J=9.0$ Hz, ArH2' and 6'), 6.92 (2H, d, $J=9.0$ Hz, ArH3' and 5'), 6.47 (1H, d, $J=1.8$ Hz, H5), 6.35 (d, $J=2.1$ Hz), 3.88 (3H, s, ArC4 OCH₃), 3.83 (3H, s, C4 OCH₃), 3.74 (3H, s, C6 OCH₃). ^{13}C NMR δ : 159.9 (C=O), 159.8 (C6), 154.3 (C4), 150.0 (C7a), 130.8 (ArC2' and 6'), 127.4 (ArC1'), 121.1 (C2), 113.2 (ArC3' and 5'), 106.8 (C3a), 95.0 (C7), 91.7 (C5), 84.0 (CO₂C(CH₃)), (55.8, 55.4 and 55.4, 3×OCH₃), 28.4 (3×C, *t*-BOC). CIMS m/z : 383 ($[\text{M}+1]^+$, 90%), 370 (15), 327 (100), 283 (30), 268 (10); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₂₂H₂₆NO₅ 384.1811; found 384.1807.

4.3. General procedure (B) for the synthesis of bisindoles: oxidative dimerisation

Solid thallium(III) trifluoroacetate (0.53 mmol) was mixed with the solid protected indole (1.06 mmol) under a nitrogen atmosphere. Dry acetonitrile (7 mL) was added and the resultant mixture cooled to 0 °C. Boron trifluoride diethyletherate (8.0 mmol) was then added slowly into the reaction mixture, and allowed to stir for 1 h. The reaction mixture was poured into water and extracted with CH_2Cl_2 (5×50 mL). The combined organic layers were concentrated to give a dark residue.

4.3.1. 2,2'-Bi-[N-benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)]indole (**12**)

This was prepared by general procedure (B) using *N*-benzenesulfonyl-3-(4-methoxyphenyl)-4,6-dimethoxyindole **8** (0.50 g, 1.18 mmol), thallium(III) trifluoroacetate (0.32 g, 0.55 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). The residue was recrystallised from CH_2Cl_2 /methanol to give the indole **12** (0.35 g, 70%) as a white solid, mp 242–246 °C. ^1H NMR δ : 7.53 (2H, d, $J=6.0$ Hz, sulfonyl ArH2'' and 6''), 7.46 (1H, t, sulfonyl ArH4''), 7.30 (2H, t, sulfonyl ArH3'' and 5''), 7.03 (2H, d, $J=12.0$ Hz, ArH2' and 6'), 7.02 (1H, d, $J=1.8$ Hz, H7), 6.65 (2H, d, $J=12.0$ Hz, ArH3' and 5'), 6.25 (1H, d, $J=1.8$ Hz, H5), 3.76 (3H, s, C4 OCH₃), 3.75 (3H, s, C6 OCH₃), 3.56 (1H, s, ArC4 OCH₃). ^{13}C NMR δ : 159.3 (ArC4'), 159.0 (C6), 154.5 (C4), 139.9 (C7a), 139.0 (sulfonyl ArC1''), 133.9 (ArC4), 131.8 (sulfonyl ArC2'' and 6''), 129.5 (sulfonyl ArC3'' and 5''), 127.0 (C2), 126.7 (ArC2' and 6'), 126.3 (C3), 126.5 (ArC1'), 124.0 (C3a), 122.0 (C2), 112.5 (C3a), 112.0 (ArC3' and 5'), 94.5 (C5), 92.0 (C7), 56.0 (C4 OCH₃), 55.0 (C6 OCH₃). ES-MS m/z : 845 ($[\text{M}+1]^+$, 35%), 797.2 (10), 704.2 (15), 625 (5), 552.3 (15), 475.4 (10), 411.2 (15), 332.3 (50), 304.3 (100), 272.3 (50), 244.2 (80), 214.0 (40), 157.9 (75), 141.8 (45), 105.9 (90); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₄₆H₄₁N₂O₁₀S₂ 845.2203; found 845.2209.

4.3.2. 2,2'-Bi-(N-benzenesulfonyl-4,6-dimethoxy-3-phenyl)-indole (**13**)

This was prepared by general procedure (B) using *N*-benzenesulfonyl-3-phenyl-4,6-dimethoxyindole **9** (0.50 g, 1.3 mmol), thallium(III) trifluoroacetate (0.35 g, 0.65 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH_2Cl_2 /methanol gave the indole **13** (0.34 g, 69% yield) as a white solid, mp 224–226 °C. ^1H NMR δ : 7.57–7.01 (10H, m, sulfonyl ArH2'', 3'', 4' and 6'', and ArH2', 3', 4', 5' and 6'), 7.00 (1H, d, $J=1.8$ Hz, H7), 6.22 (1H, d, $J=1.8$ Hz, H5), 3.76 (3H, s, C4 OCH₃), 3.53 (3H, s, C6 OCH₃). ^{13}C NMR δ : 159.3 (C6), 154.9 (C4), 139.0 (C7a), 139.5 (sulfonyl ArC1''), 133.8 (C3), 133.4 (sulfonyl ArC4''), 130.7 (sulfonyl ArC2'' and 6''), 129.1 (sulfonyl ArC3'' and 5''), 129.0 (ArC4'), 128.5 (ArC4'), 128.0 (C2), 127.5 (ArC2' and 6'), 127.2 (ArC3' and 5'), 126.9 (ArC1'), 122.3 (C3a), 95.5 (C7), 90.8 (C5), 55.9 (C4 OCH₃), 55.4 (C6 OCH₃). ES-MS

m/z : 785.0 ($[\text{M}+1]^+$, 30%), 746.2 (20), 649.4 (25), 552.2 (20), 413, 304, 272, 244 (100); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₄₄H₃₇N₂O₈S₂ 785.1991; found 785.1978.

4.3.3. 2,2'-Bi-[N-benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxy]indole (**14**)

This was prepared using general procedure (B) using thallium(III) trifluoroacetate (0.29 g, 0.53 mmol), *N*-benzenesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole **10** (0.50 g, 1.06 mmol) and boron trifluoride diethyletherate (1.12, 7.90 mmol). The residue was recrystallised from CH_2Cl_2 /methanol to give the indole **14** (0.37 g, 74%) as a white solid, mp 263–265 °C. ^1H NMR δ : 7.90 (2H, d, $J=9.0$ Hz, sulfonyl ArH2'' and 6''), 7.55–7.34 (7H, m, sulfonyl ArH3'' and 4'', 5'' and ArH2', 3', 5' and 6'), 7.19 (1H, d, $J=1.8$ Hz, H7), 6.34 (1H, d, $J=1.8$ Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.72 (3H, s, C6 OCH₃). ^{13}C NMR δ : 159.3 (C6), 154.5 (C4), 137.8 (C7a), 137.4 (ArC4'), 134.1 (sulfonyl ArC4'), 132.8 (C2), 131.2 (sulfonyl ArC2'' and 6''), 130.7 (sulfonyl ArC3'' and 5''), 129.9 (ArC1'), 129.4 (ArC1'), 129.2 (ArC2' and 6'), 126.8 (ArC3' and 5'), 123.5 (C3), 113.0 (C3a), 95.3 (C7), 89.9 (C5), 56.0 (C4 OCH₃), 55.3 (C6 OCH₃). ES-MS m/z : 942.7 ($[\text{M}+1]^+$, ⁸¹Br, 30%), 761.3 (10), 488.0 (20), 392.2 (30), 338.4 (30), 316.3 (35), 314.2 (70), 288.3 (60), 270.2 (50), 244.2 (45), 174.8 (30), 166.9 (40), 144.9 (70), 125.8 (80), 103.9 (100); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₄₄H₃₅N₂O₈⁷⁹Br₂S₂ 941.0202; found 941.0254.

4.3.4. 2,2'-Bi-[4,6-dimethoxy-3-(4-methoxyphenyl)]indole (**16**)

A solution of 2,2'-bi-[*N*-benzenesulfonyl-4,6-dimethoxy-3-(4-methoxyphenyl)]indole **12** (0.30 g, 0.3 mmol) in dioxane (20 mL) with 20% aqueous sodium hydroxide (5 mL) was heated at reflux for 4 h. The reaction mixture was poured onto cold water (60 mL) and partitioned with CH_2Cl_2 (3×50 mL). The solvent was evaporated under vacuum and the residue recrystallised from CH_2Cl_2 /methanol to give the indole **16** (0.03 g, 15%) as a pale yellow solid, mp 197–200 °C. ^1H NMR δ : 11.2 (1H, s, NH), 6.93 (2H, d, $J=9.0$ Hz, ArH2' and 6'), 6.59 (2H, d, $J=9.0$ Hz, ArH3' and 5'), 6.41 (1H, d, $J=1.8$ Hz, H5), 6.10 (1H, d, $J=1.8$ Hz, H7), 3.74 (3H, s, C4 OCH₃), 3.66 (3H, s, C6 OCH₃), 3.60 (3H, s, ArC4' OCH₃). ^{13}C NMR (DMSO *d*₆) δ : 158.3 (ArC4'), 158.2 (C6'), 154.3 (C4'), 138.0 (C7a), 132.0 (ArC2' and 6'), 128.5 (ArC1'), 124.6 (C2), 116.3 (C3), 112.2 (ArC3' and 5'), 112.0 (C3a), 92.0 (C5), 87.2 (C7), 55.9 (C4 OCH₃), 55.6 (C6 OCH₃), 55.4 (ArC4' OCH₃); CIMS m/z : 564 ($[\text{M}+1]^+$, 60%), 338 (100), 285 (10), 197 (5), 153 (10), 137 (25), 121 (30), 97 (30); ES-HRMS m/z : calcd $[\text{M}+1]^+$ C₃₄H₃₃N₂O₆ 564.2260; found 564.2259.

4.3.5. 2,2'-Bi-(4,6-dimethoxy-3-phenyl)indole (**17**)

A solution of 2,2'-bi-(*N*-benzenesulfonyl-4,6-dimethoxy-3-phenyl)indole **13** (0.30 g, 0.4 mmol) in dioxane (20 mL) with 20% aqueous sodium hydroxide (5 mL) was heated at reflux for 4 h. The reaction mixture was poured into cold water and partitioned with CH_2Cl_2 (3×50 mL). The solvent was evaporated and the residue recrystallised from CH_2Cl_2 /methanol yielding the indole **17** (0.03 g, 15%) as a pale yellow solid, mp 174–178 °C. ^1H NMR δ : 10.00 (1H, s, NH), 7.21–7.18 (2H, m, ArH2' and 6'), 7.09–7.06 (3H, m, ArH3', 4' and 5'), 6.49 (1H, d, $J=3.0$ Hz, H7), 6.18 (1H, d, $J=3.0$ Hz, H5), 3.77 (3H, s, C4 OCH₃), 3.66 (3H, s, C6 OCH₃). ^{13}C NMR δ : 158.0 (C6), 155.0 (C4), 138.0 (C7a), 136.0 (C3), 131.0 (ArC4'), 126.7 (ArC2' and 6'), 125.2 (ArC3' and 5'), 125.0 (ArC1'), 118.0 (C2), 112.0 (C3a), 92.0 (C7), 86.2 (C5), 55.0 (C4 OCH₃), 54.5 (C6 OCH₃). ES-MS m/z : 504.2 ($[\text{M}+1]^+$, 60%), 338.4 (25), 144.9 (20), 105.9 (45), 103.9 (100), 71.0 (25); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₃₂H₂₉N₂O₄ 504.2049; found 504.2054.

4.3.6. 2,2'-Bi-[3-(4-bromophenyl)-4,6-dimethoxy]indole (**18**)

This was prepared by general procedure (B) using 4,6-dimethoxy-3-(4-bromophenyl)indole **6** (0.20 g, 0.6 mmol),

thallium(III) trifluoroacetate (0.16 g, 0.3 mmol) and boron trifluoride diethyletherate (0.43 g, 3.0 mmol). The residue was subjected to flash silica gel column chromatography (50:50 hexane/ethylacetate) yielding the indole **18** (0.04 g, 20%) as a pale yellow solid, mp 195–197 °C. ^1H NMR δ : 7.85 (2H, d, $J=7.5$ Hz, ArH3' and 5'), 7.21 (2H, d, $J=7.5$ Hz, ArH2' and 6'), 7.28 (1H, d, $J=1.8$ Hz, H7), 6.32 (1H, s, H5), 3.84 (3H, s, C4 OCH₃), 3.49 (3H, s, C6 OCH₃). ^{13}C NMR (CDCl₃) δ : 153.5 (C-6 and C-4), 140.0 (C-7a), 133.0 (C-2), 132.8 (C-3), 130.4 (ArC-4'), 128.0 (ArC2' and 6'), 127.8 (ArC2' and 5'), 126.0 (C-3a), 121.6 (ArC1'), 94.3 (C-7), 92.0 (C-5), 57.8 (C-4-OCH₃), 56.0 (C-6-OCH₃). ES-MS m/z : 663 ([M+1]⁺, ^{80}Br 60%), 607 (10), 468.5 (15), 332 (40), 316 (50), 304 (50), 288 (70), 244 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₃₂H₂₇N₂O₄⁹⁹Br₂ 661.0338; found 661.0331.

4.3.7. 2,2'-Bi-[3-(4-bromophenyl)-(N-p-methylbenzenesulfonyl)-4,6-dimethoxy]indole (**25**)

This was prepared by general procedure (B) using *N*-(4-methylphenylsulfonyl)-3-(4-bromophenyl)-4,6-dimethoxyindole **19** (0.50 g, 1.0 mmol), thallium(III) trifluoroacetate (0.28 g 0.50 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **25** (0.36 g 73%) as a yellow solid, mp 219 °C. ^1H NMR δ : 7.45 (2H, d, $J=8.4$ Hz, sulfonyl ArH3' and 5'), 7.17 (2H, d, $J=8.7$ Hz, ArH2' and 6'), 7.14 (2H, d, $J=8.7$ Hz, ArH3' and 5'), 7.04 (1H, d, $J=2.1$ Hz, H7), 6.86 (2H, d, $J=8.4$ Hz, sulfonyl ArH2' and 6'), 6.26 (1H, d, $J=2.1$ Hz, H5), 3.79 (3H, s, C6 OCH₃), 3.57 (3H, s, C4 OCH₃), 2.36 (3H, s, CH₃). ^{13}C NMR δ : 159.6 (C6), 154.8 (C4), 144.7 (sulfonyl ArC4'), 138.7 (C7a), 136.7 (ArC4'), 132.9 (ArC1'), 132.2 (sulfonyl ArC2' and 6'), 130.1 (ArC2' and 6'), 129.8 (ArC3' and 5'), 127.2 (sulfonyl ArC3' and 5'), 126.0 (sulfonyl ArC1'), 124.0 (C2), 121.4 (C3), 112.3 (C3a), 95.4 (C7), 90.8 (C5), 55.8 (C4 OCH₃), 55.3 (C6 OCH₃), 21.8 (Me). ES-MS m/z : 970.7 ([M+1]⁺, ^{81}Br , 20%), 501.7 (20), 304.3 (30), 288.2 (35), 272.2 (40), 244.3 (100), 225.1 (45), 103.7 (80). ES-HRMS m/z : calcd for [M+1]⁺ C₄₆H₃₇N₂O₂S₂⁹⁹Br₂ 969.0515; found 969.0524.

4.3.8. 2,2'-Bi-[3-(4-bromophenyl)-4,6-dimethoxy-(N-p-nitrobenzenesulfonyl)]indole (**26**)

This was prepared by general procedure (B) using *N*-(p-nitrobenzenesulfonyl)-3-(4-bromophenyl)-4,6-dimethoxyindole **20** (0.50 g, 1.1 mmol), thallium(III) trifluoroacetate (0.30 g, 0.55 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **26** (0.33 g, 67%) as a yellow solid, mp 232–235 °C. ^1H NMR δ : 8.14 (2H, d, $J=8.7$ Hz, ArH3 and 5), 7.48 (2H, d, $J=9.0$ Hz, ArH2 and 6), 7.20 (2H, d, $J=8.7$ Hz, sulfonyl ArH2' and 6'), 7.02 (1H, d, $J=1.8$ Hz, H7), 7.03 (2H, d, $J=9.0$ Hz, sulfonyl ArH3' and 5'), 6.39 (1H, d, $J=1.8$ Hz, H4), 3.82 (3H, s, C4 OCH₃), 3.63 (s, C6 OCH₃). ^{13}C NMR δ : 160.6 (C6), 155.2 (C4), 150.3 (sulfonyl ArC4), 144.4 (sulfonyl ArC1), 139.1 (C7a), 132.3 (ArC1), 132.2 (sulfonyl ArC3 and 5), 130.4 (ArC2 and 6), 130.0 (C2), 127.9 (sulfonyl ArC2 and 6), 124.51 (ArC3 and 50), 122.9 (ArC4), 122.0 (C30), 112.4 (C3a), 96.0 (C7), 90.8 (C5), 56.0 (C4 OCH₃), 55.4 (C6 OCH₃ and ArC4 OCH₃). ES-MS m/z : 1031 ([M+1]⁺, ^{81}Br , 100%), 887.3 (10), 757.9 (30), 705.9 (60), 662.0 (50), 629.8 (50), 585.9 (80), 564.0 (95), 498.8 (70), 451.6 (45), 391.6 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₄₄H₃₃N₄O₁₂S₂⁹⁹Br₂ 1030.9903; found 1030.9919.

4.3.9. 2,2'-Bi-(N-methanesulfonyl-4,6-dimethoxy-3-phenyl)-indole (**27**)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-phenyl-4,6-dimethoxyindole **22** (0.50 g, 1.5 mmol), thallium(III) trifluoroacetate (0.41 g, 0.75 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **27** (0.33 g, 67%) as a white solid, mp 229–231 °C. ^1H NMR δ : 7.25–7.16 (4H, m, ArH2', 3', 4', 5' and 6'), 7.11 (1H, d, $J=3.0$ Hz, H5), 6.32 (1H, d, $J=3.0$ Hz, H7), 3.87 (3H, s, C4 OCH₃), 3.59 (3H, s, C6 OCH₃), 2.90 (3H, s, CH₃). ^{13}C NMR δ : 160.0

(C7), 155.0 (C4), 148.0 (C7a), 134.0 (ArC1'), 130.0 (ArC2' and 6'), 127.3 (C3), 127.1 (ArC3' and 5'), 127.0 (C2), 126.5 (ArC4'), 113.0 (C3a), 95.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.6 (C6 OCH₃), 40.0 (Me). ES-MS m/z : 661.1 ([M+1]⁺, 50%), 348.1 (20), 132.8 (18), 105.9 (30), 103.9 (40), 71.0 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₃₄H₃₃N₂O₈S₂ 661.1678; found 661.1674.

4.3.10. 2,2'-Bi-[3-(4-bromophenyl)-N-methanesulfonyl-4,6-dimethoxy]indole (**28**)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-(4-bromophenyl)-4,6-dimethoxyindole **23** (0.50 g, 1.2 mmol), thallium(III) trifluoroacetate (0.33 g, 0.6 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **28** (0.40 g, 80%) as a white solid, mp 179–182 °C. ^1H NMR δ : 7.31 (2H, d, $J=9.0$ Hz, ArH2 and 6), 7.08 (1H, d, $J=3.0$ Hz, H5), 7.02 (2H, d, $J=9.0$ Hz, ArH3 and 5), 6.33 (1H, d, $J=3.0$ Hz, H7), 3.89 (3H, s, C4 OCH₃), 3.61 (3H, s, C6 OCH₃), 3.06 (3H, s, CH₃). ^{13}C NMR δ : 160.2 (C6), 155.0 (C4), 138.0 (C7a), 132.7 (ArC4'), 132.0 (ArC3' and 5'), 130.3 (ArC2' and 6'), 126.3 (ArC1'), 123.3 (C2), 121.4 (C3), 112.6 (C3a), 95.6 (C7), 89.9 (C5), 56.8 (C4 OCH₃), 56.0 (C6 OCH₃), 40.0 (CH₃). ES-MS m/z : 817.9 ([M+1]⁺, ^{81}Br , 15%), 466.8 (25), 464.7 (10), 343.8 (10), 341.4 (15), 338.4 (30), 275.9 (15), 204.9 (20), 144.9 (20), 132.8 (25), 105.9 (40), 103.9 (100), 71.0 (35), 60.1 (50); ES-HRMS m/z : calcd for [M+1]⁺ C₃₄H₃₁N₂O₈S₂⁷⁹Br₂ 816.9889; found 816.9863.

4.3.11. 2,2'-Bi-[N-methanesulfonyl-4,6-dimethoxy-(4-methoxyphenyl)]indole (**29**)

This was prepared by general procedure (B) using *N*-methanesulfonyl-3-(4-methoxyphenyl)-4,6-dimethoxyindole **24** (0.50 g, 1.3 mmol), thallium(III) trifluoroacetate (0.36 g 0.65 mmol) and boron trifluoride diethyletherate (1.12 g, 7.9 mmol). Recrystallisation from CH₂Cl₂/methanol gave the indole **29** (0.38 g, 76%) as a white solid, mp 219–222 °C. ^1H NMR δ : 7.18 (2H, d, $J=9.0$ Hz, ArH2' and 6'), 7.12 (1H, d, $J=1.8$ Hz, H7), 6.74 (2H, d, $J=9.0$ Hz, ArH3' and 5'), 6.33 (1H, d, $J=1.8$ Hz, H5), 3.88 (3H, s, C4 OCH₃), 3.74 (3H, s, C6 OCH₃), 3.62 (3H, s, ArC4 OCH₃), 2.93 (3H, s, CH₃). ^{13}C NMR δ : 159.9 (ArC4'), 158.5 (C6), 155.0 (C4), 138.0 (C7a), 132.0 (ArC2' and 6'), 126.0 (ArC1'), 123.8 (C3), 123.4 (C2), 113.5 (C3a), 113.0 (Ar3' and 5'), 96.0 (C7), 90.0 (C5), 56.0 (C4 OCH₃), 55.7 (C6 OCH₃), 55.6 (ArC4 OCH₃), 40.0 (CH₃); ES-MS m/z : 721.2 ([M+1]⁺, 40%), 547.4 (15), 503.9 (10), 376.1 (20), 360.4 (25), 338.4 (70), 304.2, 244.1, 103.9 (50), 71.0 (50), 60.1 (100); ES-HRMS m/z : calcd for [M+1]⁺ C₃₆H₃₇N₂O₁₀S₂ 721.1889; found 721.1883.

4.3.12. 2,2'-Bi[N-acetyl-4,6-dimethoxy-3-(4-nitrophenyl)]indole (**35**)

This was prepared by general procedure (B) using *N*-acetyl-4,6-dimethoxy-3-(4-nitrophenyl)indole **32** (0.20 g, 0.6 mmol), thallium(III) trifluoroacetate (0.16 g, 0.3 mmol) and boron trifluoride diethyletherate (1.02 mL, 7.2 mmol). The residue was subjected to flash silica gel column chromatography (70:30 hexane/ethylacetate) yielding the indole **36** (0.06 g, 30%) as a pale yellow solid, mp 179–181 °C. ^1H NMR δ : 8.20 (2H, d, $J=9.0$ Hz, ArH3' and 5'), 7.61 (1H, d, $J=2.1$ Hz, H7), 7.57 (1H, d, $J=9.0$ Hz, ArH2' and 6'), 6.36 (1H, d, $J=2.1$ Hz, H5), 3.89 (3H, s, C4 OCH₃), 3.70 (3H, s, C6 OCH₃), 2.59 (3H, s, acetyl). ^{13}C NMR δ : 171.0 (C=O), 163.9 (C6), 157.8 (C4), 147.5 (ArC1'), 146.1 (ArC4'), 142.3 (C7a), 126.3 (ArC2' and 6'), 124.0 (ArC3' and C5'), 122.5 (C2), 104.0 (C3a), 96.5 (C7), 96.0 (C5), 56.0 (C6 OCH₃), 55.7 (C4 OCH₃), 26.7 (COCH₃). EIMS m/z : 689 ([M+1]⁺, 7%), 663 (100).

4.3.13. 2-(4,6-Dimethoxy-3-(4-nitrophenyl)-N-trifluoroacetyl-1H-indol-7-yl)-4,4',6,6'-tetramethoxy-3,3'-bi(4-nitrophenyl)-N,N'-di(trifluoroacetyl)-1H,1'H-7,7'-biindole (**36**)

This was prepared by general procedure (B) using thallium(III) trifluoroacetate (0.14 g, 0.25 mmol), *N*-trifluoro-4,6-dimethoxy-

3-(4'-nitrophenyl)-1-trifluoroacetateindole **34** (0.20 g, 0.50 mmol) and boron trifluoride diethyletherate (0.29 g, 2.0 mmol). The residue was subjected to flash silica gel column chromatography (60:40 hexane/ethylacetate) yielding the trimer **37** (0.70 g, 35%) as a yellow solid, mp 165–168 °C. ^1H NMR δ : 8.24 (2H, d, $J=5.1$ Hz, ArH2' and 6'), 8.21 (2H, d, $J=5.1$ Hz, ArH2'' and 6''), 7.94 (2H, d, $J=5.1$ Hz, ArH2 and 6), 7.66 (2H, d, $J=4.8$ Hz, ArH3' and 5'), 7.55 (2H, d, $J=4.8$ Hz, ArH3 and 5), 6.95 (1H, s, H2'), 6.73 (1H, s, H5'), 6.66 (1H, s, H5), 6.44 (1H, s, H5'), 3.95 (C4 OCH₃''), 3.92 (C4 OCH₃), 3.84 (C6 OCH₃''), 3.81 (C4 OCH₃'), 3.79 (C6 OCH₃), 3.72 (C6 OCH₃'), ^{13}C NMR δ : 160.2 (C6'), 157.4 (C6''), 156.8 (C6), 156.6 (C4'), 154.8 (C4''), 154.0 (C4), 147.4 (ArC4'), 147.3 (ArC4''), 146.4 (ArC4), 142.0 (ArC1'), 141.1 (ArC1''), 140.0 (ArC1), 137.2 (C7a''), 137.2 (C7a), 137.0 (C7a'), 130.4 (ArC2' and 6', ArC2' and 6'', and ArC2 and 6), 128.0 (C2), 124.6 (C3''), 124.3 (C3'), 123.0 (ArC3' and 5' and ArC3'' and 5''), 122.4 (ArC3 and 5), 121.9 (C2''), 121.3 (C2'), 121.0 (C3), 114.2 (C3a''), 113.1 (C3a'), 112.8 (C3a), 106.2 (C7''), 105.5 (C7), 100.2 (C7'), 94.5 (C5''), 92.6 (C5), 91.8 (C5'), 57.2 (C4 OCH₃''), 57.1 (C4 OCH₃), 56.0 (C6 OCH₃'), 55.6 (C4 OCH₃'), 55.5 (C6 OCH₃''), 55.3 (C6 OCH₃). ES-MS m/z : 1179.2 ($[\text{M}+1]^+$, 25%), 1082.3 (20), 986.3 (30), 416.2 (20), 400.3 (20), 338.4 (30), 313.2 (10), 218.9 (30), 179.0 (50), 104.0 (100); ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₅₄H₃₆N₆O₁₅F₉ 1179.2095; found 1179.2089.

4.3.14. 4,6-Dimethoxy-2-[4',6'-dimethoxy-3'-(4-nitrophenyl)indol]-7'-yl]-3-(4-nitrophenyl)indole (**37**)

This was prepared by general procedure (B) using 4,6-dimethoxy-3-(4'-nitrophenyl)indole **6** (0.20 g, 0.67 mmol), thallium(III) trifluoroacetate (0.18 g, 0.34 mmol) and boron trifluoride diethyletherate (0.47 g, 3.3 mmol). The crude residue was subjected to flash silica gel column chromatography (70:30, hexane/ethylacetate) to give the indole **38** (0.04 g, 20%) as a yellow solid, mp 245–248 °C. ^1H NMR δ : 8.56 (1H, s, NH), 8.18 (2H, d, $J=5.1$ Hz, ArH3'' and 5''), 8.02 (2H, d, $J=5.4$ Hz, ArH3' and 5'), 7.94 (1H, s, NH'), 7.67 (2H, d, $J=5.1$ Hz, ArH2'' and 6''), 7.55 (2H, d, $J=5.4$ Hz, ArH2' and 6'), 6.88 (1H, s, H2'), 6.57 (1H, d, $J=1.8$ Hz, H7), 6.41 (1H, s, H5'), 6.32 (1H, d, $J=1.8$ Hz, H5), 3.91 (3H, s, C6 OCH₃), 3.88 (3H, s, C6' OCH₃), 3.79 (3H, s, C4 OCH₃), 3.75 (3H, s, C4' OCH₃), ^{13}C NMR δ : 158.3 (C6), 155.6 (C6'), 154.5 (C4), 155.3 (C4'), 145.7 (ArC1''), 145.4 (ArC1'), 144.9 (ArC4'), 143.7 (ArC4''), 138.2 (C7a), 137.7 (C7a'), 130.9 (ArC2' and 6'), 129.8 (ArC2'' and 6''), 127.5 (C3), 122.8 (ArC3' and 5'), 123.3 (ArC3'' and 5''), 122.9 (C2'), 117.3 (C3'), 114.9 (C2), 111.2 (C3a), 110.1 (C3a'), 97.1 (C7), 92.7 (C5), 89.7 (C5'), 86.9 (C7), 55.0 (C4 OCH₃), 53.5 (C4' OCH₃), 53.1 (C6' OCH₃), 52.0 (C6). ES-MS m/z : 594.2 ($[\text{M}+1]^+$, 15%), 338.5 (50), 336.2 (100), 147.1 (90). ES-HRMS m/z : calcd for $[\text{M}+1]^+$ C₃₂H₂₆N₄O₈ 595.1829; found 595.1832.

4.4. Structure determinations

Full spheres of CCD diffractometer data were measured (monochromatic Mo K α radiation, $\lambda=0.71073$ Å; ω -scans; T ca. 153 K) yielding N_{total} reflections, these merging to N unique (R_{int} cited) after 'empirical'/multiscan absorption correction (proprietary software), N_0 with $F>4\sigma(F)$ being considered 'observed'. All were used in the full matrix least squares refinement on F^2 , refining

anisotropic displacement parameters for the non-hydrogen atoms, hydrogen atom treatment following a riding model; reflection weights were $(\sigma^2(F^2) + (aP)^2)^{-1}$ ($P = (F_0^2 + 2F_c^2)/3$). Neutral atom complex scattering factors were used within the SHELXL 97 program.¹⁴ Pertinent results are given in Figures 1 and 2 and Ref. 15; full .cif depositions (without structure factor amplitudes) are deposited with the Cambridge Crystallographic Data Centre, CCDC 680107 and 680108.

Acknowledgements

NRY would like to thank the University of Wollongong for a scholarship.

References and notes

- For general reviews of biaryl synthesis, see: (a) Bringmann, G.; Walter, R.; Weirich, R. *Angew. Chem.* **1990**, 29, 977–991; (b) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, 102, 1359–1469.
- For reviews on the atropselective synthesis of biaryls, see: (a) Bringmann, G.; Price Mortimer, A. J.; Keller, P. A.; Garner, J.; Gresser, M. J.; Breuning, M. *Angew. Chem.* **2005**, 44, 5384–5427; (b) Wallace, T. *Org. Biomol. Chem.* **2006**, 4, 3197–3210; (c) Lloyd-Williams, P.; Giralt, E. *Chem. Soc. Rev.* **2001**, 30, 145–157.
- (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, 95, 2457–2483; (b) Suzuki, A. *J. Organomet. Chem.* **1999**, 576, 147–168; (c) Kotha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, 58, 9633–9695; (d) Daudin, O. *Eur. J. Org. Chem.* **2005**, 4223–4229.
- See Ref. 1 and (a) Gu, R.; Van Hecke, K.; Van Meervelt, L.; Toppet, S.; Dehaen, W. *Org. Biomol. Chem.* **2006**, 4, 3785–3789; (b) Pezzella, A.; Panzella, L.; Natangelo, A.; Arzillo, M.; Napolitano, A.; d'Ischia, M. *J. Org. Chem.* **2007**, 72, 9225–9230 and references therein.
- (a) Black, D. S.; Keller, P. A.; Kumar, N. *Tetrahedron Lett.* **1989**, 30, 5807–5808; (b) Keller, P. A.; Birch, C.; Leach, S. P.; Tyssen, D.; Griffith, R. J. *Mol. Graphics* **2003**, 21, 365–373; (c) Bremner, J. B.; Coates, J. A.; Coghlan, D. R.; David, D. M.; Keller, P. A.; Pyne, S. G. *New J. Chem.* **2002**, 26, 1549–1552.
- See Ref. 4a and (a) Abbiati, G.; Arcadi, A.; Beccalli, E.; Bianchi, G.; Marinelli, F.; Rossi, E. *Tetrahedron* **2006**, 62, 3033–3039; (b) Wada, Y.; Nagasaki, H.; Tokuda, M.; Orito, K. *J. Org. Chem.* **2007**, 72, 2008–2014; (c) Merlic, C. A.; You, Y.; McInnes, D. M.; Zechman, A. L.; Miller, M. M.; Deng, Q. *Tetrahedron* **2001**, 5199–5212 and references therein; (d) Koza, D. J.; Euler, W. B. *Heterocycl. Commun.* **1999**, 5, 399–402.
- (a) Black, D. StC.; Bowyer, M. C.; Catalano, M. M.; Ivory, A. J.; Keller, P. A.; Kumar, N.; Nugent, S. J. *Tetrahedron* **1994**, 50, 10497–10508; (b) Keawin, T.; Rajviroongit, S.; Black, D. StC. *Tetrahedron* **2005**, 61, 853–861.
- McKillop, A.; Turrell, A. G.; Young, D. W.; Taylor, E. C. *J. Am. Chem. Soc.* **1980**, 102, 6504–6512.
- Banerji, A.; Ray, R.; Pal, S. C.; Banerji, D.; Maiti, K. K. *J. Indian Chem. Soc.* **1998**, 75, 698–704.
- Tholander, J.; Bergman, J. *Tetrahedron* **1999**, 55, 12595–12602.
- Black, D. S. C.; Bowyer, M. C.; Bowyer, P. K.; Ivory, A. J.; Kim, M.; Kumar, N.; McConnell, D. B.; Popielek, M. *Aust. J. Chem.* **1994**, 47, 1741–1750.
- Black, D. StC.; Ivory, A. J.; Kumar, N. *Tetrahedron* **1996**, 52, 7003–7012.
- (a) D'Ischia, M.; Napolitano, A.; Tsiakas, K.; Prota, G. *Tetrahedron* **1990**, 46, 5789–5796; (b) Napolitano, A.; D'Ischia, M.; Prota, G.; Schultz, T. M.; Wolfram, L. J. *Tetrahedron* **1989**, 45, 6749–6760.
- Sheldrick, G. M. *SHELXL 97—A Program for Crystal Structure Refinement*; University of Göttingen: Göttingen, 1997.
- 12.** C₄₆H₄₀N₂O₁₀S₂, $M=845.0$. Monoclinic, space group $P2_1/n$, $a=12.725(1)$, $b=17.671(1)$, $c=18.158(1)$ Å, $\beta=91.449(2)^\circ$, $V=4082$ Å³, D_c ($Z=4$)=1.375 g cm⁻³, $\mu_{\text{Mo}}=0.19$ mm⁻¹; specimen: $0.10\times0.09\times0.08$ mm; $T_{\text{min/max}}=0.82$, $2\theta_{\text{max}}=70^\circ$; $N_{\text{t}}=67457$, $N=18054$ ($R_{\text{int}}=0.077$), $N_0=8927$; $R1=0.055$, $wR2=0.14$ ($a=0.067$). $|\Delta\rho_{\text{max}}|=1.38$ e Å⁻³. **28.** C₃₄H₃₀Br₂N₂O₈S₂·2(CH₃)₂CO, $M=934.7$. Monoclinic, space group $C2/c$, $a=20.766(2)$, $b=9.7298(8)$, $c=22.543(2)$ Å, $\beta=117.101(2)^\circ$, $V=4055$ Å³, D_c ($Z=4$)=1.531 g cm⁻³, $\mu_{\text{Mo}}=2.2$ mm⁻¹; specimen: $0.28\times0.15\times0.12$ mm; $T_{\text{min/max}}=0.72$, $2\theta_{\text{max}}=75^\circ$; $N_{\text{t}}=42853$, $N=10294$ ($R_{\text{int}}=0.045$), $N_0=5440$; $R1=0.037$, $wR2=0.077$ ($a=0.034$). $|\Delta\rho_{\text{max}}|=0.95$ e Å⁻³.