



Cyclodehydrogenation of hetero-oligophenylenes

Samuthirapandian Nagarajan[†], Cécile Barthes, André Gourdon^{*}

NanoSciences Group, CNRS, Université de Toulouse, CEMES (Centre d'Elaboration des Matériaux et d'Etudes Structurales), BP 94347, 29 rue J. Marvig, F-31055 Toulouse, France

ARTICLE INFO

Article history:

Received 8 January 2009
Received in revised form 11 February 2009
Accepted 13 February 2009
Available online 21 February 2009

Keywords:

Arenes
Nitrogen heterocycles
Cycloaddition
Cross-coupling
Scholl reaction

ABSTRACT

Pyrimidyl-penta-phenylbenzenes have been synthesized by Diels–Alder addition of phenyl-ethynylpyrimidines and tetraphenylcyclopentadienones under microwave irradiation. Scholl reactions of these compounds led to two types of hetero polyaromatic hydrocarbons: (a) partial cyclization by creation of two C–C bonds *ortho* to the pyrimidine nitrogen atoms gave substituted tribenzo[e,gh,j]perimidine (N-1/3HSB) in high yields; (b) when the position 2 of the pyrimidine ring was substituted by a *tert*-butyl group, the Scholl reaction was complete and provided the first example of a diaza-hexa-peribenzocoronene.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Polyaryl and polycyclic hydrocarbons are currently attracting a lot of interest for their potential applications as molecular materials for electronic and optoelectronic applications. In the course of a study of the mechanical and chemical properties of molecular devices,¹ we were interested in planar heterosuperbenzene as candidates for molecular moulding,² metal atom manipulation,³ electrode–molecule contact⁴ or controlled rotation⁵ experiments at the single molecule level. In the present contribution, the investigation of the synthesis of some pyrimidyl-penta-phenylbenzenes and the influence of the substituents on the cyclodehydrogenation reaction to access heterosuperbenzene is explored.

2. Results and discussion

2.1. Synthesis of the pyrimidyl-penta-phenylbenzenes

The general synthetic route towards (partially or fully) cyclodehydrogenated hetero-oligophenylenes is shown on Scheme 1. The propeller-like pyrimidyl-penta-phenylbenzenes were obtained by [2+4] Diels–Alder cycloaddition of tetraphenylcyclopentadienones to phenylethynylpyrimidines.

These latter were prepared by Sonogashira coupling of bromopyrimidines with ethynylbenzenes. All these syntheses were done

by microwave (μ W) heating, which revealed as a very efficient tool for cross-couplings and Diels–Alder reactions in terms of reaction times and yields.⁶ Typically, the cross-coupling reactions were completed in less than 0.5 h in good yields (61–93%).^{7–9} As pointed out by Erdélyi and Gogoll,¹⁰ the reaction temperature was maintained just below 120 °C to avoid decomposition of the palladium catalyst. The Diels–Alder reactions, and the subsequent in-situ oxidations, were done in diphenylether at 260 °C for 45 min giving the hexaaryl compounds in 40–50% yields.

The unsubstituted (5-pyrimidyl)pentaphenylbenzene **7** was obtained in 44% yield by reaction of commercial 2,3,4,5-tetraphenylcyclopenta-2,4-dienone with 5-(phenylethynyl)pyrimidine **5** (Scheme 2).


This latter compound **5** was prepared by coupling phenylacetylene and 5-bromopyrimidine in good yield (93%), significantly higher than by conventional heating.^{11–13} Similarly, the tri-substituted (5-pyrimidyl)pentaphenylbenzene **11** was obtained by [2+4] cycloaddition of 5-((4-*tert*-butylphenyl)ethynyl)pyrimidine **9** and 3,4-bis(4-*tert*-butylphenyl)-2,5-diphenylcyclopenta-2,4-dienone **10**^{14–16} (Scheme 3).

The tolane **9** was synthesized from 1-*tert*-butyl-4-ethynylbenzene **8**. This latter compound^{17,18} was prepared in 88% yield by cross-coupling under microwave irradiation of 1-bromo-4-*tert*-butylbenzene and trimethylsilylacetylene, and deprotection by aqueous NaOH in methanol.

The penta-*tert*-butyl propeller **13** (Scheme 4) was correspondingly synthesized by cycloaddition of **9** with the known cyclopentadienone **12**,^{14,15} obtained by double Knoevenagel condensation of 4,4'-di-*tert*-butylbenzil and 1,3-bis(3,5-di-*tert*-butylphenyl)-2-propanone.

^{*} Corresponding author. Tel.: +33 (0)5 62 25 78 59; fax: +33 (0)5 62 25 79 99.
E-mail address: gourdon@cemes.fr (A. Gourdon).

[†] Present address: Department of Chemistry, Annamalai University, Annamalai-nagar 608002, India.



Scheme 1. General route towards pyrimidyl-penta-phenylbenzenes and their cyclodehydrogenation products. (a) Sonogashira coupling, μw ; (b) [2+4] cycloaddition, Ph_2O , μw ; (c) cyclodehydrogenation, FeCl_3 , DCM, CH_3NO_2 .

Finally, the fully substituted propeller **16** was obtained by condensation of **15** and **12**.

The preparation of **15** requires the synthesis of 5-bromo-2-*tert*-butylpyrimidine **14**, which was made by condensation of malondialdehyde bis(dimethylacetal) and 2,2-dimethylpropanamidine¹⁹ to give the 2-*tert*-butylpyrimidine, subsequently brominated in position 5 by bromine in acetic acid²⁰ (Scheme 5).

2.2. Cyclodehydrogenations


With the objective to prepare series of heterosuperbenzene, we have investigated the intramolecular Scholl reaction on these contiguous hexaarylbenzenes **7**, **11**, **13** and **16**. The Scholl reaction, i.e., the condensation of aromatics by reaction with Lewis acids, is known to be very efficient for the cyclodehydrogenation of oligophenylene compounds. Müllen and co-workers have extensively used this reaction to convert 3D-polyphenylene structures to all benzenoid 2D-hydrocarbons,²¹ and in particular for the preparation of hexabenzocoronenes from hexaphenylbenzenes.²² In this Scholl reaction, the oxidative aryl–aryl coupling is effected in general by transition metal halides such as $\text{AlCl}_3/\text{Cu(II)}$, FeCl_3 , MoCl_5 but also by $\text{PhI}(\text{O}_2\text{CCF}_3)_2/\text{BF}_3 \cdot \text{OEt}_2$,^{23,24} which act in the same time as oxidants and Lewis acids. This synthetic route has been recently extended by Draper and co-workers to the cyclodehydro-

genation of 1,2-dipyrimidyl-3,4,5,6-tetra(4-*tert*-butylphenyl)benzene (Scheme 6).^{25,26}


Remarkably, whereas Scholl cyclodehydrogenation by $\text{AlCl}_3/\text{CuCl}_2$ in CS_2 gave solely the fully cyclized heterosuperbenzene N-HSB, the reaction employing a milder catalyst (FeCl_3 /nitromethane/DCM) gave a mixture of N-HSB (35%) and of ‘half-cyclized’ N-1/2HSB (32%). Surprisingly, treatment of N-1/2HSB by FeCl_3 , or under forcing conditions by $\text{AlCl}_3/\text{CuCl}_2$, did not give N-HSB, which indicates that the partially cyclodehydrogenated compound is not a parent intermediate of the heterosuperbenzene in this reaction. It is worth noting that some partially cyclodehydrogenated oligophenylenes can be fully cyclodehydrogenated to give HBC in quantitative yield and without chlorination.²⁷

2.3. Scholl reaction on the hexaarylbenzenes **7**, **11**, **13** and **16**

Attempts to cyclodehydrogenate these four compounds by $\text{AlCl}_3/\text{CuCl}_2$ in CS_2 or MoCl_5 in dichloromethane in various conditions gave invariably fully insoluble products, containing polychlorinated products as shown by mass spectroscopy. For **7** and **11**, it is possible that the presence of unsubstituted phenyl groups in *para*-position to the central benzene favours oligomerization. Furthermore the pyrimidine could maintain some solubility to the reaction intermediates by complexation with the aluminium, which would favour the polymerization and the chlorination. And



Scheme 2. Reagents and conditions: (a) DIPA, PPh_3 , $\text{PdCl}_2(\text{PPh}_3)_2$, CuI , DMF, μw , 115 °C, 93%; (b) Ph_2O , 260 °C, 44%; (c) $\text{FeCl}_3/\text{CH}_3\text{NO}_2$, DCM, 94%.



Scheme 3. Reagents and conditions: (a) DIPA, PPh_3 , $\text{PdCl}_2(\text{PPh}_3)_2$, CuI, DMF, μw , 115 °C, 61%; (b) Ph_2O , 260 °C, 52%; (c) $\text{FeCl}_3/\text{CH}_3\text{NO}_2$, DCM, 90%.

the presence of a dipole in the reaction products due to the pyrimidine ring decreases further the solubility of the final oligomers. The impossibility to fully cyclodehydrogenate **13** in the conditions used to prepare N-HSB from the very similar 1,2-dipyrimidyl-3,4,5,6-tetra(4-*tert*-butylphenyl)benzene (**Scheme 6**) is very puzzling. In contrast, reaction of **7**, **11** and **13** in dichloromethane with FeCl_3 /nitromethane gave very good yields of partially cyclodehydrogenated **1** (94%), **2** (90%) and **3** (85%), respectively (**Scheme 1**). In this reaction, and very similarly to what is observed for bipyrimidines (**Scheme 6**), the only C–C bonds that have been created are those in *ortho* position with respect to the pyrimidine nitrogen, leading to tribenzo[e,gh,j]perimidine cores (N-1/3HSB according to Draper's nomenclature).²⁶ The cyclodehydrogenation mechanisms are therefore probably similar and could involve arenium intermediates and/or the coordination of the iron.²⁴ Very similarly to what was observed with N-1/2HSB, attempts to complete the cyclodehydrogenation of these N-1/3HSB by reaction with AlCl_3 or FeCl_3 were unsuccessful. In contrast, reaction of the *per-tert*-butylated **16** with FeCl_3 , in the same conditions as above for **7**, **11**, **13**, provided the N-heterosuperbenzene **4** in 60% yield. The substitution of the hydrogen atom in position 2 in the pyrimidine ring (in **13**) by a *tert*-butyl group (in **16**) has a profound effect on the reactivity of these compounds. This substitution by a bulky, electron donating group has several consequences: (a) it increases the pyrimidine nucleophilicity, (b) blocks the position 2 and could hinder oligomerization at this position, (c) increases the


solubility of reagents, intermediates and products, and (d) prevents complexation by the nitrogen atoms by steric crowding. At this stage, further work is required to unveil the cyclodehydrogenation mechanisms leading to these very different chemical behaviours.

In conclusion, pyrimidyl-penta-phenylbenzenes have been synthesized by Diels–Alder addition of phenylethynylpyrimidines and tetraphenylcyclopentadienones under microwave irradiation. Scholl reactions of these compounds (ferric chloride/nitromethane/dichloromethane) led to two types of hetero polyaromatic hydrocarbons: (a) partial cyclization by creation of two C–C bonds *ortho* to the pyrimidine nitrogen atoms gave substituted tribenzo[e,gh,j]perimidine (N-1/3HSB) in high yields; (b) when the position 2 of the pyrimidine ring was substituted by a *tert*-butyl group, the Scholl reaction was complete and provided the first example of a diaza-hexa-*peri*-benzocoronene.


3. Experimental

3.1. General

¹H and ¹³C NMR spectra were recorded with a Bruker WF-250/300 MHz in CDCl_3 or CD_2Cl_2 solutions at 20 °C using solvent residue as the internal standard (CDCl_3 at δ_{H} : 7.25 ppm, δ_{C} : 77.0 ppm, CD_2Cl_2 at δ_{H} : 5.30 ppm, δ_{C} : 53.8 ppm). Mass Spectroscopy was performed with a Nermag R10-R10 (DCI). Elemental analyses were done by the Service d'Analyse de l'ICSN (Paris) and by the Service



Scheme 4. Reagents and conditions: (a) Ph_2O , 260 °C, 46%; (b) $\text{FeCl}_3/\text{CH}_3\text{NO}_2$, DCM, 85%.



Scheme 5. Reagents and conditions: (a) DIPA, PPh_3 , $\text{PdCl}_2(\text{PPh}_3)_2$, CuI, DMF, μw , 115°C , 78%; (b) Ph_2O , 260°C , 62%; (c) $\text{FeCl}_3/\text{CH}_3\text{NO}_2$, DCM, 60%.

d'Analyse du LCC (Toulouse). Note that, as observed by others, the carbon analysis of extended polyaromatic compounds **1–3** is inaccurate, due to incomplete combustion. The cyclopentadienones **10** and **12** were prepared according to Refs. **14** and **15**. THF was distilled from Na/benzophenone. Dichloromethane was distilled over calcium hydride. Nitromethane was dried on molecular sieves prior use. Other solvents and reagents were used as obtained in the best quality available. The microwave heating was carried out in closed vials with a CEM-Discover monomode microwave apparatus under the conditions (power, temperature, time) given here. After completion of the reaction, the vessel was cooled down rapidly to 60°C .

3.2. Synthesis of 5-phenylethynylpyrimidine **5**


A mixture of 102 mg of phenylacetylene (1 mmol) and 159 mg (1 mmol) 5-bromopyrimidine in 0.5 mL of dry DMF was degassed and 1.5 mL of dry DIPA, triphenylphosphine (23 mg, 0.087 mmol), $\text{Pd}(\text{P}(\text{C}_6\text{H}_5)_3)_2$ (15 mg, 0.021 mmol) and CuI (4 mg, 0.02 mmol) were added and the mixture was irradiated at 150 W, 115°C , 25 min. After cooling the brown-orange coloured suspension was extracted with 2×50 mL ether and washed with 2×25 mL saturated NH_4Cl , 2×25 mL water then dried with MgSO_4 . The crude product was further purified by chromatographic column in silica gel using

DCM as eluent. Drying under vacuum in ice gave pale yellow powder. Yield: 167 mg (93%).

^1H NMR (250 MHz, CD_2Cl_2): δ =7.41 (m, 3H, Ar-H), 7.57 (m, 2H, Ar-H), 8.86 (s, 2H, H4, H6), 9.11 (s, 1H, H2); ^{13}C NMR (75 MHz, CD_2Cl_2) δ =82.6, 95.9, 119.8, 121.9, 128.6, 129.4, 131.8, 156.8, 158.7; MS (CI, NH_3): m/z calcd for $\text{C}_{12}\text{H}_8\text{N}_2$: 180.0; found: 181 [$\text{M}^+ + 1$]. Anal. Calcd for $\text{C}_{12}\text{H}_8\text{N}_2$: C, 79.98; H, 4.47; N, 15.55. Found: C, 79.88; H, 4.44; N, 14.55.

3.3. Synthesis of 1-tert-butyl-4-ethynylbenzene **8**

To a degassed mixture of 3 g of 1-bromo-4-tert-butylbenzene (14.08 mmol), 485 mg of $\text{Pd}(\text{P}(\text{C}_6\text{H}_5)_3)_4$ (3%), 160 mg of CuI (6%) in 10 mL DIPA and 30 mL dry THF was added 2 mL of TMSA (1.38 g, 14 mmol). The mixture was irradiated at 150 W, 33 min, 110°C . After cooling the mixture was filtered over Celite; the Celite was washed with diethylether until colourless and the solvent evaporated under vacuum. The residue was then dissolved in ether, washed with HCl 1%, brine then dried on MgSO_4 . Purification by chromatography using hexane as eluent yielded 3.08 g of (4-tert-butylphenylethynyl)-trimethylsilane, which was then dissolved in 150 mL of methanol. To this solution was added 14 mL of aqueous 2.5 M NaOH. After 15 min of stirring the mixture was acidified to neutrality with 1 M HCl.



Scheme 6. Reagents and conditions from Refs. **25** and **26**. (a) AlCl_3 , CuCl_2 , CS_2 , rt, 49%; (b) $\text{FeCl}_3/\text{CH}_3\text{NO}_2$, DCM, rt; N-HSB (35%)+N-1/2HSB (32%); (c) $\text{AlCl}_3/\text{CuCl}_2$ or FeCl_3 .

Then the product was extracted with hexane (3×150 mL) and washed with water (100 mL), dried over MgSO₄, giving 1.96 g of pure **8** as orange oil (yield for the two steps: 88%). Analyses and spectroscopic data were in agreement with the literature.^{17,18}

3.4. Synthesis of **9**

To a degassed solution of **8** (158 mg, 1 mmol) and 5-bromopyrimidine (156 mg, 1 mmol) in 0.5 mL DMF and 1.5 mL dry DIPA were added triphenylphosphine (23 mg, 0.087 mmol), Pd(P(C₆H₅)₃)Cl₂ (15 mg, 0.0213 mmol) and CuI (4 mg, 0.021 mmol). The mixture was irradiated at 150 W, 115 °C, 25 min. After cooling the brown-orange suspension was extracted with 2×50 mL ether and filtered. The solution was washed with saturated NH₄Cl (2×25 mL), water (2×25 mL) and finally dried over MgSO₄. The crude product was purified by chromatography (silica gel, eluent: DCM/ethylacetate 5%). Yield: 145 mg of pale yellow powder (61%).

¹H NMR (300 MHz, CD₂Cl₂): δ=1.33 (s, 9H, *tert*-Bu), 7.29 (d, ³J_{H-H}=8.7 Hz, 2H, Ar-H), 7.51 (d, ³J_{H-H}=8.7 Hz, 2H, Ar-H), 8.84 (s, 2H, H₄, H₆), 9.09 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=31.0, 81.8, 96.1, 118.8, 120.6, 125.6, 131.4, 131.8, 152.9, 156.6, 158.5; MS (CI, NH₃): *m/z* calcd for C₁₆H₁₆N₂: 236.1; found: 237 [M⁺+1]. Anal. Calcd for C₁₆H₁₆N₂: C, 81.32; H, 6.32; N, 11.85. Found: C, 81.19; H, 6.99; N, 11.87.

3.5. Synthesis of **15**

To a degassed mixture of 158 mg (1 mmol) of 1-*tert*-butyl-4-ethynylbenzene (**8**), 5-bromo-2-*tert*-butylpyrimidine (**14**) (215 mg, 1 mmol) in 0.5 mL of dry DMF and 1.5 mL of dry DIPA were added triphenylphosphine (23 mg, 0.087 mmol), Pd(P(C₆H₅)₃)Cl₂ (15 mg, 0.021 mmol) and CuI (4 mg, 0.02 mmol), and the mixture was irradiated at 150 W, 115 °C, 25 min. After cooling the brown suspension was extracted with 2×50 mL of diethylether, filtered and washed with 2×25 mL saturated NH₄Cl, 2×25 mL water then dried with MgSO₄. The crude product was further purified by chromatographic column in silica gel using a gradient of petroleum ether/DCM as eluent (0–20% DCM). Yield: 205 mg (77%).

¹H NMR (300 MHz, CD₂Cl₂): δ=1.30 (s, 9H, *tert*-Bu), 1.37 (s, 9H, *tert*-Bu), 7.39 (d, ³J_{H-H}=8.7 Hz, 2H, Ar-H), 7.47 (d, ³J_{H-H}=8.7 Hz, 2H, Ar-H), 8.75 (s, 2H, H₄, H₆); ¹³C NMR (75 MHz, CD₂Cl₂): δ=29.2, 30.8, 82.5, 94.9, 116.2, 119.1, 125.6, 131.4, 152.6, 158.2, 175.6; MS (CI, NH₃): *m/z* calcd for C₂₀H₂₄N₂: 292.4; found: 293.3 [M⁺+1]. Anal. Calcd for C₂₀H₂₄N₂: C, 82.15; H, 8.27; N, 9.58. Found: C, 82.38; H, 8.39; N, 9.21.

3.6. General procedure for the synthesis of **7**, **11**, **13** and **16**

A mixture of phenylethynylpyrimidine (0.2 mmol) and cyclopentadienone (0.2 mmol) in diphenylether (1.5 g) was irradiated at 300 W, 260 °C, 45 min. After cooling, the dark-red mixture was diluted with dichloromethane (1 mL), poured in methanol (50 mL) and stirred; the off white precipitate was filtered, washed with methanol and vacuum dried. The product was recrystallized by partial evaporation (24 h) of a mixture of DCM (2.5 mL) and ethanol (5 mL).

3.6.1. Compound **7**

Yield 44%; ¹H NMR (300 MHz, CD₂Cl₂): δ=6.80–7.00 (m, 25H, Ar-H), 8.2 (s, 2H, H₄, H₆), 8.60 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=125.8, 126.2, 126.9, 127.5, 129.3, 131.5, 133.3, 135.1, 139.8, 140.3, 140.5, 142.0, 141.0, 155.6, 158.0, 206.8; MS (CI, NH₃): *m/z* calcd for C₄₀H₂₈N₂: 536.6; found: 537 [M⁺+1]. Anal. Calcd for C₄₀H₂₈N₂: C, 89.52; H, 5.26; N, 5.22. Found: C, 88.09; H, 5.26; N, 5.04.

3.6.2. Compound **11**

Yield 52%; ¹H NMR (300 MHz, CD₂Cl₂): δ=1.12 (br m, 27H, 3×*tert*-Bu), 6.70–6.97 (m, 20H, Ar-H), 8.14 (s, 2H, H₄, H₆), 8.59 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=31.2, 34.3, 123.6, 124.2, 124.2, 126.1, 126.8, 127.4, 131.1, 131.3, 131.5, 131.6, 135.4, 136.8, 137.4, 137.6, 140.2, 140.6, 140.8, 141.3, 142.4, 148.4, 149.1, 155.4, 158.0; MS (CI, NH₃): *m/z* calcd for C₅₂H₄₈N₂: 700.4; found: 701.4 [M⁺+1]. Anal. Calcd for C₅₂H₄₈N₂: C, 88.59; H, 7.43; N, 3.57. Found: C, 88.42; H, 7.21; N, 3.27.

3.6.3. Compound **13**

Yield: 46%; ¹H NMR (300 MHz, CD₂Cl₂): δ=1.10 (s, 45H, 5×*tert*-Bu), 6.70–6.77 (m, 10H, Ar-H), 6.84–6.90 (m, 6H, Ar-H), 6.937 (br d, ³J_{H-H}=8.4 Hz, 4H, Ar-H), 8.16 (s, 2H, H₄, H₆), 8.59 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=31.6, 34.7, 124.1, 124.6, 131.5, 131.7, 137.5, 138.1, 141.3, 141.7, 148.8, 149.4, 155.8, 158.5; MS (CI, NH₃): *m/z* calcd for C₆₀H₆₈N₂: 817.2; found: 817 [M⁺]. Anal. Calcd for C₆₀H₆₈N₂: C, 88.18; H, 8.39; N, 3.43. Found: C, 87.72; H, 8.33; N, 3.29.

3.6.4. Compound **16**

Yield: 62%; ¹H NMR (300 MHz, CD₂Cl₂): δ=1.08–1.11 (m, 54H, 6×*tert*-Bu), 6.72–6.75 (m, 10H, Ar-H), 6.83–6.93 (m, 10H, Ar-H), 8.04 (s, 2H, H₄, H₅); ¹³C NMR (75 MHz, CD₂Cl₂): δ=29.8, 31.3, 31.6, 124.0, 124.5, 131.5, 134.4, 137.7, 138.1, 141.5, 142.4, 148.8, 149.2, 157.8, 173.9; MS (CI, NH₃): *m/z* calcd for C₆₄H₇₆N₂: 872.6; found: 873.7 [M⁺+1]. Anal. Calcd for C₆₄H₇₆N₂: C, 88.02; H, 8.77; N, 3.21. Found: C, 88.03; H, 8.35; N, 3.15.

3.7. General procedure for cyclodehydrogenation of **7**, **11**, **13** and **16**

The hexaaryl compound was dissolved in anhydrous DCM. A stream of argon saturated with DCM was bubbled into the solution during the reaction to reduce chlorination. A solution of anhydrous FeCl₃ (1:30 mol ratio) in anhydrous nitromethane was quickly added. After a reaction time of 90 min, the reaction was quenched with methanol. Part of the dichloromethane was removed under reduced pressure. The precipitate was filtered and washed with aqueous HCl and methanol.

3.7.1. Compound **1**

Yield 94%; product was precipitated with methanol; ¹H NMR (300 MHz, CD₂Cl₂): δ=6.90 (m, 5H, Ar-H), 7.17 (m, 12H, Ar-H), 7.61 (m, 4H, Ar-H), 9.40 (d, ³J_{H-H}=9 Hz, 2H, Ar-H), 9.75 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=125.2, 125.4, 126.3, 126.6, 127.0, 127.1, 128.2, 128.9, 129.7, 130.5, 131.3, 131.8, 134.1, 139.8, 140.4, 142.7, 152.9, 155.6; MS (CI, CH₄): *m/z* calcd for C₄₀H₂₄N₂: 532.2; found: 533.2 [M⁺+1]. Anal. Calcd for C₄₀H₂₄N₂: C, 90.20; H, 4.54; N, 5.26. Found: C, 89.75; H, 4.73; N, 4.93. *R*_f: 0.52 in dichloromethane, silica gel.

3.7.2. Compound **2**

Yield 90%; product was precipitated with methanol; ¹H NMR (300 MHz, CD₂Cl₂): δ=1.15 (s, 9H, *tert*-Bu), 1.26 (s, 9H, *tert*-Bu), 1.45 (s, 9H, *tert*-Bu), 6.69 (d, ³J_{H-H}=8.4 Hz, 2H, Ar-H), 6.90 (d, ³J_{H-H}=8.4 Hz, 2H, Ar-H), 6.96 (d, ³J_{H-H}=8.4 Hz, 2H, Ar-H), 7.21–7.28 (m, 8H, Ar-H), 7.33 (dd, ³J_{H-H}=9.0 Hz, ⁴J_{H-H}=2.3 Hz, 1H, Ar-H), 7.59 (d, ³J_{H-H}=6.9 Hz, 1H, Ar-H), 7.64 (d, ³J_{H-H}=9 Hz, 1H, Ar-H), 7.79 (d, ³J_{H-H}=8.4 Hz, 1H, Ar-H), 9.41 (d, ³J_{H-H}=6.9 Hz, 1H, Ar-H), 9.48 (d, ⁴J_{H-H}=2.4 Hz, 1H, Ar-H), 9.76 (s, 1H, H₂); ¹³C NMR (75 MHz, CD₂Cl₂): δ=30.9, 30.0, 30.1, 123.0, 124.8, 125.1, 126.5, 126.9, 127.1, 128.1, 128.8, 129.5, 129.9, 130.7, 131.2, 131.3, 137.3, 139.8, 150.1, 155.3; MS (CI, CH₄): *m/z* calcd for C₅₂H₄₈N₂: 700.4; found: 701.4 [M⁺+1]. Anal. Calcd for C₅₂H₄₈N₂: C, 89.0; H, 6.90; N, 4.00. Found: C, 88.12; H, 7.01; N, 3.80. *R*_f: 0.45 in dichloromethane, silica gel.

3.7.3. Compound 3

Yield: 85%; product was precipitated with methanol; ^1H NMR (300 MHz, CD_2Cl_2): δ =1.14 (s, 9H, *tert*-Bu), 1.27 (s, 18H, $2\times$ *tert*-Bu), 1.47 (s, 18H, $2\times$ *tert*-Bu), 6.58 (d, $^3J_{\text{H-H}}=8.3$ Hz, 2H), 6.87 (d, $^3J_{\text{H-H}}=8.4$ Hz, 2H, Ar-H), 6.93 (d, $^3J_{\text{H-H}}=8.3$ Hz, 4H, Ar-H), 7.19 (d, $^3J_{\text{H-H}}=8.5$ Hz, 4H, Ar-H), 7.49 (dd, $^3J_{\text{H-H}}=9.1$ Hz, $^4J_{\text{H-H}}=2.1$ Hz, 2H, Ar-H), 7.83 (d, $^3J_{\text{H-H}}=9.1$ Hz, 2H, Ar-H), 9.43 (d, $^4J_{\text{H-H}}=2.1$ Hz, 2H, Ar-H), 10.09 (s, 1H, H₂); ^{13}C NMR (75 MHz, CD_2Cl_2) δ =29.7, 34.4, 35.2, 123.1, 125.1, 129.9, 130.4, 130.5, 130.9, 138.1, 138.9, 139.0, 139.8, 141.5, 150.3, 158.8; MS (CI, CH_4): m/z calcd for $\text{C}_{60}\text{H}_{64}\text{N}_2$: 812.5; found: 812.6 [M^+]. Anal. Calcd for $\text{C}_{60}\text{H}_{64}\text{N}_2$: C, 88.62; H, 7.93; N, 3.44. Found: C, 88.36; H, 7.72; N, 4.19. R_f : 0.70 in dichloromethane, silica gel.

3.7.4. Compound 4

Yield: 60%; product was further purified using column chromatography, silica gel, petroleum ether–dichloromethane (4: 1); ^1H NMR (300 MHz, CD_2Cl_2): δ =1.75 (s, 9H, *tert*-Bu), 1.85 (s, 27H, $3\times$ *tert*-Bu), 1.93 (s, 18H, $2\times$ *tert*-Bu), 9.41 (br s, 6H, Ar-H), 9.56 (d, $^4J_{\text{H-H}}=1.8$ Hz, 2H, Ar-H), 10.11 (d, $^4J_{\text{H-H}}=1.8$ Hz, 2H, Ar-H); ^{13}C NMR (75 MHz, CD_2Cl_2) δ =30.6, 31.7, 31.8, 125.9, 126.7, 128.5, 129.1, 130.3, 131.3, 132.3, 148.4, 150.0, 150.3, 166.2; MS (CI, CH_4): m/z calcd for $\text{C}_{64}\text{H}_{64}\text{N}_2$: 860.5; found: 861.7 [M^++1]. Anal. Calcd for $\text{C}_{64}\text{H}_{64}\text{N}_2$: C, 89.26; H, 7.49; N, 3.25. Found: C, 88.74; H, 7.04; N, 3.64. R_f : 0.83 in petroleum ether–dichloromethane (4:1), silica gel.

Acknowledgements

Partial support of this work by the European Commission within the projects CHIC (Contract no. IST-2001-33578) and PicoInside (Contract no. IST-015847) is gratefully acknowledged. One of the authors (S.N.) thanks the Department of Science and Technology, Government of India, for awarding a BOYSCAST fellowship.

References and notes

- Moresco, F.; Gourdon, A. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 8809–8814.
- Otero, R.; Rosei, F.; Naitoh, Y.; Jiang, P.; Thosttrup, P.; Gourdon, A.; Laegsgaard, E.; Stensgaard, L.; Joachim, C.; Besenbacher, F. *Nano Lett.* **2004**, *4*, 75–78.
- Gross, L.; Rieder, K. H.; Moresco, F.; Stojkovic, S. M.; Gourdon, A.; Joachim, C. *Nat. Mater.* **2005**, *4*, 892–895.
- Grill, L.; Moresco, F.; Rieder, K. H.; Stojkovic, S. M.; Gourdon, A.; Joachim, C. *Nano Lett.* **2005**, *5*, 859–863.
- Chiaravallotti, F.; Gross, L.; Rieder, K. H.; Stojkovic, S. M.; Gourdon, A.; Joachim, C.; Moresco, F. *Nat. Mater.* **2007**, *6*, 30–33.
- Thierney, J.-P.; Lidström, P. *Microwave Assisted Organic Synthesis*; Blackwell, CRC: Oxford, UK, 2005.
- de Souza, A. L. F.; da Silva, L. C.; Oliveira, B. L.; Antunes, O. A. C. *Tetrahedron Lett.* **2008**, *49*, 3895–3898.
- Togninelli, A.; Gevariya, H.; Alongi, M.; Botta, M. *Tetrahedron Lett.* **2007**, *48*, 4801–4803.
- Nilsson, P.; Ofsson, K.; Larhed, M. *Top. Curr. Chem.* **2006**, *266*, 103–144.
- Erdélyi, M.; Gogoll, A. *J. Org. Chem.* **2001**, *66*, 4165–4169.
- Feuerstein, M.; Doucet, H.; Santelli, M. *Tetrahedron Lett.* **2005**, *46*, 1717–1720.
- Soerensen, U. S.; Pombo-Villar, E. *Tetrahedron* **2005**, *61*, 2697–2704.
- Tang, B.-X.; Wang, F.; Li, J.-H.; Xie, Y.-X.; Zhang, M. B. *J. Org. Chem.* **2007**, *72*, 6294–6297.
- Iyer, V. S.; Wehmeier, M.; Diedrich Brand, J.; Keegstra, M. A.; Müllen, K. *Angew. Chem., Int. Ed.* **1997**, *36*, 1604–1607.
- Sadhukhan, S. K.; Viala, C.; Gourdon, A. *Synthesis* **2003**, *10*, 1521–1525.
- Perfetti, T. A.; Ogliaruso, M. A. *J. Org. Chem.* **1978**, *43*, 884–888.
- John, J. A.; Tour, J. M. *Tetrahedron* **1997**, *53*, 15515–15534.
- Leznoff, C. C.; Suchozak, B. *Can. J. Chem.* **2001**, *79*, 878–887.
- Wang, T.; Cloudsdale, I. S. *Synth. Commun.* **1997**, *27*, 2521–2526.
- Pews, R. G. *Heterocycles* **1990**, *31*, 109–114.
- Berresheim, A. J.; Muller, M.; Müllen, K. *Chem. Rev.* **1999**, *99*, 1747–1785 and references therein.
- (a) Rathore, R.; Burns, C. I. *J. Org. Chem.* **2003**, *68*, 4071–4074; (b) Chebny, V. J.; Gwengo, C.; Gardinier, J. R.; Rathore, R. *Tetrahedron Lett.* **2008**, *49*, 4869–4872.
- Rempala, P.; Kroulik, J.; King, B. T. *J. Org. Chem.* **2006**, *71*, 5067–5081.
- King, B. T.; Kroulik, J.; Robertson, C. R.; Rempala, P.; Cameron, C. L.; Korinek, J. D.; Gortari, L. M. *J. Org. Chem.* **2007**, *72*, 2279–2288.
- Draper, S. M.; Gregg, D. J.; Madathil, R. *J. Am. Chem. Soc.* **2002**, *124*, 3486–3487.
- Gregg, D. J.; Bothe, E.; Höfer, P.; Passaniti, P.; Draper, S. M. *Inorg. Chem.* **2005**, *44*, 5654–5660.
- Feng, X.; Wu, J.; Enkelmann, V.; Müllen, K. *Org. Lett.* **2006**, *8*, 1145–1148.