



# Water works: an efficient palladium-catalyzed cross-coupling reaction between boronic acids and bromoacetate with aminophosphine ligand

Zhi-Yong Peng<sup>a</sup>, Jian-Ping Wang<sup>a</sup>, Jiang Cheng<sup>b</sup>, Xiao-min Xie<sup>a</sup>, Zhaoguo Zhang<sup>a,c,\*</sup>

<sup>a</sup> School of Chemistry and Chemical Engineering, Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai 200240, China

<sup>b</sup> College of Chemistry and Materials Engineering, Wenzhou University, Wenzhou, 325027, China

<sup>c</sup> State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China

## ARTICLE INFO

### Article history:

Received 6 July 2010

Received in revised form 19 August 2010

Accepted 19 August 2010

Available online 9 September 2010

### Keywords:

Aminophosphine

Boronic acids

Ethyl bromoacetate

Palladium-catalyzed

Water

## ABSTRACT

Water greatly restrained the formation of self-coupling of boronic acids in a palladium-catalyzed cross-coupling reaction between boronic acids and ethyl bromoacetate with an aminophosphine ligand; good to excellent yields of cross-coupling product were obtained.


© 2010 Elsevier Ltd. All rights reserved.

## 1. Introduction

$\alpha$ -Aryl carboxylic acids are of significant importance since many of them are commercially used as non-steroidal anti-inflammatory and analgesic drugs.<sup>1</sup> Continuing demands exist for general and efficient procedures to supersede the current syntheses that need multiple steps, special or toxic reagents, and usually are conducted under harsh reaction conditions. Transition metal-catalyzed C–C bond-forming reactions, characterized by mild reaction conditions, that are high yielding and tolerate a wide range of functional groups, have been widely applied in organic syntheses.

There are two general reaction pathways to realize the arylation of corresponding esters to generate  $\alpha$ -aryl carboxylic acid derivatives reported by transition metal-catalyzed reactions (Scheme 1). Reactions via pathway (i), which involves a palladium-catalyzed coupling reaction between aryl halide electrophiles and enolates, were extensively studied by Buchwald<sup>2</sup> and Hartwig.<sup>3</sup> In these reactions, bulky, electron-rich phosphine ligands and/or carbene ligands were employed. Pathway (ii) is another promising alternative to achieve the coupling reaction by employing  $\alpha$ -halocarbonyl compounds and aryl metal as the substrates. Because arylboronic acids are readily available, fairly stable, and broadly compatible with an array of

functional groups,<sup>4</sup> they were usually used as the aryl metallic reagents.




Scheme 1.

Gooßen published the first paper that efficiently fulfilled the cross-coupling reaction between arylboronic acids and bromoacetic acid derivatives by employing trisnaphthyl phosphine as ligand<sup>5</sup> without the traditional utilization of highly toxic thallium salt as base.<sup>6</sup> Deng discovered that Cu<sub>2</sub>O served as co-catalyst to suppress the homo-coupling reaction of arylboronic acids and thus enhanced the efficiency for the cross-coupling reaction between arylboronic acids and ethyl bromoacetate or bromoacetamide with commercially available Pd(PPh<sub>3</sub>)<sub>4</sub> as catalyst.<sup>7</sup> Luo found that hydroquinone accelerated the reaction remarkably,<sup>8</sup> and this offers another entry to achieve cross-coupling of  $\alpha$ -halogenated carbonyl compounds and boronic acids rather than self-coupling of boronic acids.<sup>9</sup> Employing a nickel catalyst, Lei reported an efficient cross-coupling between arylboronic acids and ethyl bromoacetate.<sup>10</sup>

\* Corresponding author. Tel./fax: +86 21 54748925; e-mail address: [zhaoguo@sjtu.edu.cn](mailto:zhaoguo@sjtu.edu.cn) (Z. Zhang).

Aminophosphine ligands (Scheme 2) that we had applied to the Suzuki–Miyaura and Sonogashira coupling reactions<sup>11</sup> are easily-accessible, finely-tunable, efficient, and cheap ligands; however, they were only recently employed as ligands in transition metal-catalyzed cross-coupling reactions.<sup>12</sup> A few reports on the coordinative chemistry of aminophosphine compounds revealed that the function of amino groups was more diversified from alkoxy groups in phosphates.<sup>13</sup> In mono- and di-aminophosphines, alkyl and/or aryl amino groups served as strong electron-donating groups, making the phosphines stronger  $\sigma$ -donor ligands. Hartwig reported that both the enolate itself and ligand structure determine the connection pattern of the enolate to palladium during his study toward the reductive elimination of a palladium intermediate in palladium-catalyzed  $\alpha$ -arylation of carbonyl compounds.<sup>14</sup> Gooßen found that the sterically bulky and moderately electron-donating phosphines were suitable ligands for the cross-coupling reaction between arylboronic acids and bromoacetate. This enlightening discovery prompted us to test our aminophosphine ligands in this reaction.




Scheme 2.

## 2. Results and discussion

We chose phenylboronic acid (**1a**) and ethyl bromoacetate (**2**) as substrates, **L1a** as ligand and Pd<sub>2</sub>(dba)<sub>3</sub> as palladium species to investigate the cross-coupling reaction. A degassed Schlenk reaction tube was charged with ethyl bromoacetate (2 mmol), phenylboronic acid (3 mmol), potassium carbonate (6 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (0.05 mmol), ligand **L1a** (0.15 mmol), and THF (5 mL). The resulting suspension was refluxed for 8 h under nitrogen atmosphere. Besides the desired cross-coupling product, the homo-coupling product of phenylboronic acid, was also formed (Table 1, entry 1).

**Table 1**  
Effect of solvents using **L1a** as ligand<sup>a</sup>



	<b>1a</b>	<b>2</b>	<b>3a</b>	<b>4a</b>
Entry	Solvent		<b>3a</b> (%) <sup>b</sup>	<b>4a</b> (mg) <sup>c</sup>
1	THF		75	70
2	Dioxane		34	88
3	Toluene		33	33
4	Ethanol		66	49
5	DMF		16	51
6	Acetone		38	65
7	THF/H <sub>2</sub> O <sup>d</sup>		91	41
8	Dioxane/H <sub>2</sub> O <sup>d</sup>		65	67
9	Toluene/H <sub>2</sub> O <sup>d</sup>		70	78
10	Ethanol/H <sub>2</sub> O <sup>d</sup>		35	34

<sup>a</sup> All reactions were run with phenylboronic acid (3 mmol), ethyl bromoacetate (2 mmol), K<sub>2</sub>CO<sub>3</sub> (6 mmol), Pd (0.05 mmol), and ligand **L1a** (0.15 mmol) in 5 mL of solvent at 65 °C for 8 h.


<sup>b</sup> Isolated yields based on ethyl bromoacetate.

<sup>c</sup> Isolated mass.

<sup>d</sup> Organic solvent 4.5 mL, water 0.5 mL.

To optimize the reaction conditions, we first tested the solvent effect in reactions between phenylboronic acid and bromoacetate. We found that THF and ethanol were suitable solvents, and water plays a dramatic role in controlling the selectivity between homo-coupling and cross-coupling reactions. When 10% (v/v) of H<sub>2</sub>O was added, the yield of cross-coupling was greatly increased and the quantity of biphenyl was obviously decreased. This is quite different from previous reports, in which water was detrimental to the cross-coupling.<sup>5,8,9</sup>

To explain why water can improve the selectivity, the equilibrium between the C–Pd and O–Pd species should be considered (Scheme 3). After the oxidative addition of Pd(0) with ethyl bromoacetate, two possible palladium species may be formed. If a C–Pd bond-containing complex **A** is the only species, transmetalation in the presence of phenylboronic acid will proceed via route c to generate complex **D**, which is transformed to ethyl phenylacetate after reductive elimination. However, if complex **B** (a palladium enolate) is the predominant species (equilibrium a), or its presence is not negligible, the transmetalation step might be a different story. Because boron is oxophilic, the B(OH)<sub>2</sub> moiety could replace PdBr in complex **B** to generate a new boron-containing enolate and a palladium complex PhPdBr (**E**). **E** is formally the direct oxidative addition product of Pd(0) and bromobenzene, which undergoes further transmetalation with phenylboronic acid to yield biphenyl as the product. When water was added, there would exist OH<sup>−</sup> in the basic system, which displaces the halide in species **A** to give an hydroxopalladium species **C** in solution (equilibrium b),<sup>4a,15</sup> transmetalation in the presence of phenylboronic acid will proceed via route e to generate complex **D**, which also undergoes reductive elimination to give ethyl phenylacetate. The addition of water favors equilibrium b, and thus leads to the decrease of the formation of biphenyl product and the increase of cross-coupling product.




Scheme 3.

We then turned our attention to ligands, palladium species, and the effect of bases. The results are summarized in Table 2. Only **L1a** is a highly effective ligand in this reaction (Table 2, entry 1), while other ligands are inferior. In our previous work,<sup>11</sup> ligands **L1b** and **L1c** were workable ligands in the Suzuki–Miyaura cross-coupling reaction; however, they gave poor yields in this reaction (Table 2, entries 3–8).

When different palladium sources were tested, tris(dibenzylideneacetone) dipalladium(0) turned out to be the best palladium (0) precursors. Palladium(II) acetate gave products with moderate yields, PdCl<sub>2</sub>(MeCN)<sub>2</sub> and Pd(PPh<sub>3</sub>)<sub>4</sub> gave products with poor yields along with much more biaryl product (Table 2, entries 9–11).

The choice of the base also affects the product selectivity (Table 2, entries 12–15). K<sub>2</sub>CO<sub>3</sub> is the most suitable base, K<sub>3</sub>PO<sub>4</sub> and KF are also workable bases, NaOAc and NEt<sub>3</sub> are significantly less active.

**Table 2**The effects of ligands, palladium sources, and bases<sup>a</sup>


Entry	Palladium source	Ligand	Base	3a (%) <sup>b</sup>	4a (mg) <sup>c</sup>
1	Pd <sub>2</sub> (dba) <sub>3</sub>	L1a	K <sub>2</sub> CO <sub>3</sub>	91	41
2	Pd <sub>2</sub> (dba) <sub>3</sub>	L1b	K <sub>2</sub> CO <sub>3</sub>	27	139
3	Pd <sub>2</sub> (dba) <sub>3</sub>	L1c	K <sub>2</sub> CO <sub>3</sub>	15	232
4	Pd <sub>2</sub> (dba) <sub>3</sub>	L1d	K <sub>2</sub> CO <sub>3</sub>	18	182
5	Pd <sub>2</sub> (dba) <sub>3</sub>	L2a	K <sub>2</sub> CO <sub>3</sub>	42	81
6	Pd <sub>2</sub> (dba) <sub>3</sub>	L2b	K <sub>2</sub> CO <sub>3</sub>	25	143
7	Pd <sub>2</sub> (dba) <sub>3</sub>	L2c	K <sub>2</sub> CO <sub>3</sub>	11	231
8	Pd <sub>2</sub> (dba) <sub>3</sub>	L2d	K <sub>2</sub> CO <sub>3</sub>	16	147
9	Pd(OAc) <sub>2</sub>	L1a	K <sub>2</sub> CO <sub>3</sub>	79	87
10	PdCl <sub>2</sub> (MeCN) <sub>2</sub>	L1a	K <sub>2</sub> CO <sub>3</sub>	58	72
11	Pd(PPh <sub>3</sub> ) <sub>4</sub>	L1a	K <sub>2</sub> CO <sub>3</sub>	42	124
12	Pd <sub>2</sub> (dba) <sub>3</sub>	L1a	K <sub>3</sub> PO <sub>4</sub>	69	52
13	Pd <sub>2</sub> (dba) <sub>3</sub>	L1a	KF	62	33
14	Pd <sub>2</sub> (dba) <sub>3</sub>	L1a	NaOAc	32	11
15	Pd <sub>2</sub> (dba) <sub>3</sub>	L1a	NEt <sub>3</sub>	21	27

<sup>a</sup> All reactions were run with phenylboronic acid (3 mmol), ethyl bromoacetate (2 mmol), base (6 mmol), Pd (0.05 mmol) and ligand (0.15 mmol) in 5 mL of solvent (THF/H<sub>2</sub>O=9:1, v/v) at 65 °C for 8 h.

<sup>b</sup> Isolated yields based on ethyl bromoacetate.

<sup>c</sup> Isolated mass.

The generality and selectivity of the reaction were investigated using a number of arylboronic acids and alkenylboronic acids in combination with ethyl bromoacetate; the results are summarized in Table 3.

**Table 3**Pd-catalyzed synthesis of arylacetic acid derivatives<sup>a</sup>

Entry	R-B(OH) <sub>2</sub>	3	Yield <sup>b</sup> (%)
1	Phenyl	3a	90
2	<i>o</i> -Tolyl	3b	91
3	<i>m</i> -Tolyl	3c	90
4	<i>p</i> -Tolyl	3d	98
5	<i>p</i> -Chlorophenyl	3e	98
6	<i>p</i> -Formylphenyl	3f	92
7	<i>p</i> -Trifluoromethylphenyl	3g	85
8	<i>p</i> -Trifluoromethoxyphenyl	3h	84
9	<i>p</i> -Methoxyphenyl	3i	97
10	<i>p</i> -( <i>tert</i> -Butyl)phenyl	3j	90
11	1,3-Benzodioxol-5-yl	3k	97
12	2,6-Dimethylphenyl	3l	60
13	1-Naphthyl	3m	88
14	2-Methoxy-1-Naphthyl	3n	86
15	( <i>E</i> )-Styryl	3o	94
16	( <i>E</i> )-3-(Benzyloxy)prop-1-enyl	3p	84

<sup>a</sup> All reactions were run with phenylboronic acid (3 mmol), ethyl bromoacetate (2 mmol), K<sub>2</sub>CO<sub>3</sub> (6 mmol), Pd (0.05 mmol), and 1a (0.15 mmol) in 5 mL of solvent (THF/H<sub>2</sub>O=9:1, v/v), at 65 °C for 8 h.

<sup>b</sup> Isolated yields based on ethyl bromoacetate.

As illustrated in Table 3, most substrates gave good to excellent yields. Electron-poor and electron-rich compounds are equally suitable for the transformation (Table 3, entries 1–11), sterically hindered compounds gave moderate yields (Table 3, entry 12), 1-naphthylboronic acid, 2-methoxy-1-naphthylboronic acid, and alkenylboronic acid also gave good yields (Table 3, entries 13–16).

### 3. Conclusions

In conclusion, we have applied aminophosphine as an efficient ligand in a palladium-catalyzed cross-coupling reaction between

boronic acids and ethyl bromoacetate successfully. We optimized the reaction condition and found that water was crucial in the cross-coupling reaction. A number of arylboronic acids and alkenylboronic acids were tested and good to excellent yields were obtained.

## 4. Experimental section

### 4.1. Ethyl 2-phenylacetate (3a)

A degassed Schlenk reaction tube was charged in sequence with PhB(OH)<sub>2</sub> (3 mmol), potassium carbonate (6 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (0.05 mmol), ligand L1a (0.15 mmol), THF (4.5 mL), water (0.5 mL), and ethyl bromoacetate (2 mmol). The resulting mixture was degassed by freeze–thaw techniques and was kept stirring for 8 h at 65 °C. The reaction was quenched with water (5 mL), and the solution was extracted with ethyl acetate (3×10 mL). The organic layers were combined, and dried over sodium sulfate. The solvent evaporated in vacuo to give the crude product, which was purified by flash chromatography (PE then 2% EA/PE), biphenyl (41.2 mg, a white solid) and 3a (295.2 mg, 90%, a colorless oil) were obtained in sequence. Characterization of 3a: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.35–7.30 (m, 5H), 4.17 (q, *J*=7.0 Hz, 2H), 3.63 (s, 2H), 1.27 (t, *J*=7.2 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.6, 134.1, 129.2, 128.5, 127.0, 60.8, 41.4, 14.1.

### 4.2. Ethyl 2-*o*-tolylacetate (3b)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.19–7.17 (m, 4H), 4.16 (q, *J*=7.2 Hz, 2H), 3.63 (s, 2H), 2.32 (s, 3H), 1.25 (t, *J*=7.0 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.7, 137.1, 133.2, 130.6, 130.4, 127.6, 126.3, 61.0, 39.5, 19.8, 14.4.

### 4.3. Ethyl 2-*m*-tolylacetate (3c)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.23–7.20 (m, 1H), 7.11–7.07 (m, 3H), 4.16 (q, *J*=7.2 Hz, 2H), 3.57 (s, 2H), 2.34 (s, 3H), 1.25 (t, *J*=7.0 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.9, 138.4, 134.3, 130.2, 128.7, 128.0, 126.5, 61.0, 41.6, 21.6, 14.4.

### 4.4. Ethyl 2-*p*-tolylacetate (3d)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.18–7.11 (m, 4H), 4.14 (q, *J*=7.2 Hz, 2H), 3.56 (s, 2H), 2.32 (s, 3H), 1.24 (t, *J*=7.0 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.6, 136.4, 130.9, 129.0, 128.9, 60.6, 40.8, 20.9, 14.0.

### 4.5. Ethyl 2-(4-chlorophenyl)acetate (3e)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.30–7.26 (m, 2H), 7.23–7.20 (m, 2H), 4.16 (q, *J*=7.1 Hz, 2H), 3.58 (s, 2H), 1.25 (t, *J*=7.0 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.4, 133.2, 132.8, 130.9, 128.9, 61.2, 40.9, 14.4.

### 4.6. Ethyl 2-(4-formylphenyl)acetate (3f)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.0 (s, 1H), 7.85 (d, *J*=8.4 Hz, 2H), 7.46 (d, *J*=8.0 Hz, 2H), 4.16 (q, *J*=7.2 Hz, 2H), 3.71 (s, 2H), 1.26 (t, *J*=7.0 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 192.0, 170.8, 141.2, 135.5, 130.2, 130.1, 61.3, 41.6, 14.3.

### 4.7. Ethyl 2-(4-(trifluoromethyl)phenyl)acetate (3g)

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.58 (d, *J*=8.0 Hz, 2H), 7.40 (d, *J*=8.0 Hz, 2H), 4.16 (q, *J*=7.2 Hz, 2H), 3.68 (s, 2H), 1.26 (t, *J*=7.2 Hz, 3H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): 171.0, 138.3, 129.9, 129.4, 125.7, 123.0, 61.4, 41.3, 14.3.

**4.8. Ethyl 2-(4-(trifluoromethoxy)phenyl)acetate (3h)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.31 (d,  $J=8.4$  Hz, 2H), 7.17 (d,  $J=8.4$  Hz, 2H), 4.16 (q,  $J=7.2$  Hz, 2H), 3.62 (s, 2H), 1.26 (t,  $J=7.2$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.3, 148.5, 133.0, 130.8, 121.2, 119.4, 61.2, 40.7, 14.2. IR (KBr):  $\nu$  2358, 2339, 1735, 1509, 1253, 1217, 1153, 1030, 922  $\text{cm}^{-1}$ . HRMS (MICROMASS GCT-MS EI) exact mass calcd for ( $\text{C}_{11}\text{H}_{11}\text{F}_3\text{O}_3$ ):  $m/z$  248.0660, found  $m/z$  248.0633.

**4.9. Ethyl 2-(4-methoxyphenyl)acetate (3i)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.20 (m, 2H), 6.86 (m, 2H), 4.15 (q,  $J=6.8$  Hz, 2H), 3.79 (s, 3H), 3.55 (s, 2H), 1.25 (t,  $J=7.2$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 172.2, 158.9, 130.5, 126.5, 114.2, 61.0, 55.4, 40.7, 14.4.

**4.10. Ethyl 2-(4-*tert*-butylphenyl)acetate (3j)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.34 (d,  $J=8.4$  Hz, 2H), 7.22 (d,  $J=8.4$  Hz, 2H), 4.15 (q,  $J=7.2$  Hz, 2H), 3.58 (s, 2H), 1.31 (s, 9H), 1.26 (t,  $J=7.2$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 172.0, 150.0, 131.4, 129.2, 125.7, 61.0, 41.1, 34.7, 31.6, 14.5.

**4.11. Ethyl 2-(benzo[d][1,3]dioxol-5-yl)acetate (3k)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  6.79–6.72 (m, 3H), 5.94 (s, 2H), 4.15 (q,  $J=7.2$  Hz, 2H), 3.51 (s, 2H), 1.25 (t,  $J=7.0$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.9, 147.9, 146.9, 127.9, 122.6, 109.9, 108.4, 101.2, 61.1, 41.2, 14.4.

**4.12. Ethyl 2-(2,6-dimethylphenyl)acetate (3l)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.09–7.02 (m, 3H), 4.15 (q,  $J=7.2$  Hz, 2H), 3.68 (s, 2H), 2.33 (s, 6H), 1.24 (t,  $J=6.8$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.5, 137.4, 132.0, 128.3, 127.2, 60.9, 36.7, 20.5, 14.5.

**4.13. Ethyl 2-(naphthalen-1-yl)acetate (3m)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.01–7.98 (m, 1H), 7.86–7.77 (m, 2H), 7.54–7.41 (m, 4H), 4.15 (q,  $J=7.2$  Hz, 2H), 4.05 (s, 2H), 1.21 (t,  $J=6.8$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.9, 134.1, 132.4, 131.0, 129.0, 128.3, 128.2, 126.8, 126.1, 125.8, 124.1, 61.2, 39.6, 14.5.

**4.14. Ethyl 2-(2-methoxynaphthalen-1-yl)acetate (3n)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.73–7.70 (m, 2H), 7.62 (s, 1H), 7.42–7.39 (m, 1H), 7.34–7.29 (m, 1H), 7.10 (s, 1H), 4.16 (q,  $J=7.2$  Hz, 2H), 3.90 (s, 3H), 3.76 (s, 2H), 1.24 (t,  $J=6.8$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.9, 156.3, 134.4, 130.2, 128.9, 127.6, 126.7, 126.4, 126.3, 125.3, 124.0, 105.4, 60.9, 55.7, 37.0, 14.5.

**4.15. (E)-Ethyl 4-phenylbut-3-enoate (3o)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.38–7.20 (m, 5H), 6.49 (d,  $J=15.6$  Hz, 1H), 6.30 (dt,  $J=16.0$ , 7.2 Hz, 1H), 4.17 (q,  $J=7.2$  Hz, 2H), 3.24 (dd,  $J=6.8$ , 1.2 Hz, 2.0 H), 1.28 (t,  $J=7.2$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.8, 137.1, 133.6, 128.8, 127.7, 126.5, 122.1, 61.0, 38.7, 14.5.

**4.16. (E)-Ethyl 5-(benzyloxy)pent-3-enoate (3p)**

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.34–7.24 (m, 5H), 5.88–5.69 (m, 2H), 4.49 (s, 2H), 4.13 (q,  $J=7.0$  Hz, 2H), 4.00 (d,  $J=7.2$  Hz, 2H), 3.09 (d,  $J=6.6$  Hz, 2H), 1.25 (t,  $J=7.0$  Hz, 3H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ): 171.3, 138.0, 130.2, 128.2, 127.6, 127.4, 125.5, 71.9, 70.0, 60.5, 37.6, 14.0.

**Acknowledgements**

We thank the National Natural Science Foundation of China, the Science and Technology Commission of Shanghai Municipality, and the Education Commission of Shanghai Municipality for financial support.

**References and notes**

- (a) Lloyd-Jones, G. C. *Angew. Chem., Int. Ed.* **2002**, 41, 953–956; (b) Millard, A. A.; Rathke, M. W. *J. Am. Chem. Soc.* **1977**, 99, 4833–4835.
- (a) Palucki, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1997**, 119, 11108–11109; (b) Fox, J. M.; Huang, X.; Chieffi, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2000**, 122, 1360–1370; (c) Moradi, W. A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2001**, 123, 7996–8002; (d) Gaertzen, O.; Buchwald, S. L. *J. Org. Chem.* **2002**, 67, 465–475; (e) Vogl, E. M.; Buchwald, S. L. *J. Org. Chem.* **2002**, 67, 106–111; (f) Chae, J.; Yun, J.; Buchwald, S. L. *Org. Lett.* **2004**, 6, 4809–4812.
- (a) Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1997**, 119, 12382–12383; (b) Lee, S.; Beare, N. A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2001**, 123, 8410–8411; (c) Su, W.; Raders, S.; Verkade, J. G.; Liao, X.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2006**, 45, 5852–5855; (d) Hama, T.; Culkun, D. A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2006**, 128, 4976–4985.
- (a) Ishiyama, T.; Murata, M.; Miyaura, N. *J. Org. Chem.* **1995**, 60, 7508–7510; (b) Murphy, J. M.; Tzschucke, C. C.; Hartwig, J. F. *Org. Lett.* **2007**, 9, 757–760.
- Gooßen, L. J. *Chem. Commun.* **2001**, 669–670.
- Sato, M.; Miyaura, N.; Suzuki, A. *Chem. Lett.* **1989**, 1405–1408.
- (a) Liu, X.-X.; Deng, M.-Z. *Chem. Commun.* **2002**, 622–623; (b) Duan, Y.-Z.; Deng, M.-Z. *Tetrahedron Lett.* **2003**, 44, 3423–3426; (c) Duan, Y.-Z.; Zhang, J.-S.; Yang, J.; Deng, M.-Z. *Chin. J. Chem.* **2009**, 27, 179–182.
- Lu, T.-Y.; Xue, C.; Luo, F.-T. *Tetrahedron Lett.* **2003**, 44, 1587–1590.
- (a) Lei, A.; Zhang, X. *Tetrahedron Lett.* **2002**, 43, 2525–2528; (b) Shi, W.; Liu, C.; Yu, Z.; Lei, A. *Chem. Commun.* **2007**, 2342–2344.
- (a) Liu, C.; He, C.; Shi, W.; Chen, M.; Lei, A. *Org. Lett.* **2007**, 9, 5601–5604.
- (a) Cheng, J.; Wang, F.; Xu, J.; Pan, Y.; Zhang, Z. *Tetrahedron Lett.* **2003**, 44, 7095–7098; (b) Cheng, J.; Sun, Y.; Wang, F.; Guo, M.; Xu, J.; Pan, Y.; Zhang, Z. *J. Org. Chem.* **2004**, 69, 5428–5432; (c) Zhang, W.; Cheng, J.; Huang, Z.; Zhou, Q.; Chen, X.; Qian, J. *J. Chem. Res.* **2006**, 12, 781–782; (d) Zhang, W.; Wu, H.; Liu, Z.; Zhong, P.; Zhang, L.; Huang, X.; Cheng, J. *Chem. Commun.* **2006**, 46, 4826–4828.
- (a) Clarke, M. L.; Cole-Hamilton, D. J.; Woolins, J. D. *J. Chem. Soc., Dalton Trans.* **2001**, 2721–2723; (b) Schareina, T.; Kempe, R. *Angew. Chem., Int. Ed.* **2002**, 41, 1521–1523; (c) Urgaonkar, S.; Nagarajan, M.; Verkade, J. G. *Tetrahedron Lett.* **2002**, 43, 8921–8924; (d) Clarke, M. L.; Cole-Hamilton, D. J.; Slawin, A. M. Z.; Woolins, J. D. *Chem. Commun.* **2000**, 2065–2066; (e) Urgaonkar, S.; Nagarajan, M.; Verkade, J. G. *J. Org. Chem.* **2003**, 68, 452–459; (f) Priya, S.; Balakrishna, M. S.; Mobin, S. M.; McDonald, R. *J. Organomet. Chem.* **2003**, 688, 227–235; (g) David, B.; Vladica, B.; Kurt, M.; Karl, K. *Organometallics* **2006**, 25, 3817–3823; (h) Bolliger, J. L.; Blaque, O.; Frech, C. M. *Angew. Chem., Int. Ed.* **2007**, 46, 6514–6517; (i) Cho, S. D.; Kim, H. K.; Yim, H. S.; Kim, M. R.; Lee, J. K.; Kim, J. J.; Yoon, Y. *J. Tetrahedron* **2007**, 63, 1345–1352; (j) Biricik, N.; Durap, F.; Kayan, C.; Gungum, B.; Gurbuz, N.; Ozdemir, I.; Ang, W. H.; Fei, Z.; Scopelliti, R. *J. Organomet. Chem.* **2008**, 693, 2693–2699; (k) Bolliger, J. L.; Frech, C. M. *Adv. Synth. Catal.* **2009**, 351, 891–902.
- (a) Rømming, C.; Songstad, J. *Acta Chem. Scand. Ser. A* **1978**, 32, 689–699; (b) Rømming, C.; Songstad, J. *Acta Chem. Scand. Ser. A* **1979**, 33, 187–197; (c) Rømming, C.; Songstad, J. *Acta Chem. Scand. Ser. A* **1982**, 36, 665–671; (d) Moloy, K. G.; Petersen, J. L. *J. Am. Chem. Soc.* **1995**, 117, 7696–7710; (e) Socol, S. M.; Jacobson, R. A.; Verkade, J. G. *Inorg. Chem.* **1984**, 23, 88–94; (f) Zijp, E. J.; Vlugt, J. I. v. d.; Tooke, D. M.; Spek, A. L.; Vogt, D. *J. Chem. Soc., Dalton Trans.* **2005**, 512–517.
- Culkun, D. A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2001**, 123, 5816–5817.
- (a) Grushin, V. V.; Alper, H. *Organometallics* **1993**, 12, 1890–1901; (b) Moriya, T.; Miyaura, N.; Suzuki, A. *Synlett* **1994**, 149–151; (c) Sodeoka, M.; Ohrai, K.; Shibasaki, M. *J. Org. Chem.* **1995**, 60, 2648–2649.