



Improved synthesis of UDP-2-(2-ketopropyl)galactose and a first synthesis of UDP-2-(2-ketopropyl)glucose for the site-specific linking of biomolecules via modified glycan residues using glycosyltransferases

Andrés E. Dulcey^{a,*}, Pradman K. Qasba^b, Jeffrey Lamb^a, Gary L. Griffiths^a

^aImaging Probe Development Center, National Heart, Lung, and Blood Institute, National Institutes of Health, 9800 Medical Center Drive, Rockville, MD 20850, USA

^bStructural Glycobiology Section, Nanobiology Program, Center for Cancer Research, NCI-Frederick, Frederick, MD 21702, USA

ARTICLE INFO

Article history:

Received 10 December 2010

Received in revised form 25 January 2011

Accepted 26 January 2011

Available online 2 February 2011

Keywords:

Carbohydrate chemistry

Organic synthesis

Modified sugars

Glycosyltransferases

ABSTRACT

The potential of wild-type and mutant glycosyltransferases to produce glycoconjugates carrying sugar moieties with chemical handles has made it possible to conjugate biomolecules with orthogonal reacting groups at specific sites. The synthesis of UDP-2-(2-ketopropyl)galactose has been previously carried out, albeit with difficulty and low efficiency. A modified approach has been developed for the synthesis of UDP-2-(2-ketopropyl)glucose and UDP-2-(2-ketopropyl)galactose, allowing better access to the desired test compounds, the UDP-2-(2-ketopropyl)glucose and UDP-2-(2-ketopropyl)galactose analogs were synthesized in eight steps and 4.8% and 5.3% overall yield, respectively, an improvement over the first generation synthesis involving eight steps and an overall yield of 0.7%.

Published by Elsevier Ltd.

1. Introduction

Structural information on glycosyltransferases has revealed that the specificity of these enzymes can be broadened to include modified sugars with chemical handles that can then be utilized for site-specific conjugation chemistry. As such, this approach represents a powerful general method for the site-specific modification of glycoproteins. Substitution of Tyr289 to Leu in the catalytic pocket of bovine β -1,4-galactosyltransferase generates a novel glycosyltransferase, which can transfer not only galactose, but also modified galactosyl moieties, such as GalNAc¹ and C2-modified galactose containing various chemical handles, from the corresponding uridine diphosphate (UDP) derivatives.^{2,3} Similarly, the wild-type *N*-acetylgalactosaminyltransferase, which naturally transfers GalNAc from UDP/GalNAc, can also transfer C2-modified galactose units from their UDP-derivative to the Ser/Thr residue of a polypeptide acceptor substrate tagged as a fusion peptide to a non-glycoprotein.⁴ The potential of wild-type and mutant glycosyltransferases to produce glycoconjugates carrying sugar moieties with chemical handles has made it possible to conjugate biomolecules with orthogonal reacting groups at specific sites.⁵ This methodology has enabled the

assembly of novel bio-nanoparticles useful for targeted drug delivery and as MRI contrast agents.^{6,7}

The synthesis of UDP-2-(2-ketopropyl)glucose (Fig. 1, **8-Glc**) has not been previously reported, while the synthesis of UDP-2-(2-ketopropyl)galactose (Fig. 1, **8-Gal**) was reported in low yields. The




Fig. 1. UDP-2-(2-ketopropyl)glucose **8-Glc** and UDP-2-(2-ketopropyl)galactose **8-Gal**.

* Corresponding author. Tel.: +1 301 217 5759; fax: +1 301 217 5779; e-mail address: dulceyan@mail.nih.gov (A.E. Dulcey).

lack of efficient syntheses of these substrates represents a roadblock to the continuing development of this promising technology, as both are required for the site-specific conjugation of biomolecules via glycan residues using glycosyltransferases. The linking technique enables conjugation of the glycoprotein-appended 2-(2-ketopropyl) sugar to aminoxy derivatives, such as aminoxybiotin and has wide potential for future conjugation of cytotoxic drugs, cytokines, radionuclides for imaging and radioimmunotherapy, and lipids for the assembly of liposomes for targeted drug delivery.^{6,7}

2. Results and discussion

'2-(2-Ketopropyl)sugars', which are C2-carbon isosteres of 2-*N*-acetamidoglycans, were first introduced by Bertozzi⁸ as novel analogs possessing a ketone group for chemoselective reaction with aminoxy- or hydrazido-reagents. The first generation synthesis of UDP-2-(2-ketopropyl)galactose (**8-Gal**) was carried out by Hsieh-Wilson's group,² albeit with difficulty and low efficiency. The modified approach for the synthesis of UDP-2-(2-ketopropyl)glucose (**8-Glc**) and UDP-2-(2-ketopropyl)galactose (**8-Gal**), developed at the *Imaging Probe Development Center* is displayed in *Scheme 1*.

Thus, from commercially available tri-*O*-acetyl-*D*-glucal (**1-Glc**) or tri-*O*-acetyl-*D*-galactal (**1-Gal**), equatorial iodination was effected with *N*-iodosuccinimide (NIS) and acetic acid in dry dichloromethane to afford equatorial iodosugars **2-Glc** and **2-Gal** in 25% and 35% yield, respectively, with an α : β ratio of \sim 1:10 for the Glc series and \sim 1:2 for the Gal series. Keck radical allylation⁹ with methylallyltributylstannane and catalytic AIBN (2,2'-azobis(2-methylpropionitrile)) afforded the desired equatorial methallyl compounds **3-Glc** and **3-Gal** in 70% and 68% yield, respectively, along with <5% of the axial epimer, separable chromatographically.

Anomeric deprotection was effected with hydrazine acetate in DMF, obtaining lactol **4-Glc** and **4-Gal**, which was coupled with dibenzyl *N,N*-diisopropylphosphoramidite at -30 °C. Compounds

5-Glc and **5-Gal** were not isolated. After a quick workup, **5-Glc** and **5-Gal** were subjected to ozonolysis at -78 °C followed by the addition of dimethyl sulfide to obtain α -*D*-glucopyranoside phosphate **6-Glc** in 68% yield from **4-Glc**, and α -*D*-glucopyranoside phosphate **6-Glc** in 60% yield from **4-Gal**.


Hydrogenolysis of the benzyl groups in **6-Glc** and **6-Gal**, followed by condensation with UMP-morpholidate and deprotection of the acetyl groups afforded the target UDP-2-(2-ketopropyl)glucose (**8-Glc**) in 49% yield from **6-Glc**, and UDP-2-(2-ketopropyl)galactose (**8-Gal**) in 53% yield from **6-Gal**.

A major obstacle was presented by the sequence illustrated in *Scheme 2*, employed in the first generation synthesis.² The sequence was troubled by low-yielding transformations and unwanted byproduct formation due to the instability of the intermediates, along with a need to perform two oxidation steps.


The yield reported in the first generation synthesis from intermediate **3-Gal** to intermediate **6-Gal** was of only 26%. These shortcomings would prove detrimental if the synthesis was to be scaled up eventually, as requested. The new approach shown in *Scheme 1* applies redox economy in organic synthesis,¹⁰ and illustrates how unwanted byproduct formation (Aldol condensation) was avoided by performing just one combined oxidation step¹¹ after the phosphoramidite coupling had taken place (**4** to **5** to **6** in *Scheme 1*). This aided in scale up and afforded increased yields for key intermediates **6-Gal** and **6-Glc** with the yields from intermediate **3** to intermediate **6** now 42% in the galactose series and 54% in the glucose series (*Scheme 1*).

3. Conclusion

The modified approach developed for the synthesis of UDP-2-(2-ketopropyl)glucose and UDP-2-(2-ketopropyl)galactose, herein described in full experimental detail, has allowed much improved access to the desired test substrates. The UDP-2-(2-ketopropyl)



Scheme 1. Reagents and conditions. (a) *N*-Iodosuccinimide (1.5 equiv), AcOH (2 equiv) CH₂Cl₂, rt, 45 min. (b) Methylallyltributylstannane (2 equiv), AIBN (0.15 equiv), benzene, reflux, 3 h. (c) Hydrazine acetate (1.2 equiv), DMF, rt, 2 h. (d) 1-*H*-Tetrazole (4 equiv), dibenzyl *N,N*-diisopropylphosphoramidite (2.5 equiv), CH₂Cl₂, -30 °C to rt 3 h. (e) (i) Ozone, CH₂Cl₂–CH₃OH, -78 °C, 40 min. Then (ii) Me₂S (excess), -78 °C to rt overnight. (f) (1) Pd/C (10%, 10 wt %), triethylamine (0.6 equiv), CH₃OH, H₂, 24 h. (2) UMP-morpholidate 4-morpholine-*N,N'*-dicyclohexylcarboxamide salt (1.5 equiv), 1-*H*-tetrazole (4 equiv), pyridine, 4 Å MS, 3 days. (g) CH₃OH, H₂O, Et₃N (4:2:1), 24 h.



Scheme 2. Problematic step and unwanted side-product generation via Aldol condensation.

glucose analog **8-Glc** was synthesized in eight steps and 4.8% overall yield, and the UDP-2-(2-ketopropyl)galactose analog **8-Gal** was obtained in the same number of steps with and overall yield of 5.3%. With the latter previously described compound a 7.5-fold yield improvement was obtained over the first generation synthesis,² which also involved eight steps but resulted in an overall yield of only 0.7%.

4. Experimental section

4.1. General

All commercially available organic precursors and dry solvents were obtained from Sigma–Aldrich, and used as received unless otherwise noted. Reactions were magnetically stirred under an argon atmosphere and monitored by thin layer chromatography (TLC) with 0.25 mm Sigma–Aldrich pre-coated aluminum-backed silica gel plates with fluorescent indicator. TLC visualization was achieved using 254 nm or 360 nm UV lamp detection and/or staining with cerium molybdate (Hannesian's stain), phosphomolybdic acid (PMA), or potassium permanganate. Flash column chromatography was performed on an AnaLogix IntelliFlash 280 system, using Biotage® SNAP Cartridges and SNAP Samplet Cartridges with KP-Silica 60 μ m. Analytical HPLC analyses were performed on an Agilent 1200 Series instrument equipped with multi-wavelength detectors using a Zorbax Stable Bond C-18 column (4.6 \times 50 mm, 3.5 μ m) with a flow rate of 0.5 mL/min or 1.0 mL/min. Solvent A was 0.05% trifluoroacetic acid (TFA) in water (H₂O), solvent B was 0.05% TFA in acetonitrile (ACN), and a linear gradient of 5–95% B over 10 min was used. ESI or APCI mass spectrometry (MS) was performed on an LC/MSD TrapXCI Agilent Technologies instrument or on a 6130 Quadrupole LC/MS Agilent Technologies instrument equipped with a diode array detector. High resolution mass spectrometry (HRMS) was performed by The Scripps Research Institute Center for Metabolomics and Mass Spectrometry (La Jolla, CA). ¹H, ¹³C and ³¹P NMR spectra were recorded on a Varian spectrometer operating at 400 MHz, 100 MHz and 162 MHz, respectively. Chemical shifts are reported relative to either chloroform (δ 7.26), dimethyl sulfoxide (δ 2.50), acetone (δ 2.05) or deuterium oxide (δ 4.79) for ¹H NMR and chloroform (δ 77.0), dimethyl sulfoxide (δ 39.5) or acetone (δ 29.8) for ¹³C NMR.

4.1.1. 1,3,4,6-Tetra-O-acetyl-2-deoxy-2-iodo-D-glucopyranose (2-Glc). *N*-Iodosuccinimide (NIS) (2.04 g, 9.07 mmol) was added at room temperature to a solution of tri-O-acetyl-D-galactal (1.51 g, 5.55 mmol) and acetic acid (AcOH) (635 μ L, 11.1 mmol) in dry dichloromethane (CH₂Cl₂) (20 mL) in an aluminum foil-wrapped flask. The resulting reaction mixture was allowed to stir in the dark for 45 min, after which TLC [2:1 hexane (Hex)/ethyl acetate (EtOAc)] showed completion. The reaction was quenched by the addition of a saturated aqueous solution of sodium thiosulfate (Na₂S₂O₃) and allowed to stir for 15 min. The reaction mixture was diluted with CH₂Cl₂ and washed with Na₂S₂O₃ (one time), H₂O (one time) and brine, dried over magnesium sulfate (MgSO₄), and concentrated. The

crude mixture was purified by flash column chromatography on silica gel eluted with a gradient of 3:1 \rightarrow 2:1 Hex/EtOAc to afford the unwanted 2-iodomannose isomer (lower *R_f*) as the major product, and the desired 2-iodoglucose derivative **2-Glc** (higher *R_f*) as the minor product (634 mg, 25% yield) as a clear syrup with an α : β anomeric ratio of 1:10. β anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.85 (d, *J*=9.6 Hz, 1H), 5.32 (t, *J*=10.0 Hz, 1H), 4.99 (t, *J*=10.0 Hz, 1H), 4.30 (dd, *J*=12.4, 4.4 Hz, 1H), 4.07 (dd, *J*=12.4, 2.0 Hz, 1H), 3.97 (t, *J*=11.2 Hz, 1H), 3.86 (ddd, *J*=10.0, 4.4, 2.0 Hz, 1H), 2.15 (s, 3H), 2.08 (s, 3H), 2.06 (s, 3H), 2.00 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.5, 169.7, 169.4, 168.4, 93.8, 75.1, 72.9, 68.4, 61.4, 25.6, 20.6, 20.5; MS (*m/z*)=459.7 (*M*+1)⁺; HRMS (ESI) calcd for C₁₄H₁₉IO₉Na⁺ [*M*+Na]⁺ 480.9966, found 480.9966.

4.1.2. 1,3,4,6-Tetra-O-acetyl-2-deoxy-2-iodo-D-galactopyranose (2-Gal). Tri-O-acetyl-D-galactal (2.30 g, 8.45 mmol), NIS (2.28 g, 10.1 mmol), AcOH (1 mL, 16.9 mmol) and, CH₂Cl₂ (25 mL) to obtain **2-Gal** (1.34 g, 34% yield) as syrup, minor product with higher *R_f*. β anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.88 (d, *J*=9.2 Hz, 1H), 5.32 (m, 1H), 5.22 (d, *J*=4 Hz, 1H), 5.12 (dd, *J*=12.0, 3.6 Hz, 1H), 4.33 (t, *J*=6.8 Hz, 1H), 4.11 (m, 1H), 4.04 (m, 1H), 2.15 (s, 3H), 2.11 (s, 3H), 2.04 (s, 3H), 2.01 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.2, 169.8, 169.5, 168.4, 93.3, 73.0, 72.3, 68.8, 61.1, 24.7, 20.5, 19.5; MS (*m/z*)=459.1 (*M*+1)⁺; HRMS (ESI) calcd for C₁₄H₁₉IO₉Na⁺ [*M*+Na]⁺ 480.9966, found 480.9973.

4.1.3. 1,3,4,6-Tetra-O-acetyl-2-deoxy-2-(methallyl)-D-glucopyranose (3-Glc). A solution of **2-Glc** (1.30 g, 2.84 mmol) and methallyltributylstannane⁹ (1.96 g, 5.67 mmol) in benzene (15 mL) was thoroughly degassed by bubbling argon through it for 20 min. 2,2'-Azobis(2-methylpropionitrile) (AIBN) (70.0 mg, 0.43 mmol) was then added in one portion and degassing continued for another 5 min. The resulting reaction mixture was refluxed for 3 h, after which TLC (2:1 Hex/EtOAc) showed completion. The reaction mixture was allowed to cool to room temperature and concentrated under reduced pressure. The residue was taken up in acetonitrile and washed with hexanes (three times). The acetonitrile layer was concentrated and the residue purified by flash column chromatography on silica gel using 2:1 Hex/EtOAc to afford **3-Glc** as a clear oil (768 mg, 70% yield). β anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.54 (d, *J*=8.8 Hz, 1H), 5.01 (t, *J*=6.8 Hz, 1H), 4.67 (s, 1H), 4.62 (s, 1H), 4.31 (dd, *J*=12.4, 4.4 Hz, 1H), 4.06 (dd, *J*=12.4, 2.4 Hz, 1H), 3.79–3.74 (m, 1H), 2.10–2.07 (m, 10H), 2.00 (s, 3H), 1.98 (s, 3H), 1.73 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.2, 169.8, 168.9, 142.3, 112.2, 94.3, 72.3, 69.1, 61.9, 42.1, 37.1, 21.6, 20.8, 20.7, 20.6; MS (*m/z*)=387.9 (*M*+1)⁺; HRMS (ESI) calcd for C₁₈H₂₆O₉Na⁺ [*M*+Na]⁺ 409.1469, found 409.1467.

4.1.4. 1,3,4,6-Tetra-O-acetyl-2-deoxy-2-(methallyl)-D-galactopyranose (3-Gal). **2-Gal** (2.20 g, 4.80 mmol), methallyltributylstannane⁸ (4.14 g, 12.0 mmol), AIBN (118 mg, 0.72 mmol), and benzene (30 mL) afforded **3-Gal** (1.26 g, 68% yield) as a clear oil. β anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.54 (d, *J*=9.2 Hz, 1H), 5.36–5.33 (m, 1H), 5.26 (d, *J*=2.4 Hz, 1H), 4.82 (dd, *J*=11.6, 3.2 Hz, 1H), 4.66 (s, 1H), 4.59 (s, 1H), 4.16–4.03 (m, 3H), 3.94 (t, *J*=6.8 Hz, 1H), 2.12–2.09 (m, 4H), 2.05 (s, 3H), 2.01 (s, 3H), 1.95 (s, 3H), 1.72 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.2, 169.9, 168.8, 142.6, 113.6, 113.0, 112.1, 93.7, 71.4, 68.9, 67.0, 66.8, 37.9, 36.8, 35.7, 21.8, 20.9, 20.7, 20.3, 19.7; MS (*m/z*)=387.5 (*M*+1)⁺; HRMS (ESI) calcd for C₁₈H₂₆O₉Na⁺ [*M*+Na]⁺ 409.1469, found 409.1475.

4.1.5. 3,4,6-Tri-O-acetyl-2-deoxy-2-(methallyl)-D-glucopyranose (4-Glc). Hydrazine acetate (102 mg, 1.11 mmol) was added in one portion to a solution of **3-Glc** (357 mg, 0.92 mmol) in *N,N*-dimethylformamide (DMF) (5 mL). The reaction was allowed to stir at room temperature for 2 h. TLC (2:1 Hex/EtOAc) showed completion. The reaction mixture was partitioned between H₂O and EtOAc,

and the organic layer washed with H₂O (three times), followed by brine, dried over MgSO₄, and concentrated under reduced pressure. Product was purified by flash column chromatography on silica gel with 2:1 Hex/EtOAc to afford the lactol **4-Glc** as a syrup (265 mg, 83% yield). α anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.27 (t, J =10.0 Hz, 1H), 5.15 (d, J =3.2 Hz, 1H), 4.96 (t, J =9.6 Hz, 1H), 4.78–4.68 (m, 3H), 4.26–4.21 (m, 1H), 4.17–4.12 (m, 1H), 4.09–4.02 (m, 1H), 3.33 (d, J =3.6 Hz, 1H), 2.07–2.04 (m, 5H), 2.01 (s, 3H), 2.01 (s, 3H), 1.98 (s, 3H), 1.96 (s, 3H), 1.69 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.9, 170.6, 170.1, 142.4, 114.0, 112.5, 93.6, 72.7, 71.3, 70.5, 69.3, 68.5, 67.7, 65.7, 62.8, 42.2, 40.0, 35.9, 23.1, 21.6, 20.6, 19.7; MS (m/z)=345.8 (M+1)⁺; HRMS (ESI) calcd for C₁₆H₂₄O₈Na⁺ [M+Na]⁺ 367.1363, found 367.1368.

4.1.6. 3,4,6-Tri-O-acetyl-2-deoxy-2-(methallyl)-D-galactopyranose (4-Gal). Hydrazine acetate (145 mg, 1.56 mmol), **3-Gal** (550 mg, 1.42 mmol), and DMF (5 mL) to obtain **4-Gal** (343 mg, 70% yield). α anomer: ¹H NMR (400 MHz, CDCl₃) δ 5.31 (d, J =2.8 Hz, 1H), 5.23 (t, J =3.2 Hz, 1H), 5.14 (dd, J =11.6, 3.2 Hz, 1H), 4.77 (d, J =10.0 Hz, 2H), 4.41 (t, J =6.4 Hz, 1H), 4.13–4.11 (m, 1H), 4.07 (t, J =5.6 Hz, 1H), 2.98 (d, J =3.6 Hz, 1H), 2.14–2.10 (m, 5H), 2.03 (s, 3H), 1.97 (s, 3H), 1.75–1.69 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.4, 142.2, 112.5, 93.2, 69.5, 67.1, 66.5, 62.4, 36.4, 35.3, 22.6, 22.3, 20.7 ($\times 2$); MS (m/z)=345.6 (M+1)⁺; HRMS (ESI) calcd for C₁₆H₂₄O₈Na⁺ [M+Na]⁺ 367.1363, found 367.1369.

4.1.7. Dibenzyl (2-acetyl-2-deoxy-3,4,5-tri-O-acetyl- α -D-glucopyranosyl) phosphate (6-Glc). A solution of lactol **4-Glc** (250 mg, 0.73 mmol) and 1H-tetrazole (200 mg, 2.90 mmol) in dry CH₂Cl₂ (5 mL) was cooled to –30 °C and treated with dibenzyl *N,N*-diisopropylphosphoramidite (627 mg, 1.81 mmol). The resulting reaction mixture was allowed to slowly warm from –30 °C to room temperature for 3 h. TLC (2:1 Hex/EtOAc) showed complete disappearance of lactol **4-Glc**. The reaction mixture was diluted with Et₂O and washed with cold brine (two times), dried over MgSO₄, and concentrated under reduced pressure. The residue was then taken up in dry CH₂Cl₂ (6 mL) and cooled to –78 °C. A stream of ozone was bubbled through the reaction mixture until a light blue color became evident (~30–40 min). Argon was then bubbled through the reaction mixture until the blue color disappeared and Me₂S (1 mL) added. The resulting reaction mixture was allowed to stir overnight as the temperature rose to ambient. The solvent was removed under reduced pressure and the residue purified by flash column chromatography: silica gel, 1:1 Hex/EtOAc to afford **6-Glc** as a syrup (298 mg, 68% yield). ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.32 (m, 10H), 5.81 (dd, J =6.4, 3.2 Hz, 1H), 5.37 (t, J =3.2 Hz, 1H), 5.23 (t, J =10.2 Hz, 1H), 5.09–4.97 (m, 4H), 4.21 (t, J =4.4 Hz, 1H), 4.14–4.06 (m, 2H), 4.04–3.99 (m, 1H), 3.88 (dd, J =12.4, 2.0 Hz, 1H), 3.31 (d, J =2.4 Hz, 1H), 2.13 (s, 3H), 2.03 (s, 3H), 2.00 (s, 3H), 1.92 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 205.5, 170.5, 170.4, 169.5, 135.3, 128.7 ($\times 2$), 128.0, 97.1, 70.5, 69.7 ($\times 2$), 69.6, 68.6, 61.6, 39.7, 29.7, 20.6, 20.5; ³¹P NMR (162 MHz, CDCl₃) δ –2.31; MS (m/z)=607.9 (M+1)⁺; HRMS (ESI) calcd for C₂₉H₃₅O₁₂P⁺ [M+H]⁺ 607.1939, found 607.1930.

4.1.8. Dibenzyl (2-acetyl-2-deoxy-3,4,5-tri-O-acetyl- α -D-galactopyranosyl) phosphate (6-Gal). **4-Gal** (60 mg, 0.17 mmol), 1H-tetrazole (49 mg, 0.70 mmol), dibenzyl *N,N*-diisopropylphosphoramidite (150 mg, 0.44 mmol), ozone, and Me₂S (1 mL) to afford **6-Gal** as a syrup (63 mg, 60% yield). ¹H NMR (400 MHz, CDCl₃) δ 7.35–7.30 (m, 10H), 5.86 (dd, J =6.0, 2.8 Hz, 1H), 5.29 (s, 1H), 5.09–5.00 (m, 4H), 4.92 (dd, J =12.4, 3.2 Hz, 1H), 4.25 (t, J =6.4 Hz, 1H), 4.08–4.03 (m, 2H), 3.95–3.91 (m, 1H), 2.36–2.34 (m, 2H), 2.10 (s, 3H), 1.97 (s, 3H), 1.92 (s, 3H), 1.88 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 205.9, 170.2 ($\times 2$), 170.0, 135.3, 128.7 ($\times 2$), 128.6, 128.0, 97.8, 92.9, 69.6, 69.5 ($\times 2$), 68.9, 66.7, 66.1, 65.9, 61.7, 40.8, 39.7, 34.4, 30.3, 29.8, 20.6, 20.4; ³¹P NMR (162 MHz, CDCl₃) δ –2.24; MS (m/z)=

607.3 (M+1)⁺; HRMS calcd for C₂₉H₃₅O₁₂P⁺ [M+H]⁺ 607.1939, found 607.1935.

4.1.9. Uridine 5'-diphospho-2-acetyl-2-deoxy- α -D-glucopyranose diammonium salt (8-Glc). Pd/C (10%, 30 mg) was added to a solution of **6-Glc** (290 mg, 0.48 mmol) and triethylamine (101 mg, 0.29 mmol) in dry CH₃OH (4 mL). The resulting reaction mixture was subjected to an H₂ atmosphere via a balloon and allowed to stir for 20 h. TLC (1:1 Hex/EtOAc) showed complete disappearance of **6-Glc** starting material. The reaction mixture was filtered through Celite™ and concentrated to dryness. The residue was taken up in dry pyridine (3 mL) and UMP-morpholidate 4-morpholine-*N,N'*-dicyclohexylcarboxamide salt (492 mg, 0.72 mmol) was added along with activated 4 Å molecular sieves (~300 mg), followed by the addition of 1H-tetrazole (134 mg, 1.91 mmol) in pyridine (2 mL). The resulting reaction mixture was allowed to stir at room temperature for 3 days and then filtered through Celite™. The filtrate was concentrated under reduced pressure and the residue taken up in CH₃OH (4 mL). H₂O (2 mL) was added, followed by Et₃N (1 mL). The resulting reaction mixture was allowed to stir at room temperature for 24 h then diluted with H₂O and extracted with CH₂Cl₂ (two times). The aqueous layer was lyophilized and the crude mixture purified by HPLC using a preparative Agilent Zorbax XDB C-18 column, 21.4×250 mm, 5 μ m, eluted with 100 mM ammonium bicarbonate buffer (NH₄HCO₃, pH=7.9) at a rate of 20 mL/min. The desired product eluted at a retention time of 4.1–6.2 min to afford **8-Glc** (150 mg, 49% yield) as a white powder after lyophilization. ¹H NMR (400 MHz, D₂O) δ 7.93 (d, J =8.0 Hz, 1H), 6.00–5.95 (m, 2H), 5.57 (dd, J =7.2, 3.2 Hz, 1H), 4.40–4.34 (m, 2H), 4.28–4.26 (m, 1H), 4.24–4.16 (m, 2H), 3.91–3.78 (m, 3H), 3.63 (t, J =9.2 Hz, 1H), 3.46 (t, J =9.2 Hz, 1H), 2.86 (dd, J =18.0, 5.2 Hz, 1H), 2.79–2.72 (m, 2H), 2.62 (s, 3H); ¹³C NMR (100 MHz, D₂O) δ 215.5, 167.2, 151.0, 141.9, 103.5, 96.2, 88.5, 84.6, 74.3, 72.1, 69.6, 68.2, 65.2, 63.9, 63.2, 44.5, 40.6, 30.1; ³¹P NMR (162 MHz, D₂O) δ –11.24 (d, J =18.0 Hz), –12.33 (d, J =19.2 Hz); MS (m/z)=605.0 (M+1)⁺; HRMS (ESI) calcd for C₁₈H₂₆N₂O₁₇P₂⁺ [M–H][–] 605.0785, found 605.0801.

4.1.10. Uridine 5'-diphospho-2-acetyl-2-deoxy- α -D-galactopyranose diammonium salt (8-Gal). **6-Gal** (60 mg, 99.0 μ mol), triethylamine (21 mg, 59.4 μ mol), 10% Pd/C (8 mg), and CH₃OH (3 mL). Then UMP-morpholidate 4-morpholine-*N,N'*-dicyclohexylcarboxamide salt (102 mg, 0.15 mmol), 1H-tetrazole (28 mg, 0.40 mmol), pyridine (3 mL), activated 4 Å molecular sieves (~300 mg). Then CH₃OH (2 mL), H₂O (1 mL), Et₃N (0.5 mL). Preparative HPLC conditions same as for **8-Glc**, afforded **8-Gal** as white powder (34 mg, 53% yield) after lyophilization. ¹H NMR (400 MHz, D₂O) δ 7.95 (d, J =8.1 Hz, 1H), 5.97–5.95 (m, 2H), 5.57 (dd, J =7.5, 3.4 Hz, 1H), 4.35–4.30 (m, 2H), 4.25–4.23 (m, 1H), 4.21–4.16 (m, 2H), 4.11 (t, J =5.3 Hz, 1H), 3.91–3.88 (m, 1H), 3.78–3.70 (m, 3H), 2.80–2.75 (m, 1H), 2.56 (m, 1H), 2.60 (s, 3H); ¹³C NMR (100 MHz, D₂O) δ 214.5, 167.9, 152.0, 142.1, 104.1, 96.9, 88.4, 84.1, 74.3, 72.1, 69.8, 68.3, 65.2, 64.0, 63.3, 44.6, 40.1, 30.5; ³¹P NMR (162 MHz, D₂O) δ –10.24 (d, J =19.0 Hz), –12.43 (d, J =19.9 Hz); MS (m/z)=605.4 (M+1)⁺; HRMS (ESI) calcd for C₁₈H₂₆N₂O₁₇P₂⁺ [M–H][–] 605.0785, found 605.0798.

Acknowledgements

This work was supported by the NIH Roadmap for Medical Research Initiative through its establishment of the Imaging Probe Development Center, administered by the National Heart, Lung, and Blood Institute.

Supplementary data

Supplementary data associated with this article can be found in online version at doi:10.1016/j.tet.2011.01.081. These data include

MOL files and InChIKeys of the most important compounds described in this article.

References and notes

1. Ramakrishnan, B.; Qasba, P. K. *J. Biol. Chem.* **2002**, *277*, 20833–20839.
2. Khidekel, N.; Arndt, S.; Lamarre-Vincent, N.; Lippert, A.; Poulin-Kerstien, K. G.; Ramakrishnan, B.; Qasba, P. K.; Hsieh-Wilson, L. C. *J. Am. Chem. Soc.* **2003**, *125*, 16162–16163.
3. Boeggeman, E.; Ramakrishnan, B.; Kilgore, C.; Khidekel, N.; Hsieh-Wilson, L. C.; Simpson, J. T.; Qasba, P. K. *Bioconjugate Chem.* **2007**, *18*, 806–814.
4. Ramakrishnan, B.; Boeggeman, E.; Qasba, P. K. *Bioconjugate Chem.* **2007**, *18*, 1912–1918.
5. Pradman, K. Q.; Elizabeth, B.; Boopathy, R. *Biotechnol. Prog.* **2008**, *24*, 520–526.
6. Qasba, P. K.; Ramakrishnan, B.; Boeggeman, E. *Am. Ass. Pharm. Soc. J.* **2006**, *8*, 190–195.
7. Ramakrishnan, B.; Boeggeman, E.; Qasba, P. K. *Expert Opin. Drug Delivery* **2008**, *5*, 149–153.
8. Hang, H. C.; Bertozzi, C. R. *J. Am. Chem. Soc.* **2001**, *123*, 1242–1243.
9. Keck, G. E.; Enholm, E. J.; Yates, J. B.; Wiley, M. R. *Tetrahedron* **1985**, *41*, 4079–4094.
10. Burns, N. Z.; Baran, P. S.; Hoffmann, R. W. *Angew. Chem., Int. Ed.* **2009**, *48*, 2–16.
11. Campbell, R. E.; Tanner, M. E. *J. Org. Chem.* **1999**, *64*, 9487–9492.