

Prediction of Aquaporin Function by Integrating Evolutionary and Functional Analyses

Juliana Perez Di Giorgio · Gabriela Soto · Karina Alleva · Cintia Jozefkowicz ·
Gabriela Amodeo · Jorge Prometeo Muschietti · Nicolás Daniel Ayub

Received: 10 September 2013 / Accepted: 9 November 2013 / Published online: 29 November 2013
© Springer Science+Business Media New York 2013

Abstract Aquaporins (AQPs) are a family of channel proteins, which transport water and/or small solutes across cell membranes. AQPs are present in Bacteria, Eukarya, and Archaea. The classical AQP evolution paradigm explains the inconsistent phylogenetic trees by multiple transfer events and emphasizes that the assignment of orthologous AQPs is not possible, making it difficult to integrate functional information. Recently, a novel phylogenetic framework of eukaryotic AQP evolution showed congruence between eukaryotic AQPs and organismal trees

identifying 32 orthologous clusters in plants and animals (Soto et al. Gene 503:165–176, 2012). In this article, we discuss in depth the methodological strength, the ability to predict functionality and the AQP community perception about the different paradigms of AQP evolution. Moreover, we show an updated review of AQPs transport functions in association with phylogenetic analyses. Finally, we discuss the possible effect of AQP data integration in the understanding of water and solute transport in eukaryotic cells.

Keywords Aquaporin · Evolution · Function · Integration

Juliana Perez Di Giorgio and Gabriela Soto have contributed equally to this work.

J. Perez Di Giorgio · G. Soto · J. P. Muschietti
Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Dr. Héctor Torres (INGEBI-CONICET), Vuelta de Obligado 2490, C1428ADN Buenos Aires, Argentina

G. Soto · N. D. Ayub (✉)
Instituto de Genética Ewald A. Favret (CICVyA-INTA), De los reseros S/N, Castelar, CC25 (1712) Buenos Aires, Argentina
e-mail: nayub@cnia.inta.gov.ar

K. Alleva · C. Jozefkowicz · G. Amodeo
Instituto de Biodiversidad y Biología Experimental y Aplicada (IBBEA, CONICET-UBA), Intendente Güiraldes 2160, Ciudad Universitaria, Pabellón II, C1428EGA Buenos Aires, Argentina

K. Alleva · C. Jozefkowicz · G. Amodeo · J. P. Muschietti (✉)
Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Intendente Güiraldes 2160, Ciudad Universitaria, Pabellón II, C1428EGA Buenos Aires, Argentina
e-mail: prometeo@dna.uba.ar

N. D. Ayub
Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Avda. Rivadavia 1917, C1033AAJ Buenos Aires, Argentina

General Background of Aquaporins

Osmolarity and the corresponding water movement represent one of the crucial environmental factors that lead to cellular homeostasis. For many years, water was considered to enter and leave the cells exclusively through the lipid membrane. It was proposed later, the existence of *hydrophilic pores* that would facilitate water and ion transport through the membranes (Stein and Danielli 1956). Those pores were later named aquaporin (AQP) (Preston et al. 1992). Since then more than 7,000 articles have been indexed under the tag “aquaporin” in PubMed resulted in a detailed picture of what a water channel is. AQPs are not only specific water channels but also solute transporters; i.e., most AQPs transport other molecules such as glycerol, urea, and arsenic, among others.

AQPs are transmembrane channels. Thus, the ability of a molecule to cross an AQP channel depends on its own characteristics (size, polarity, charge) and on the features of the AQP involved. As other channels relevant in physiology, AQPs are passive transporters. Water crosses an AQP

channel or a lipid membrane, and in both cases the driven force for the movement of water molecules is the osmotic gradient. However, the kinetics of water transport depends on the permeability characteristics of each pathway, generally being AQPs more permeable than the lipid bilayer, which implies that AQPs allow a passage of larger flow of water in shorter time intervals.

The Perception of AQPs as a Protein Family

Since the discovery of mammalian AQPs in 1992 and until now much effort has been made to understand the structure and function of AQPs. However, the idea of AQPs as a protein family was formally introduced with a phylogenetic analysis of putative proteins that can transport water and solutes in bacteria and eukaryotes, hypothetically evolutionarily related to previously characterized AQPs (Zardoya and Villalba 2001). In this work, it was showed for the first time a powerful bioinformatic tool, where a putative functional domain was used to automatically identify new AQPs. This superfamily of integral membrane channel proteins was called major intrinsic protein (MIP). Although this work constituted a fundamental step in the understanding of the structure and function of AQPs, it also introduced confusion regarding the definition of AQPs as a family of proteins. This probably occurred because the MIP superfamily is a structural and a functional family with no clear evidence of being a family of homologous genes. As will be discussed later, a phylogenetic analysis using several proteins and showing congruence (same topology) between gene and species trees is a strong evidence for the presence of an evolutionary family; but this type of congruence was not reported for the AQP family. The AQP family could be constituted by non-homologous proteins with similar structure and motifs due to evolutionary convergence.

The lack of precision described above continues to exist. Here, we discuss the consequences of assuming, when reconstructing AQPs' history, that AQPs are a family of monophyletic proteins affecting the classification of AQPs and prediction of their function.

Structure and Motifs in AQPs

As previously mentioned, the AQP denomination is linked to the MIP definition. But what is the MIP superfamily? MIP is considered a conserved domain (cd00333) by The National Center for Biotechnology Information (Marchler-Bauer et al. 2013). According to this definition, the MIP superfamily is divided into two families based on the distinct primary sequences: AQPs and glycerol uptake facilitators (GlpFs) (Zardoya and Villalba 2001). However, there is no

clear division between families in terms of functionality; therefore, MIP and AQPs are considered synonymous and are used interchangeably. MIPs have six transmembrane helices (TMH1–6) and two additional membrane-embedded domains (Maurel et al. 1993) (Fig. 1). The monomeric MIP units contain functional pores that can be stably assembled as tetramers (Verbavatz et al. 1993) (Fig. 2). Nevertheless, it was demonstrated that each monomer in the tetramer is a functional unit (Preston et al. 1992).

The sequences of the amino- and carboxy-terminal halves of MIP genes are similar to each other and are arranged as tandem repeats, apparently originated from the duplication of a half-sized gene (Quigley et al. 2002; Zardoya and Villalba 2001). Each half of the molecule bears one hydrophobic loop which includes two highly conserved Asn–Pro–Ala (NPA) motifs (Fig. 1) involved in the primary selectivity of these transporters (Johanson et al. 2001; Zardoya and Villalba 2001). NPA motifs play a critical function in charge and size obstacle (Wallace and Roberts 2004). After the first NPA motif, there is a second motif known as aromatic/arginine (ar/R) constriction (for comprehensive review see Wu and Beitz 2007). This motif is composed of four residues, two from the helices 2 (H2) and 5 (H5) and two from loop E (LE1 and the invariant R; Fig. 1). This filter seems to be the narrow part of the pore, involved in the rejection of large molecules (Fu 2000; Gomes et al. 2009; Sui et al. 2001; Wallace and Roberts 2004). It has been proposed that the selectivity of the ar/R constriction region is related to proton repulsion and to the binding, through hydrogen bonds, to uncharged molecules such as water and glycerol (Wallace and Roberts 2004). One further step forward in the selectivity, i.e., the discrimination between molecules such as water and glycerol, seems to be given by the P1–P5 motif, which is composed of five amino acid residues located in extracellular loop regions and integral membrane domains of AQPs (Fig. 1). Finally, the AEF motif (Ala–Glu–Phe) is located in the TMH1 of AQPs (Fig. 1) (Zardoya and Villalba 2001). Although this motif is conserved in AQP proteins, its function is still unknown. Regarding AQPs gating and/or regulation, some AQP, are regulated by protonation (Tournaire-Roux et al. 2003). It has been described that protonation (but also, phosphorylation and cation binding) directly affects protein conformation, modifying their transport activity. For the plant AQP SoPIP2;1, the mechanism of a transition from an open to a closed state involves the protonation of conserved histidine residues that moves the LD loop to a position that blocks the water pore (Hedfalk et al. 2006). However, the gating mechanisms for other AQPs are not yet clear. It has been described another putative pH sensing motif $-\mu H\phi\phi\phi$ ($-\phi$: acidic residue, μ : hydrophobic residue, H: histidine, ϕ : polar non-charged residue) that is located in the




Fig. 1 Schematic representation of the classical structure of AQPs. An AQP monomer showing the six transmembrane helices (TMH1–6) connected by two intracellular (LB and LD) and three extracellular (LA, LC and LE) loops. The conserved residues from the first

selective filter (NPA motifs) and the second filter (ar/R constriction) are shown in red and blue, respectively. The P1–P5 residue positions and the putative pH sensing motif— $\mu\text{H}\phi\phi\phi$ are shown in red and gray, respectively (Color figure online)

intracellular LD loop of many plant AQPs and in the extracellular LC loop of *Arabidopsis thaliana* TIP5;1 (Soto et al. 2010). However, the functional validation of this motif in the external and internal sense of pH changes remains to be explored in more detail. AQPs can also form homo- or hetero-oligomers (Jozefkowicz et al. 2013; Neely et al. 1999; Zelazny et al. 2007), but the conditions and/or the motifs involved are not completely understood. Despite the identification of new specific AQP motifs (e.g. AEF, ar/R, and P1–P5– $\mu\text{H}\phi\phi\phi$) described through the last decade, the NPA residues and the six TMHs are still the ones used to identify new classes of putative AQPs.

Classical Clustering of AQPs

AQPs are present in the three domains of life: Bacteria, Eukarya, and Archaea. In Eukarya, the greatest AQP family

diversification occurred in vertebrates and plants. While the classification of animal AQPs (AQP0–12) is broadly consistent and reflects their evolutionary relationships, for plant AQPs is different. In plants, AQP subfamilies were initially named and organized considering their putative subcellular localization, but different subcellular localization within each of the subfamilies has already been reported (for comprehensive reviews see Bienert and Chaumont 2013; Hachez et al. 2013; Ishibashi et al. 2011; Wudick et al. 2009). Currently, plant AQPs are classified into seven subfamilies: plasma membrane intrinsic proteins (PIPs), tonoplast intrinsic proteins (TIPs), NOD26-like intrinsic proteins (NIPs), small basic intrinsic proteins (SIPs), x intrinsic proteins (XIPs), hybrid intrinsic proteins (HIPs), and GlpF-like intrinsic proteins (GIPs) (Danielson and Johanson 2008; Johanson et al. 2001). However, as previously described for the bacterial and eukaryotic AQPs, there is no work supporting the evolutionary integrity of the seven plant AQP subfamilies.

Fig. 2 Cartoon representation of the crystal structure of the spinach aquaporin SoPIP2;1 (Hedfalk et al. 2006) in an open conformation to 3.9 Å resolution (2B5F.PDB from www.pdb.org). Side view (on top) and end on view from the extracellular surface of the tetramer (bottom), each monomer is indicated in a different color (chain A in yellow, chain B in green, chain C in blue and chain D in red). MIP tridimensional structure performed by the Cn3D macromolecular structure viewer software (<http://www.ncbi.nlm.nih.gov/Structure/CN3D/cn3d.shtml>). The MIP structure consists of six trans-membrane helical protein segments lying parallel to the membrane plane (left). A view showing the MIP pore oriented nearly perpendicular to the bilayer plane (right) (Color figure online)




Figure 3 shows the randomness of plant AQPs classification at different levels such as individual proteins and groups within subfamilies. After the release of new genomic data, classification of novel AQPs was done using different criteria. In some cases, new AQPs were named according to the order of availability of the sequence in the database, or to the amino acid identity regarding the AQPs of *A. thaliana* as taxonomic criteria, while new phylogenetic trees containing reference proteins of *Arabidopsis* and/or other plant species were built.

The Hypothetical Lateral Transfer of AQPs Between Bacteria and Eukarya

In the recent years, several reports have suggested that all proteins with a MIP functional domain from Bacteria and Eukarya are actually homologous despite their very low amino acid identity (e.g. <5 %). Under this assumption, AQPs from extremely distant taxa (e.g., Bacteria and Eukarya domains) have been included in the same phylogenetic tree (Fig. 4), obtaining incongruence (i.e. different




Fig. 3 Nomenclature of plant aquaporins under Johansson et al. framework (2001) and the alternative view of this topology. Both possibilities are arbitrary

topology) between AQPs and organismal trees (Danielson 2010; Danielson and Johanson 2008; Heymann and Engel 1999; Johanson et al. 2001; Quigley et al. 2002; Zardoya 2005; Zardoya et al. 2002; Zardoya and Villalba 2001). It has been suggested that these inconsistent patterns are due to the existence of multiple lateral transfer events, such as genetic exchange of AQP genes between unrelated organisms, such as plants, vertebrates, and bacteria. For example, it has been proposed that the animal AQP3 subfamily and

the plant NIP subfamily clusters were acquired by lateral transfer, hypothetically derived from the glycerol facilitator (EcGlpF) from *Escherichia coli* and bacterial NIP-like proteins, respectively (Danielson 2010; Park and Saier 1996; Zardoya et al. 2002). Nevertheless, an unexpected position of a protein within a phylogenetic tree may also be explained by gene duplication, lineage-specific gene loss events, and large amino acid distances (Andersson 2005; Delsuc et al. 2005; Koonin 2003).

The relationships described between AQPs from Bacteria and Eukarya are supported by statistics (high bootstrap values; Danielson 2010; Park and Saier 1996); this information was used to support the lateral transfer hypothesis. However, it is important to point out that obtaining a strongly supported tree does not necessarily indicate that the tree is correct (Delsuc et al. 2005). It is possible to obtain an inaccurate, but statistically supported, phylogenetic tree if the method used does not correctly handle the properties of the data (Delsuc et al. 2005). To avoid overestimation of gene transfer, general congruence with the organismal tree, except for the transfer event, must be observed (Phillips 2006); it is also necessary to find an independent evidence such as

localization within genomic islands (Ayub et al. 2007; Yan et al. 2008), or to take advantage of powerful algorithms specifically developed to statistically support gene transfer events (Abby et al. 2010).

The comparative study of genomes from Bacteria and Eukarya indicates that a major fraction of the genes in the prokaryotic genomes have been acquired by horizontal transfer (Koonin et al. 2001). The quantity of horizontal transfer of genes is often associated with the microorganism lifestyle. In addition, the transfer of genes in eukaryotic cells is commonly associated with symbiotic or parasitic relationships with bacteria (Novichkov et al. 2004). Fixation and long-term persistence of horizontally transferred genes




Fig. 4 Schematic representation of evolutionary relationships between eukaryotic and bacterial MIPs based on previously phylogenetic analyses (Zardoya and Villalba 2001)




Fig. 5 Schematic representation of homologous, orthologous, and paralogous genes

imply that these genes present a selective advantage on the recipient organism (Phillips 2006). Therefore, works proposing the acquisition of bacterial genes in the eukaryotic host should explore the evolutionary advantage of this hypothetical transfer event. This exploration should consist of empirical evidence predicting functionality.

In conclusion, lateral transfer of AQPs between Bacteria and Eukarya could be an artifact. Thus, the inclusion of these distant or unrelated proteins within the same phylogenetic tree (e.g. used as an out-group) can produce a negative impact on the reconstruction of AQP evolution.

Essential Concepts for a Critical View of Evolutionary Studies of AQPs

The most rigorous way to represent the evolutionary history is through the construction of phylogenetic trees. The

main assumption is that the genes analyzed within the tree are homologous genes (Fig. 5). On the other hand, because it is difficult to know which genes are actually homologous, some type of approximation to preselect homologous genes to be included in a phylogenetic tree is needed. The most common approach is to constrain the phylogenetic analysis to proteins that, as a whole, have more than 25 % of amino acid identity (Hughes et al. 2005). But, no sampling criterion (for example >25 % of amino acid identity) is sufficient to ensure that two genes are homologous. It is possible that two genes with high amino acid identity (e.g. 50 %) are non-homologous (by convergent evolution) or genes with low amino acid identity (e.g. 30 %) are homologous (by functional divergence). This type of criteria (e.g. amino acid identity) is used because there is no statistical elements to designate homology, and this occurs because systematics is a historical science with particular epistemological limitations (Cleland 2002). For example,

the information used for the evolutionary reconstruction arises from the observation, and not from the design and execution of experiments. This means that, in many cases, the results of phylogenetic analysis are accepted by the scientific community but with certain precautions.

The divergence of orthologous genes coincides with and is a product of the divergence of the species in which they are included (Fig. 5). Naturally, orthologous genes are evolutionarily more closely related and are therefore expected to have a similar biological and biochemical function. Robust methods for finding orthologs are based on the analysis of phylogenetic trees. For orthologous assignment, the trees have to be congruent (same topology) with the species tree (Fig. 5).

Different phylogenetic reconstruction methods, such as maximum parsimony, minimum evolution, and neighbor-joining, are available within freely accessible evolutionary software packages such as MEGA (Tamura et al. 2011). These methods compare the nucleotide or amino acid sequences using different parameters of genetic distance and parsimony but, in all cases, independent of the reconstruction method used, the number of possible phylogenetic trees increases exponentially with the number of sequences (Li 1997). Since a congruent pattern is only one topology, the probability that a congruent pattern occurs by chance is practically null. For example, when a phylogenetic tree has only 20 sequences, the probability to obtain a congruent pattern by chance (congruent pattern/possible trees) is $1/2 \times 10^{21} \approx 0$ (Fig. 6).

As described previously works on AQP evolution contained numerous inaccuracies in methods and/or


interpretation of results, but this does not imply that by correcting these problems would be possible to obtain a consistent phylogeny, especially considering that AQPs constitute a broadly diversified family of genes.

The Novel Paradigm of AQP Evolution: Vertical Transfer

In this complex background, the phylogeny of eukaryotic AQPs has been recently re-evaluated by restricting the analysis to proteins with high amino acid identity (>25 %) and using sequences from well-characterized species of flowering plants and vertebrates (Soto et al. 2012). Since members of the subfamilies PIPs, TIPs, NIPs, and SIPs from flowering plants and AQPs plus aquaglyceroporins from vertebrates met the requirement of amino acid identity (>25 %), were included in the analysis. In contrast, the subfamilies XIPs, GIPs, and HIPs were not analyzed because they did not meet that requirement. As previously explained, this result does not imply that the XIPs, GIPs, and HIPs subfamilies are not homologous to the rest of eukaryotic AQPs, but shows that it was not possible to analyze their evolution according to the strict criterion of selection of proteins included within a same phylogenetic tree.

This strict criterion has shown congruence between AQPs (210 proteins) and organismal (13 species of eukaryotes) trees (Soto et al. 2012). The probability of finding a congruent pattern (vertical transfer) by chance was practically null. The advantage of this new perspective is that its congruence allowed defining clusters of

Fig. 6 Exponential function describing the relationship between number of possible unrooted phylogenetic trees and number of genes or proteins analyzed






Fig. 7 Schematic representation of orthologous gene clusters in plants based on Soto et al. (2012)




Fig. 8 Schematic representation of orthologous gene clusters in animals according to Soto et al. (2012)

orthologous genes for both flowering plants (19 clusters; Fig. 7) and vertebrates (13 clusters; Fig. 8). This leads to the description of specific conserved motifs for each orthologous cluster as a useful tool for automatic assignment of orthologs (Fig. 9) (Soto et al. 2012). The identification of conserved motifs in each subfamily and in each cluster of orthologous genes offers a framework for studying the possible functional implication of such motifs. This is a powerful tool, and the availability of new genomes of flowering plants and vertebrates could serve to define these motifs more precisely.

This paradigm offers the opportunity to establish a new classification of eukaryotic AQPs based on their evolutionary relationships. In this way, the current nomenclature differs significantly from a nomenclature based on the identification of orthologous genes. Although the nomenclature is important per se, it also has significant influence in the functional field. For example, given the current nomenclature, AtPIP2;6 of *Arabidopsis* can be considered the equivalent protein (ortholog) of OsPIP2;6 of *Oryza sativa*, suggesting an equivalent role in both plants. However, AtPIP2;7 and AtPIP2;8 are equally related to OsPIP2;6 expecting similar functions for all these proteins (Soto et al. 2012). Another advantage of having clusters of orthologous genes is the possibility of evaluating evolutionary constraints. The fact that PIPs have greater evolutionary constraints than TIPs, NIPs, and SIPs, support the prediction of greater functional constraints for PIPs (Soto et al. 2012). In turn, some of the TIPs and NIPs orthologous gene clusters also showed high evolutionary constraints, suggesting functional constraints.

This AQP phylogenetic framework for flowering plants and vertebrates can be used to predict a putative function of individual AQPs on the basis of orthologous genes from *A. thaliana* and *Homo sapiens*. However, the separate identification of clusters of orthologous genes in plants and vertebrates does not allow extrapolating the function between AQPs belonging to organisms of both kingdoms. As previously suggested by other phylogenetic analyses (Cerda and Finn 2010; Finn and Cerda 2011; Tingaud-Sequeira et al. 2010), our phylogenetic framework revealed that each subfamily of plant AQPs was related to a subfamily of animal AQP, thus showing a pattern of vertical transfer which predicts the presence of at least four families of AQPs in the ancestral eukaryote from which plants and vertebrates derived (Soto et al. 2012). We suggest that the four AQPs subfamilies described in animals (AQP1-, AQP8-, AQP3-, and AQP11-like) and plants (PIP-, TIP-, NIP-, and SIP-like) are derived from four ancestral AQPs subfamilies: A–D, respectively (Fig. 10). Thus, a pattern of vertical transfer in the evolution of AQPs of animals and plants at all levels, i.e., within (Figs. 7, 8) and between kingdoms (Fig. 10), was observed.

PIP-like		
subfamily TGINPARS [FIL]		
CL1	MEGK	
CL2	M[AG]KX	
CL3	MSKE	
TIP-like		
subfamily [AG] [AGS] MNPA [CRSV] [ASV] F		
CL1	EFISTLIFVXAGX.....	NILAGGAFXGASMNPAXXF
CL2	EFXSMXIFVFAGX.....	NILXGGAFDGASMNPAVSF
CL3	EFIATLIFVFAGV.....	NILAAGPFSGGSMNPARSF
CL4	EXXXTXXFVFAXE.....	NXLXGGPFXGAXMNPARXF
CL5	EFXSTLXVFVFAGV.....	NILXAGPFSGGSMNPARSF
CL6	EXXXTFLFVFXGV.....	NXKACXXXXGASMNPARSF
CL7	EFXSTFXXVXXXX.....	XVLAAGXXXXGXSMNPAXXF
NIP-like		
subfamily [AG] S [LM] NP [AGV] R [ST] [ILV]		
CL1	EXXGTY.....	NVXXAXXXXASMNPXRXX
CL2		NVFVAGPXSGASMNPARSX
CL3	EXXGTF.....	XSIIXAGXXSGGSMMNPARTL
CL4		NILXXGPXXGXSMNPVRXL
CL5		NIXIAAGXXTXASMNPVRXL
CL6		XXLXXGXXXGXSMNPARXL
SIP-like		
subfamily [LM] NPAXXXWA		
CL1	[FY]NP[TC]	
CL2	[FY]NP[AS]	
CL3	[FY]NPL	
Blue are hydrophobic residues: ACFILVWM		
Fuchsia are large hydrophobic aminoacids: FIWLM		
Green are polar aminoacids: NQST		
Pink are negative residues: DE		
Red are positive residues: KRHGPY		

Fig. 9 Motif characterization of plant aquaporins. Illustration of amino acid motifs for each subfamily (PIPs, TIPs, NIPs, and SIPs) and each orthologous cluster (CL). The letter X represents similar amino acids. Blue, fuchsia, green, pink, and red letters represents hydrophobic (ACFILVWM), large hydrophobic (FIWLM), polar (NQST), negative (DE) and positive (KRHGPY) amino acids, respectively (Color figure online)




Fig. 10 Evolution of MIP superfamily in plants and animals. The hypothetical ancestral eukaryote has four AQP subfamilies (AD)

Functional Transfer of Eukaryotic AQPs Under the Novel Evolution Paradigm

This new framework allowed the comparison for each individual protein of the evolutionary patterns together with the described functions. We previously showed a correlation between the phylogenetic analysis and the functional information of 47 AQPs (Soto et al. 2012). Tables 1 and 2 describe the evolutionary location (subfamilies A–D), their current nomenclature, and molecules that they transport, the organisms that contain them for 106 AQPs that showed strong correlation between evolutionary and functional data.

AQPs of the subfamily A (PIP- and AQP1-like) transport water and also CO_2 as a common ancestral feature. Afterward, some AQPs of this subfamily acquired the capacity to transport other solutes, like glycerol, urea, and hydrogen peroxide. Some AQPs show poor water transport, however, they are permeable to anions as is the case of AQP6 (Table 2). Among PIPs, PIP2s transport water, whereas for PIP1s there is a great functional divergence: some reports show that PIP1s do not form functional homotetramers in plasma membrane (Fetter et al. 2004; Zelazny et al. 2007), or that it transports CO_2 , or that it has low water transport capacity (Table 1). Several reports showed that PIP1 is localized in the plasma membrane only when it is expressed together with PIP2 (Bellati et al. 2010; Fetter et al. 2004); if it is expressed alone, then it is retained in the endoplasmic reticulum (Fetter et al. 2004; Jozefkowicz et al. 2013; Zelazny et al. 2007). In this context, the possibility of PIP1–2 hetero-oligomerization became a new way of regulation of water transport.

AQPs of the subfamily B (TIP- and AQP8-like) transport water and urea. However, many reports have demonstrated that several *Arabidopsis* TIPs transport not only urea but also ammonia (Table 1). In particular, as it is discussed later, *AtTIP5;1* is an urea transporter that belongs to the most divergent cluster of *Arabidopsis* TIPs (Soto et al. 2010), suggesting that this function could be ancestral. Furthermore, the ability to transport hydrogen peroxide could also be an ancestral feature, as several members of this subfamily share this capacity. Interestingly, new features appeared after functional divergence because some AQPs of this subfamily transport glycerol (Table 1).

AQPs of the subfamily C (NIP- and AQP3-like) are named aquaglyceroporins because almost all members of this group transport water and glycerol (Tables 1, 2). The transport of metalloids may be an ancestral feature as many plant NIPs and also AQP7 and -9 transport arsenite. AQP10 was found to be expressed in the small intestine

and transport water, glycerol, and urea (Table 2). The fact that urea and boric acid transport was observed only in the animal and plant members of this subfamily, respectively, suggest that the transport of glycerol and urea might not be an ancestral feature but a putative event of functional divergence. Additionally, it was reported that many NIP-like AQPs transport other compounds, such as formamide and lactic acid (Table 1).

AQPs of the subfamily D (SIP- and AQP11-like) include the most recently identified AQPs. They are unusual because only the second NPA motif located in LE is conserved while the NPA motif in LB loop is modified, such as NPC in AQP11 and AtSIP1;2 or NPT in AQP12 and AtSIP1;1 (Soto et al. 2010). Also, their N-terminal tail is shorter than the N-terminal of the rest of the AQPs, a characteristic that has been assigned to explain their intracellular localization (Maeshima and Ishikawa 2008). Due to this, it has been difficult to establish their transport substrate specificity when expressed in *Xenopus* oocytes.

In summary, although AQPs from all subfamilies show different pattern of solute transport, all AQPs are permeable to water and so, this can be considered the ancestral feature shared by all the four subfamilies.

The potential of the proposed evolutionary framework in the prediction of functionality of plants and animals AQPs can be illustrated, with the example of the animal AQP8 and plant TIPs both members of the subfamily B (Fig. 7). Within TIPs, the cluster 7, includes *AtTIP5;1* of *A. thaliana*, which is the most divergent TIP and therefore the most similar to the ancestral protein that gave rise to all TIPs (Fig. 7). Therefore, the evolutionary framework predicts that AQP8 would have a function equivalent to that of TIP5;1. It was described that TIP5;1 is an urea transporter located in pollen tube mitochondria when overexpressed in the pollen vegetative cell of *A. thaliana* (Soto et al. 2008, 2010). Based on these results, it has been proposed that *AtTIP5;1* would be involved in the efflux of urea from mitochondria during the urea cycle (Soto et al. 2010). The urea cycle is well conserved in all living organisms (Goldraij and Polacco 2000; Kojima et al. 2006; Mobley et al. 1995; Pedrozo et al. 1996; Yu et al. 1997): urea is synthesized by a mitochondrial arginase and degraded by a cytosolic urease. Therefore, a mitochondrial transporter that would export urea from the mitochondrion into the cytosol has been predicted for many years discarding the possibility of a passive transport (Rodela et al. 2008). Due to AQP8 is the putative ortholog of *AtTIP5;1* (Fig. 10), our phylogenetic framework predicts that AQP8 is involved in the urea cycle in vertebrates as it was suggested previously (Calamita et al. 2006, 2007; Holm et al. 2005; Liu et al. 2006; Soria et al. 2013).

Table 1 Functional characteristics of plants AQPs disaggregated by subfamily

	H ₂ O	Gly	NH ₄	Urea	B	As	H ₂ O ₂	CO ₂	I	O	References
SF A											
AtPIP1;1	+					—					Hooijmaijers et al. (2012), Kammerloher et al. (1994)
AtPIP1;2	+					—	+				Heckwolf et al. (2011), Hooijmaijers et al. (2012), Kammerloher et al. (1994), Tournaire-Roux et al. (2003)
AtPIP1;3	+					—					Hooijmaijers et al. (2012), Kammerloher et al. (1994)
AtPIP1;4						—					Hooijmaijers et al. (2012)
AtPIP1;5						—					Hooijmaijers et al. (2012)
NtAQP1	+	+						+	—		Biela et al. (1999), Otto et al. (2010), Uehlein et al. (2003)
ZmPIP1;2							—				Bienert and Chaumont (2013)
ZmPIP1;5	+			+							Gaspar (2003)
SsAQP1	—		+								Moshelion (2002)
AtPIP2;1	+					—					Bienert et al. (2007), Dynowski et al. (2008), Hooijmaijers et al. (2012), Kammerloher et al. (1994)
AtPIP2;2	+					—	+				Hooijmaijers et al. (2012), Kammerloher et al. (1994), Tournaire-Roux et al. (2003)
AtPIP2;4	+					—	+				Dynowski et al. (2008), Hooijmaijers et al. (2012)
AtPIP2;3	+					—					Daniels et al. (1994), Hooijmaijers et al. (2012)
AtPIP2;5	+					—	+				Hooijmaijers et al. (2012)
AtPIP2;6						—					Hooijmaijers et al. (2012)
AtPIP2;7						—	+				Hooijmaijers et al. (2012)
AtPIP2;8						—					Hooijmaijers et al. (2012)
ZmPIP2;1	+										Fetter et al. (2004)
ZmPIP2;4											Fetter et al. (2004)
ZmPIP2;5						—					Bienert and Chaumont (2013), Chaumont et al. (2001), Fetter et al. (2004)
McPIP2;1	+	—		—							Amezcu-Romero et al. (2010)
SoPIP2;1	+										Johansson et al. (1998)
SsAQP2	+		—								Moshelion (2002)
OsPIP2;1	+										Matsumoto et al. (2009), Sakurai et al. (2008)
OsPIP2;2	+										Matsumoto et al. (2009), Sakurai et al. (2008)
OsPIP2;3	+										Matsumoto et al. (2009), Mosa et al. (2012), Sakurai et al. (2008)
OsPIP2;4	+				—	+					Matsumoto et al. (2009), Sakurai et al. (2008)
OsPIP2;5	+										Matsumoto et al. (2009), Sakurai et al. (2008)
OsPIP2;6	+				—	+					Matsumoto et al. (2009), Mosa et al. (2012)
OsPIP2;7	+				—	+					Matsumoto et al. (2009), Mosa et al. (2012)
OsPIP2;8	+										Matsumoto et al. (2009)
NtPIP2;1	+					—					Bots et al. (2005), Otto et al. (2010)
SF B											
AtTIP1;1	+	—		+		+	+	+			Bienert et al. (2007), Klebl et al. (2003), Liu et al. (2003), Maurel et al. (1993)
TgTIP1;1	+		—	+	+		—	+			Azad et al. (2008, 2012)
AtTIP1;2					—		—	—			Bienert et al. (2007), Liu et al. (2003)
OsTIP1;2	+	—	—								Li et al. (2008), Sakurai et al. (2008)
TgTIP1;2	+		—	—		—	—	—			Azad et al. (2008, 2012)
AtTIP1;3	+	—		—	—	—	—	—			Soto et al. (2008)
AtTIP2;1	+		—	—	—	—	—	—			Klebl et al. (2003), Liu et al. (2003), Loque et al. (2005), Maurel et al. (1993)
OsTIP2;1	+	—	—	—	—	—	—	—			Li et al. (2008)
TaTIP2;1	+		—	—	—	—	—	—			+
TaTIP2;2	+		—	—	—	—	—	—			Holm et al. (2005), Jahn et al. (2004)
											Bertl and Kaldenhoff (2007)

Table 1 continued

	H ₂ O	Gly	NH ₄	Urea	B	As	H ₂ O ₂	CO ₂	I	O	References
AtTIP2;3			+	+			+				Dynowski et al. (2008), Loque et al. (2005)
AtTIP3;1	+										Eckert et al. (1999)
OsTIP3;2	-	+									Li et al. (2008)
OsTIP4;1	+	+									Li et al. (2008)
AtTIP5;1	+	-		+	-						Soto et al. (2008)
NtTIPa	+	+		+							Gerbeau et al. (1999)
SF C											
AtNIP1;1	+	+				+	-			+	Dynowski et al. (2008), Kamiya and Fujiwara (2009), Kamiya et al. (2009), Weig and Jakob (2000)
AtNIP1;2		+					+				Dynowski et al. (2008), Weig and Jakob (2000)
AtNIP2;1	+	+					+			+	Choi and Roberts (2007), Mizutani et al. (2006)
AtNIP4;1	+										Soto et al. (2008)
AtNIP5;1	+	+				+				+	Bienert et al. (2008), Mitani-Ueno et al. (2011), Takano et al. (2006)
AtNIP6;1	-	+		+	+	+	+			+	Bienert et al. (2008), Tanaka et al. (2008), Wallace and Roberts (2005)
AtNIP7;1	+	+		+	+	+	+			+	Bienert et al. (2008), Li et al. (2011)
CpNIP1				+							Klebl et al. (2003), Liu et al. (2003)
GmNOD26	+	+	+	-	-				-	+	Dean et al. (1999), Hwang et al. (2010), Rivers et al. (1997), Schnurbusch et al. (2010), Wallace et al. (2012)
HvNIP2;1	+			-		+	+			+	Ligaba et al. (2011), Schnurbusch et al. (2010)
LjLIMP2	+	+								+	Guenther and Roberts (2000)
LjNIP5;1						+				+	Bienert et al. (2008)
LjNIP6;1						+				+	Bienert et al. (2008)
OsNIP1;1						+					Ma et al. (2008)
OsNIP2;1	+			+	+	+				+	Mitani-Ueno et al. (2011), Mitani et al. (2008)
OsNIP2;2	+				-	+				+	Bienert et al. (2008), Ma et al. (2006, 2008), Mitani-Ueno et al. (2011), Mitani et al. (2008)
OsNIP3;1						+					Ma et al. (2008)
OsNIP3;2					+					+	Bienert et al. (2008)
PsNIP1;1	+	+									Schuurmans et al. (2003)
PtNIP1;1	+	+									Ciavatta et al. (2001)
TaNIP2;1										+	Montpetit et al. (2012)
ZmNIP2;1			+							+	Gu et al. (2012), Mitani et al. (2009)
ZmNIP2;2										+	Mitani et al. (2009)
ZmNIP2;4			+								Gu et al. (2012)
SF D											
AtSIP1;1	+										Ishikawa et al. (2005)
AtSIP1;2	+										Ishikawa et al. (2005)
AtSIP2;1	-										Ishikawa et al. (2005)

+, presence; -, absence; +/-, controversy between authors. Plants: At, *Arabidopsis thaliana*; Cp, *Cucurbita pepo*; Gm, *Glycine max*; Hv, *Hordeum vulgare*; Lj, *Lotus japonicus*; Mc, *Mesembryanthemum crystallinum*; Nt, *Nicotiana tabacum*; Os, *Oryza sativa*; Ps, *Polygonum sibiricum*; Pt, *Pinus taeda*; So, *Spinacia oleracea*; Ss, *Samanea saman*; Ta, *Triticum aestivum*; Tg, *Tulipa gesneriana*; Zm, *Zea mays*

SF subfamily, Gly glycerol, B boric acid, As arsenic, I ions, O other compounds (e.g., formamide and lactic acid)

Table 2 Functional characteristics of vertebrates AQPs disaggregated by subfamily

	H ₂ O	Gly	NH ₄	Urea	B	As	H ₂ O ₂	CO ₂	I	O	References
SF A											
BtAQP0	+						+				Mulders et al. (1995), Yang and Verkman (1997), Zampighi et al. (1985)
HsAQP0	+							+			Chandy et al. (1997)
DrAQP0	+										Froger et al. (2010)
SaAQP0a	+										Chauvigne et al. (2013)
RnAQP1	+	+		-			+		+	-	Abrami et al. (1995), Li et al. (2011), Ma et al. (1993), Marinelli et al. (1997)
DrAQP1	+	-		-							Tingaud-Sequeira et al. (2010)
HsAQP1	+	+	+					+	+		Abrami et al. (1995), Endeward et al. (2006), Preston et al. (1992), Anthony et al. (2000), Herrera et al. (2006), Musa-Aziz et al. (2009), Nakhoul et al. (1998), Prasad et al. (1998)
SaAQP1aa/ ab	+										Chauvigne et al. (2013)
HsAQP2	+	+	-	-				-			Abrami et al. (1995), Fushimi et al. (1993), Geyer et al. (2013), Meinild (1998), Yang and Verkman (1997)
DrAQP4	+										Tingaud-Sequeira et al. (2010)
RnAQP4	+	+/-	-	-				+			Fenton et al. (2010), Geyer et al. (2013), Jung et al. (1994), Meinild (1998), Musa-Aziz et al. (2009), Yang and Verkman (1997)
RnAQP4	+	-	-	-				-			Fenton et al. (2010), Geyer et al. (2013)
RnAQP5	+										Raina et al. (1995), Yang and Verkman (1997)
HsAQP5	+	-	-					+			Meinild (1998), Musa-Aziz et al. (2009)
HsAQP6	+	+			+				+	+	Holm et al. (2004), Liu et al. (2006), Ma et al. (1996)
RnAQP6	-		+					+	+	+	Geyer et al. (2013), Hazama et al. (2002), Ikeda et al. (2002), Liu et al. (2006), Yasui et al. (1999)
SF B											
HsAQP8	+	-	+	-			+	-	+		Bienert et al. (2007), Geyer et al. (2013), Jahn et al. (2004), Liu et al. (2006)
RnAQP8	+	+/-	+	+/-					+		Holm et al. (2005), Ishibashi et al. (1997), Koyama et al. (1997), Liu et al. (2006)
DrAQP8	+	-		+							Tingaud-Sequeira et al. (2010)
SaAQP8b	+			+							Chauvigne et al. (2013)
SF C											
RnAQP3	+	+	+	+			+	-	-	+	Echevarria et al. (1994), Geyer et al. (2013), Hara-Chikuma et al. (2012), Holm et al. (2005), Ishibashi et al. (1994), Meinild (1998), Yang and Verkman (1997), Zeuthen et al. (1997)
DrAQP3	+	+			+					+	Chauvigne et al. (2011), Tingaud-Sequeira et al. (2010)
HsAQP3	+	+					-			+	Chauvigne et al. (2011), Liu et al. (2004)
HsAQP7	+		+				+		-		Geyer et al. (2013), Liu et al. (2004)
MmAQP7	+				+						Liu et al. (2002)
RnAQP7	+	+		+							Ishibashi et al. (1997), Kishida et al. (2000)
DrAQP7	+	+		+						+	Chauvigne et al. (2011), Tingaud-Sequeira et al. (2010)
SaAQP7	+	+		+							Chauvigne et al. (2013)
RnAQP9	+	+	+	+			+		+	+	Geyer et al. (2013), Liu et al. (2002), Tsukaguchi et al. (1999)
DrAQP9	+	+		+						+	Chauvigne et al. (2011), Tingaud-Sequeira et al. (2010)
HsAQP9							+				Liu et al. (2004), McDermott et al. (2010)
SaAQP9b	+	+		+							Chauvigne et al. (2013)
HsAQP10	+	+		+			-				Hatakeyama et al. (2001), Ishibashi et al. (2002), Liu et al. (2004)
DrAQP10	+	+		+							Tingaud-Sequeira et al. (2010)
SaAQP10b	+	+		+							Chauvigne et al. (2013)

Table 2 continued

	H ₂ O	Gly	NH ₄	Urea	B	As	H ₂ O ₂	CO ₂	I	O	References
SF D											
MmAQP11	+										Yakata et al. (2007, 2011)
HsAQP11	+										Ikeda et al. (2011)

+, presence; -, absence; +/-, controversy between authors. Animals: Ac, *Anomala cuprea*; Bg, *Blattella germanica*; Bt, *Bos taurus*; Ba, *Bemisia tabaci*; Dr, *Danio rerio*; Gg, *Gallus gallus*; Hc, *Hyla chrysoscelis*; Hs, *Homo sapiens*; Mm, *Mus musculus*; Rn, *Rattus norvegicus*; Sa, *Sparus aurata*

SF subfamily, Gly glycerol, B boric acid, As arsenic, I ions, O other compounds (e.g. formamide and lactic acid)

Conclusions and Prospects

The AQP vertical transfer hypothesis makes predictions that are testable and refutable, as we have demonstrated throughout the text so far. This new paradigm of evolution of plant and animal AQPs offers a novel framework to integrate functional information. It allows two distant groups, the plant and animal AQPs, to work together and support each other, especially in understanding water and solute transport. Furthermore, the availability of new clusters of orthologous genes and specific motifs associated with such clusters offers a starting point for an in-depth understanding of the consensuses and tridimensional structures associated with the functional diversity of AQPs in the specificity of transport, interaction among AQPs and with other molecules, including regulation and subcellular localization. In this context, it is expected that the new consistent evolutionary framework of eukaryotic AQPs increases the ability to properly predict biochemical and biological functions of AQPs. Functional information of individual AQPs by empirical studies is expected to grow and more sequenced genomes of plants and animals are expected to be available, positively influencing the definition and precision of motifs and functions of each cluster of orthologous genes and each AQP subfamily. However, it is necessary to point out that the extrapolation of functionality has the intrinsic restriction of the biochemical and biological diversification processes. For example, the extrapolation of the biochemical and biological functions of the ancestor of flowering plants is not always possible because each of the clusters of orthologous genes in monocotyledonous and dicotyledonous plants evolved independently, incorporating and eliminating various functions related to AQPs. Similarly, although certain features of cell functionality may have been conserved in all vertebrates, it is not expected that the AQPs of fishes and mammals have exactly the same biochemical and biological function, especially when they are exposed to different environments that would potentiate their functional divergence. Finally, the future of the experimental study of AQPs seems to have evolutionary guidance, which, despite its limitations constitutes a solid road toward a better understanding of AQPs.

References

Abby SS, Tannier E, Gouy M, Daubin V (2010) Detecting lateral gene transfers by statistical reconciliation of phylogenetic forests. *BMC Bioinform* 11:324

Abrami L, Tacnet F, Ripocer P (1995) Evidence for a glycerol pathway through aquaporin 1 (CHIP28) channels. *Pflüg Arch Eur J Physiol* 430:447–458

Amezcuia-Romero JC, Pantoja O, Vera-Estrella R (2010) Ser123 is essential for the water channel activity of McPIP2;1 from *Mesembryanthemum crystallinum*. *J Biol Chem* 285:16739–16747

Andersson JO (2005) Lateral gene transfer in eukaryotes. *Cell Mol Life Sci* 62:1182–1197

Anthony TL, Brooks HL, Boassa D, Leonov S, Yanochko GM, Regan JW, Yool AJ (2000) Cloned human aquaporin-1 is a cyclic GMP-gated ion channel. *Mol Pharmacol* 57:576–588

Ayub ND, Pettinari MJ, Mendez BS, Lopez NI (2007) The polyhydroxyalkanoate genes of a stress resistant Antarctic *Pseudomonas* are situated within a genomic island. *Plasmid* 58:240–248

Azad AK, Katsuhara M, Sawa Y, Ishikawa T, Shibata H (2008) Characterization of four plasma membrane aquaporins in tulip petals: a putative homolog is regulated by phosphorylation. *Plant Cell Physiol* 49:1196–1208

Azad AK, Yoshikawa N, Ishikawa T, Sawa Y, Shibata H (2012) Substitution of a single amino acid residue in the aromatic/arginine selectivity filter alters the transport profiles of tonoplast aquaporin homologs. *Biochim Biophys Acta* 1818:1–11

Bellati J, Alleva K, Soto G, Vitali V, Jozefkowicz C, Amodeo G (2010) Intracellular pH sensing is altered by plasma membrane PIP aquaporin co-expression. *Plant Mol Biol* 74:105–118

Bertl A, Kaldenhoff R (2007) Function of a separate NH₃-pore in Aquaporin TIP2;2 from wheat. *FEBS Lett* 581:5413–5417

Biela A, Grote K, Otto B, Hoth S, Hedrich R, Kaldenhoff R (1999) The *Nicotiana tabacum* plasma membrane aquaporin NtAQP1 is mercury-insensitive and permeable for glycerol. *Plant J* 18:565–570

Bienert GP, Chaumont F (2013) Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochim Biophys Acta*. doi:10.1016/j.bbagen.2013.09.017

Bienert GP, Moller AL, Kristiansen KA, Schulz A, Moller IM, Schjoerring JK, Jahn TP (2007) Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem* 282:1183–1192

Bienert GP, Thorsen M, Schussler MD, Nilsson HR, Wagner A, Tamas MJ, Jahn TP (2008) A subgroup of plant aquaporins facilitate the bi-directional diffusion of As(OH)₃ and Sb(OH)₃ across membranes. *BMC Biol* 6:26

Bots M, Feron R, Uehlein N, Weterings K, Kaldenhoff R, Mariani T (2005) PIP1 and PIP2 aquaporins are differentially expressed during tobacco anther and stigma development. *J Exp Bot* 56:113–121

Calamita G, Gena P, Meleleo D, Ferri D, Svelto M (2006) Water permeability of rat liver mitochondria: a biophysical study. *Biochim Biophys Acta* 1758:1018–1024

Calamita G, Moreno M, Ferri D, Silvestri E, Roberti P, Schiavo L, Gena P, Svelto M, Goglia F (2007) Triiodothyronine modulates the expression of aquaporin-8 in rat liver mitochondria. *J Endocrinol* 192:111–120

Cerda J, Finn RN (2010) Piscine aquaporins: an overview of recent advances. *J Exp Zool A* 313:623–650

Chandy G, Zampighi GA, Kreman M, Hall JE (1997) Comparison of the water transporting properties of MIP and AQP1. *J Membr Biol* 159:29–39

Chaumont F, Barrieu F, Wojcik E, Chrispeels MJ, Jung R (2001) Aquaporins constitute a large and highly divergent protein family in maize. *Plant Physiol* 125:1206–1215

Chauvigne F, Lubzens E, Cerda J (2011) Design and characterization of genetically engineered zebrafish aquaporin-3 mutants highly permeable to the cryoprotectant ethylene glycol. *BMC Biotechnol* 11:34

Chauvigne F, Boj M, Vilella S, Finn RN, Cerda J (2013) Subcellular localization of selectively permeable aquaporins in the male germ line of a marine teleost reveals spatial redistribution in activated spermatozoa. *Biol Reprod* 89:37

Choi WG, Roberts DM (2007) Arabidopsis NIP2;1, a major intrinsic protein transporter of lactic acid induced by anoxic stress. *J Biol Chem* 282:24209–24218

Ciavatta VT, Morillon R, Pullman GS, Chrispeels MJ, Cairney J (2001) An aquaglyceroporin is abundantly expressed early in the development of the suspensor and the embryo proper of loblolly pine. *Plant Physiol* 127:1556–1567

Cleland CE (2002) Methodological and epistemic differences between historical science and experimental science. *Philos Sci* 69:474–496

Daniels MJ, Mirkov TE, Chrispeels MJ (1994) The plasma membrane of arabidopsis thaliana contains a mercury-insensitive aquaporin that is a homolog of the tonoplast water channel protein TIP. *Plant Physiol* 106:1325–1333

Danielson J (2010) Plant major intrinsic proteins, natural variation and evolution. *Media-Tryck AB, Lund*

Danielson JA, Johanson U (2008) Unexpected complexity of the aquaporin gene family in the moss *Physcomitrella patens*. *BMC Plant Biol* 8:45

Dean RM, Rivers RL, Zeidel ML, Roberts DM (1999) Purification and functional reconstitution of soybean nodulin 26. An aquaporin with water and glycerol transport properties. *Biochemistry* 38:347–353

Delsuc F, Brinkmann H, Philippe H (2005) Phylogenomics and the reconstruction of the tree of life. *Nat Rev Genet* 6:361–375

Dynowski M, Schaaf G, Loque D, Moran O, Ludewig U (2008) Plant plasma membrane water channels conduct the signalling molecule H₂O₂. *Biochem J* 414:53–61

Echevarria M, Windhager EE, Tate SS, Frindt G (1994) Cloning and expression of AQP3, a water channel from the medullary collecting duct of rat kidney. *Proc Natl Acad Sci USA* 91:10997–11001

Eckert M, Biela A, Siefritz F, Kaldenhoff IR (1999) New aspects of plant aquaporin regulation and specificity. *J Exp Bot* 50:1541–1545

Endeward V, Musa-Aziz R, Cooper GJ, Chen LM, Pelletier MF, Virkki LV, Supuran CT, King LS, Boron WF, Gros G (2006) Evidence that aquaporin 1 is a major pathway for CO₂ transport across the human erythrocyte membrane. *FASEB J* 20:1974–1981

Fenton RA, Moeller HB, Zelenina M, Snaebjornsson MT, Holen T, MacAulay N (2010) Differential water permeability and regulation of three aquaporin 4 isoforms. *Cell Mol Life Sci* 67:829–840

Fetter K, Van Wilder V, Moshelion M, Chaumont F (2004) Interactions between plasma membrane aquaporins modulate their water channel activity. *Plant Cell* 16:215–228

Finn RN, Cerda J (2011) Aquaporin evolution in fishes. *Front Physiol* 2:44

Froger A, Clemens D, Kalman K, Nemeth-Cahalan KL, Schilling TF, Hall JE (2010) Two distinct aquaporin 0s required for development and transparency of the zebrafish lens. *Investig Ophthalmol Vis Sci* 51:6582–6592

Fu D (2000) Structure of a glycerol-conducting channel and the basis for its selectivity. *Science* 290:481–486

Fushimi K, Uchida S, Hara Y, Hirata Y, Marumo F, Sasaki S (1993) Cloning and expression of apical membrane water channel of rat kidney collecting tubule. *Nature* 361:549–552

Gaspar M (2003) Cloning and characterization of ZmPIP1-5b, an aquaporin transporting water and urea. *Plant Sci* 165:21–31

Gerbeau P, Güçlü J, Ripoche P, Maurel C (1999) Aquaporin Nt-TIPa can account for the high permeability of tobacco cell vacuolar membrane to small neutral solutes. *Plant J* 18:577–587

Geyer RR, Musa-Aziz R, Qin X, Boron WF (2013) Relative CO₂/NH₃ selectivities of mammalian aquaporins 0–9. *Am J Physiol Cell Physiol* 304:C985–C994

Goldraij A, Polacco JC (2000) Arginine degradation by arginase in mitochondria of soybean seedling cotyledons. *Planta* 210(4):652–658

Gomes D, Agasse A, Thiebaud P, Delrot S, Geros H, Chaumont F (2009) Aquaporins are multifunctional water and solute transporters highly divergent in living organisms. *Biochim Biophys Acta* 1788:1213–1228

Gu R, Chen X, Zhou Y, Yuan L (2012) Isolation and characterization of three maize aquaporin genes, ZmNIP2;1, ZmNIP2;4 and ZmTIP4;4 involved in urea transport. *BMB Rep* 45:96–101

Guenther JF, Roberts DM (2000) Water-selective and multifunctional aquaporins from *Lotus japonicus* nodules. *Planta* 210:741–748

Hachez C, Besserer A, Chevalier AS, Chaumont F (2013) Insights into plant plasma membrane aquaporin trafficking. *Trends Plant Sci* 18:344–352

Hara-Chikuma M, Chikuma S, Sugiyama Y, Kabashima K, Verkman AS, Inoue S, Miyachi Y (2012) Chemokine-dependent T cell migration requires aquaporin-3-mediated hydrogen peroxide uptake. *J Exp Med* 209:1743–1752

Hatakeyama S, Yoshida Y, Tani T, Koyama Y, Nihei K, Ohshiro K, Kamiie JI, Yaoita E, Suda T, Hatakeyama K, Yamamoto T (2001) Cloning of a new aquaporin (AQP10) abundantly expressed in duodenum and jejunum. *Biochem Biophys Res Commun* 287:814–819

Hazama A, Kozono D, Guggino WB, Agre P, Yasui M (2002) Ion permeation of AQP6 water channel protein. Single channel recordings after Hg²⁺ activation. *J Biol Chem* 277:29224–29230

Heckwolf M, Pater D, Hanson DT, Kaldenhoff R (2011) The *Arabidopsis thaliana* aquaporin AtPIP1;2 is a physiologically relevant CO₂ transport facilitator. *Plant J* 67:795–804

Hedfalk K, Tornroth-Horsefield S, Nyblom M, Johanson U, Kjellbom P, Neutze R (2006) Aquaporin gating. *Curr Opin Struct Biol* 16:447–456

Herrera M, Hong NJ, Garvin JL (2006) Aquaporin-1 transports NO across cell membranes. *Hypertension* 48:157–164

Heymann JB, Engel A (1999) Aquaporins: phylogeny, structure, and physiology of water channels. *News Physiol Sci* 14:187–193

Holm LM, Klaerke DA, Zeuthen T (2004) Aquaporin 6 is permeable to glycerol and urea. *Pflug Arch* 448:181–186

Holm LM, Jahn TP, Moller AL, Schjoerring JK, Ferri D, Klaerke DA, Zeuthen T (2005) NH₃ and NH₄⁺ permeability in aquaporin-expressing *Xenopus* oocytes. *Pflug Arch* 450:415–428

Hooijmair C, Rhee JY, Kwak KJ, Chung GC, Horie T, Katsuhara M, Kang H (2012) Hydrogen peroxide permeability of plasma

membrane aquaporins of *Arabidopsis thaliana*. *J Plant Res* 125:147–153

Hughes AL, Ekollu V, Friedman R, Rose JR (2005) Gene family content-based phylogeny of prokaryotes: the effect of criteria for inferring homology. *Syst Biol* 54:268–276

Hwang JH, Ellingson SR, Roberts DM (2010) Ammonia permeability of the soybean nodulin 26 channel. *FEBS Lett* 584:4339–4343

Ikeda M, Beitz E, Kozono D, Guggino WB, Agre P, Yasui M (2002) Characterization of aquaporin-6 as a nitrate channel in mammalian cells. Requirement of pore-lining residue threonine 63. *J Biol Chem* 277:39873–39879

Ikeda M, Andoo A, Shimono M, Takamatsu N, Taki A, Muta K, Matsushita W, Uechi T, Matsuzaki T, Kenmochi N, Takata K, Sasaki S, Ito K, Ishibashi K (2011) The NPC motif of aquaporin-11, unlike the NPA motif of known aquaporins, is essential for full expression of molecular function. *J Biol Chem* 286:3342–3350

Ishibashi K, Sasaki S, Fushimi K, Uchida S, Kuwahara M, Saito H, Furukawa T, Nakajima K, Yamaguchi Y, Gojobori T, Marumo F (1994) Molecular cloning and expression of a member of the aquaporin family with permeability to glycerol and urea in addition to water expressed at the basolateral membrane of kidney collecting duct cells. *Proc Natl Acad Sci USA* 91:6269–6273

Ishibashi K, Kuwahara M, Gu Y, Kageyama Y, Tohsaka A, Suzuki F, Marumo F, Sasaki S (1997) Cloning and functional expression of a new water channel abundantly expressed in the testis permeable to water, glycerol, and urea. *J Biol Chem* 272:20782–20786

Ishibashi K, Morinaga T, Kuwahara M, Sasaki S, Imai M (2002) Cloning and identification of a new member of water channel (AQP10) as an aquaglyceroporin. *Biochim Biophys Acta* 1576:335–340

Ishibashi K, Kondo S, Hara S, Morishita Y (2011) The evolutionary aspects of aquaporin family. *Am J Physiol Regul Integr Comp Physiol* 300:R566–R576

Ishikawa F, Suga S, Uemura T, Sato MH, Maeshima M (2005) Novel type aquaporin SIPs are mainly localized to the ER membrane and show cell-specific expression in *Arabidopsis thaliana*. *FEBS Lett* 579:5814–5820

Jahn TP, Moller AL, Zeuthen T, Holm LM, Klaerke DA, Mohsin B, Kuhlbrandt W, Schjoerring JK (2004) Aquaporin homologues in plants and mammals transport ammonia. *FEBS Lett* 574:31–36

Johanson U, Karlsson M, Johansson I, Gustavsson S, Sjövall S, Fraysse L, Weig AR, Kjellbom P (2001) The complete set of genes encoding major intrinsic proteins in *Arabidopsis* provides a framework for a new nomenclature for major intrinsic proteins in plants. *Plant Physiol* 126:1358–1369

Johansson I, Karlsson M, Shukla VK, Chrispeels MJ, Larsson C, Kjellbom P (1998) Water transport activity of the plasma membrane aquaporin PM28A is regulated by phosphorylation. *Plant Cell* 10:451–459

Jozefkowicz C, Rosi P, Sigaut L, Soto G, Pietrasanta LI, Amodeo G, Alleva K (2013) Loop A is critical for the functional interaction of two *Beta vulgaris* PIP aquaporins. *PLoS ONE* 8:e57993

Jung JS, Bhat RV, Preston GM, Guggino WB, Baraban JM, Agre P (1994) Molecular characterization of an aquaporin cDNA from brain: candidate osmoreceptor and regulator of water balance. *Proc Natl Acad Sci USA* 91:13052–13056

Kamiya T, Fujiwara T (2009) *Arabidopsis* NIP1;1 transports antimonic acid and determines antimonic acid sensitivity. *Plant Cell Physiol* 50:1977–1981

Kamiya T, Tanaka M, Mitani N, Ma JF, Maeshima M, Fujiwara T (2009) NIP1;1, an aquaporin homolog, determines the arsenite sensitivity of *Arabidopsis thaliana*. *J Biol Chem* 284:2114–2120

Kammerloher W, Fischer U, Piechottka GP, Schäffner AR (1994) Water channels in the plant plasma membrane cloned by immunoselection from a mammalian expression system. *Plant J* 6:187–199

Kishida K, Kuriyama H, Funahashi T, Shimomura I, Kihara S, Ouchi N, Nishida M, Nishizawa H, Matsuda M, Takahashi M, Hotta K, Nakamura T, Yamashita S, Tochino Y, Matsuzawa Y (2000) Aquaporin adipose, a putative glycerol channel in adipocytes. *J Biol Chem* 275:20896–20902

Klebl F, Wolf M, Sauer N (2003) A defect in the yeast plasma membrane urea transporter Dur3p is complemented by CpNIP1, a Nod26-like protein from zucchini (*Cucurbita pepo* L.), and by *Arabidopsis thaliana* δ -TIP or γ -TIP. *FEBS Lett* 547:69–74

Kojima S, Bohner A, von Wieren N (2006) Molecular mechanisms of urea transport in plants. *J Membr Biol* 212:83–91

Koonin EV (2003) Horizontal gene transfer: the path to maturity. *Mol Microbiol* 50:725–727

Koonin EV, Makarova KS, Aravind L (2001) Horizontal gene transfer in prokaryotes: quantification and classification. *Annu Rev Microbiol* 55:709–742

Koyama Y, Yamamoto T, Kondo D, Funaki H, Yaoita E, Kawasaki K, Sato N, Hatakeyama K, Kihara I (1997) Molecular cloning of a new aquaporin from rat pancreas and liver. *J Biol Chem* 272:30329–30333

Li WH (1997) Molecular evolution. Sinauer Associates, Sunderland

Li GW, Peng YH, Yu X, Zhang MH, Cai WM, Sun WN, Su WA (2008) Transport functions and expression analysis of vacuolar membrane aquaporins in response to various stresses in rice. *J Plant Physiol* 165:1879–1888

Li H, Chen H, Steinbronn C, Wu B, Beitz E, Zeuthen T, Voth GA (2011) Enhancement of proton conductance by mutations of the selectivity filter of aquaporin-1. *J Mol Biol* 407:607–620

Ligaba A, Katsuhara M, Shibasaki M, Djiré G (2011) Abiotic stresses modulate expression of major intrinsic proteins in barley (*Hordeum vulgare*). *C R Biol* 334:127–139

Liu Z, Shen J, Carbrey JM, Mukhopadhyay R, Agre P, Rosen BP (2002) Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9. *Proc Natl Acad Sci USA* 99:6053–6058

Liu LH, Ludewig U, Gassert B, Frommer WB, von Wieren N (2003) Urea transport by nitrogen-regulated tonoplast intrinsic proteins in *Arabidopsis*. *Plant Physiol* 133:1220–1228

Liu Z, Carbrey JM, Agre P, Rosen BP (2004) Arsenic trioxide uptake by human and rat aquaglyceroporins. *Biochem Biophys Res Commun* 316:1178–1185

Liu K, Nagase H, Huang CG, Calamita G, Agre P (2006) Purification and functional characterization of aquaporin-8. *Biol Cell* 98:153–161

Loque D, Ludewig U, Yuan L, von Wieren N (2005) Tonoplast intrinsic proteins AtTIP2;1 and AtTIP2;3 facilitate NH₃ transport into the vacuole. *Plant Physiol* 137:671–680

Ma T, Frigeri A, Tsai ST, Verbavatz JM, Verkman AS (1993) Localization and functional analysis of CHIP28k water channels in stably transfected Chinese hamster ovary cells. *J Biol Chem* 268:22756–22764

Ma T, Yang B, Kuo WL, Verkman AS (1996) cDNA cloning and gene structure of a novel water channel expressed exclusively in human kidney: evidence for a gene cluster of aquaporins at chromosome locus 12q13. *Genomics* 35:543–550

Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Katsuhara M, Ishiguro M, Murata Y, Yano M (2006) A silicon transporter in rice. *Nature* 440:688–691

Ma JF, Yamaji N, Mitani N, Xu XY, Su YH, McGrath SP, Zhao FJ (2008) Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. *Proc Natl Acad Sci USA* 105:9931–9935

Maeshima M, Ishikawa F (2008) ER membrane aquaporins in plants. *Pflug Arch* 456:709–716

Marchler-Bauer A, Zheng C, Chitsaz F, Derbyshire MK, Geer LY, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI, Lanczycki CJ, Lu F, Lu S, Marchler GH, Song JS, Thanki N, Yamashita RA, Zhang D, Bryant SH (2013) CDD: conserved domains and protein three-dimensional structure. *Nucleic Acids Res* 41:D348–D352

Marinelli RA, Pham L, Agre P, LaRusso NF (1997) Secretin promotes osmotic water transport in rat cholangiocytes by increasing aquaporin-1 water channels in plasma membrane. Evidence for a secretin-induced vesicular translocation of aquaporin-1. *J Biol Chem* 272:12984–12988

Matsumoto T, Lian HL, Su WA, Tanaka D, Liu C, Iwasaki I, Kitagawa Y (2009) Role of the aquaporin PIP1 subfamily in the chilling tolerance of rice. *Plant Cell Physiol* 50:216–229

Maurel C, Reizer J, Schroeder JI, Chrispeels MJ (1993) The vacuolar membrane protein gamma-TIP creates water specific channels in *Xenopus* oocytes. *EMBO J* 12:2241–2247

McDermott JR, Jiang X, Beene LC, Rosen BP, Liu Z (2010) Pentavalent methylated arsenicals are substrates of human AQP9. *Biometals* 23:119–127

Meinild AK (1998) Bidirectional water fluxes and specificity for small hydrophilic molecules in aquaporins 0–5. *J Biol Chem* 273:32446–32451

Mitani N, Yamaji N, Ma JF (2008) Characterization of substrate specificity of a rice silicon transporter, Lsi1. *Pflug Arch* 456:679–686

Mitani N, Yamaji N, Ma JF (2009) Identification of maize silicon influx transporters. *Plant Cell Physiol* 50:5–12

Mitani-Ueno N, Yamaji N, Zhao FJ, Ma JF (2011) The aromatic/arginine selectivity filter of NIP aquaporins plays a critical role in substrate selectivity for silicon, boron, and arsenic. *J Exp Bot* 62:4391–4398

Mizutani M, Watanabe S, Nakagawa T, Maeshima M (2006) Aquaporin NIP2;1 is mainly localized to the ER membrane and shows root-specific accumulation in *Arabidopsis thaliana*. *Plant Cell Physiol* 47:1420–1426

Mobley HL, Island MD, Hausinger RP (1995) Molecular biology of microbial ureases. *Microbiol Rev* 59:451–480

Montpetit J, Vivancos J, Mitani-Ueno N, Yamaji N, Remus-Borel W, Belzile F, Ma JF, Belanger RR (2012) Cloning, functional characterization and heterologous expression of TaLsi1, a wheat silicon transporter gene. *Plant Mol Biol* 79:35–46

Mosa KA, Kumar K, Chhikara S, McDermott J, Liu Z, Musante C, White JC, Dhankher OP (2012) Members of rice plasma membrane intrinsic proteins subfamily are involved in arsenite permeability and tolerance in plants. *Transgenic Res* 21: 1265–1277

Moshelion M (2002) Plasma membrane aquaporins in the motor cells of *Samanea saman*: diurnal and circadian regulation. *Plant Cell Online* 14:727–739

Mulders SM, Preston GM, Deen PM, Guggino WB, van Os CH, Agre P (1995) Water channel properties of major intrinsic protein of lens. *J Biol Chem* 270:9010–9016

Musa-Aziz R, Chen LM, Pelletier MF, Boron WF (2009) Relative CO_2/NH_3 selectivities of AQP1, AQP4, AQP5, AmtB, and RhAG. *Proc Natl Acad Sci USA* 106:5406–5411

Nakhoul NL, Davis BA, Romero MF, Boron WF (1998) Effect of expressing the water channel aquaporin-1 on the CO_2 permeability of *Xenopus* oocytes. *Am J Physiol* 274:C543–C548

Neely JD, Christensen BM, Nielsen S, Agre P (1999) Heterotetrameric composition of aquaporin-4 water channels. *Biochemistry* 38:11156–11163

Novichkov PS, Omelchenko MV, Gelfand MS, Mironov AA, Wolf YI, Koonin EV (2004) Genome-wide molecular clock and horizontal gene transfer in bacterial evolution. *J Bacteriol* 186:6575–6585

Otto B, Uehlein N, Sdorra S, Fischer M, Ayaz M, Belastegui-Macadam X, Heckwolf M, Lachnit M, Pede N, Priem N, Reinhard A, Siegfart S, Urban M, Kaldenhoff R (2010) Aquaporin tetramer composition modifies the function of tobacco aquaporins. *J Biol Chem* 285:31253–31260

Park JH, Saier MH Jr (1996) Phylogenetic characterization of the MIP family of transmembrane channel proteins. *J Membr Biol* 153:171–180

Pedrozo HA, Schwartz Z, Dean DD, Wiederhold ML, Boyan BD (1996) Regulation of statocilia mineralization in *Aplysia californica* in vitro. *Connect Tissue Res* 35:317–323

Phillips AJ (2006) Homology assessment and molecular sequence alignment. *J Biomed Inform* 39:18–33

Prasad GV, Coury LA, Finn F, Zeidel ML (1998) Reconstituted aquaporin 1 water channels transport CO_2 across membranes. *J Biol Chem* 273:33123–33126

Preston GM, Carroll TP, Guggino WB, Agre P (1992) Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP28 protein. *Science* 256:385–387

Quigley F, Rosenberg JM, Shachar-Hill Y, Bohnert HJ (2002) From genome to function: the *Arabidopsis* aquaporins. *Genome Biol* 3:RESEARCH0001

Raina S, Preston GM, Guggino WB, Agre P (1995) Molecular cloning and characterization of an aquaporin cDNA from salivary, lacrimal, and respiratory tissues. *J Biol Chem* 270:1908–1912

Rivers RL, Dean RM, Chandy G, Hall JE, Roberts DM, Zeidel ML (1997) Functional analysis of nodulin 26, an aquaporin in soybean root nodule symbiosomes. *J Biol Chem* 272:16256–16261

Rodela TM, Ballantyne JS, Wright PA (2008) Carrier-mediated urea transport across the mitochondrial membrane of an elasmobranch (*Raja erinacea*) and a teleost (*Oncorhynchus mykiss*) fish. *Am J Physiol Regul Integr Comp Physiol* 294:R1947–R1957

Sakurai J, Ahamed A, Murai M, Maeshima M, Uemura M (2008) Tissue and cell-specific localization of rice aquaporins and their water transport activities. *Plant Cell Physiol* 49:30–39

Schnurbusch T, Hayes J, Hrmova M, Baumann U, Ramesh SA, Tyerman SD, Langridge P, Sutton T (2010) Boron toxicity tolerance in barley through reduced expression of the multi-functional aquaporin HvNIP2;1. *Plant Physiol* 153:1706–1715

Schuurmans JA, van Dongen JT, Rutjens BP, Boonman A, Pieterse CM, Borstlap AC (2003) Members of the aquaporin family in the developing pea seed coat include representatives of the PIP, TIP, and NIP subfamilies. *Plant Mol Biol* 53:633–645

Soria LR, Marrone J, Calamita G, Marinelli RA (2013) Ammonia detoxification via ureagenesis in rat hepatocytes involves mitochondrial aquaporin-8 channels. *Hepatology* 57:2061–2071

Soto G, Alleva K, Mazzella MA, Amodeo G, Muschietti JP (2008) AtTIP1;3 and AtTIP5;1, the only highly expressed *Arabidopsis* pollen-specific aquaporins, transport water and urea. *FEBS Lett* 582:4077–4082

Soto G, Fox R, Ayub N, Alleva K, Guaimas F, Erijman EJ, Mazzella A, Amodeo G, Muschietti J (2010) TIP5;1 is an aquaporin specifically targeted to pollen mitochondria and is probably involved in nitrogen remobilization in *Arabidopsis thaliana*. *Plant J* 64:1038–1047

Soto G, Alleva K, Amodeo G, Muschietti J, Ayub ND (2012) New insight into the evolution of aquaporins from flowering plants and vertebrates: orthologous identification and functional transfer is possible. *Gene* 503:165–176

Stein WD, Danielli JF (1956) Structure and function in red cell permeability. *Discuss Faraday Soc* 21:238–251

Sui H, Han BG, Lee JK, Walian P, Jap BK (2001) Structural basis of water-specific transport through the AQP1 water channel. *Nature* 414:872–878

Takano J, Wada M, Ludewig U, Schaaf G, von Wieren N, Fujiwara T (2006) The *Arabidopsis* major intrinsic protein NIP5;1 is essential for efficient boron uptake and plant development under boron limitation. *Plant Cell* 18:1498–1509

Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739

Tanaka M, Wallace IS, Takano J, Roberts DM, Fujiwara T (2008) NIP6;1 is a boric acid channel for preferential transport of boron to growing shoot tissues in *Arabidopsis*. *Plant Cell* 20: 2860–2875

Tingaud-Sequeira A, Calusinska M, Finn RN, Chauvigne F, Lozano J, Cerdá J (2010) The zebrafish genome encodes the largest vertebrate repertoire of functional aquaporins with dual paralogy and substrate specificities similar to mammals. *BMC Evol Biol* 10:38

Tournaire-Roux C, Sutka M, Javot H, Gout E, Gerbeau P, Luu DT, Bligny R, Maurel C (2003) Cytosolic pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature* 425:393–397

Tsukaguchi H, Weremowicz S, Morton CC, Hediger MA (1999) Functional and molecular characterization of the human neutral solute channel aquaporin-9. *Am J Physiol* 277:F685–F696

Uehlein N, Lovisolo C, Siefritz F, Kaldenhoff R (2003) The tobacco aquaporin NtAQP1 is a membrane CO_2 pore with physiological functions. *Nature* 425:734–737

Verbavatz JM, Brown D, Sabolic I, Valenti G, Ausiello DA, Van Hoek AN, Ma T, Verkman AS (1993) Tetrameric assembly of CHIP28 water channels in liposomes and cell membranes: a freeze-fracture study. *J Cell Biol* 123:605–618

Wallace IS, Roberts DM (2004) Homology modeling of representative subfamilies of *Arabidopsis* major intrinsic proteins. Classification based on the aromatic/arginine selectivity filter. *Plant Physiol* 135:1059–1068

Wallace IS, Roberts DM (2005) Distinct transport selectivity of two structural subclasses of the nodulin-like intrinsic protein family of plant aquaglyceroporin channels. *Biochemistry* 44: 16826–16834

Wallace IS, Shakesby AJ, Hwang JH, Choi WG, Martinkova N, Douglas AE, Roberts DM (2012) *Acyrthosiphon pisum* AQP2: a multifunctional insect aquaglyceroporin. *Biochim Biophys Acta* 1818:627–635

Weig AR, Jakob C (2000) Functional identification of the glycerol permease activity of *Arabidopsis thaliana* NLM1 and NLM2 proteins by heterologous expression in *Saccharomyces cerevisiae*. *FEBS Lett* 481:293–298

Wu B, Beitz E (2007) Aquaporins with selectivity for unconventional permeants. *Cell Mol Life Sci* 64:2413–2421

Wudick MM, Luu DT, Maurel C (2009) A look inside: localization patterns and functions of intracellular plant aquaporins. *New Phytol* 184:289–302

Yakata K, Hiroaki Y, Ishibashi K, Sohara E, Sasaki S, Mitsuoka K, Fujiyoshi Y (2007) Aquaporin-11 containing a divergent NPA motif has normal water channel activity. *Biochim Biophys Acta* 1768:688–693

Yakata K, Tani K, Fujiyoshi Y (2011) Water permeability and characterization of aquaporin-11. *J Struct Biol* 174:315–320

Yan Y, Yang J, Dou Y, Chen M, Ping S, Peng J, Lu W, Zhang W, Yao Z, Li H, Liu W, He S, Geng L, Zhang X, Yang F, Yu H, Zhan Y, Li D, Lin Z, Wang Y, Elmerich C, Lin M, Jin Q (2008) Nitrogen fixation island and rhizosphere competence traits in the genome of root-associated *Pseudomonas stutzeri* A1501. *Proc Natl Acad Sci USA* 105:7564–7569

Yang B, Verkman AS (1997) Water and glycerol permeabilities of aquaporins 1–5 and MIP determined quantitatively by expression of epitope-tagged constructs in *Xenopus* oocytes. *J Biol Chem* 272:16140–16146

Yasui M, Hazama A, Kwon TH, Nielsen S, Guggino WB, Agre P (1999) Rapid gating and anion permeability of an intracellular aquaporin. *Nature* 402:184–187

Yu JJ, Smithson SL, Thomas PW, Kirkland TN, Cole GT (1997) Isolation and characterization of the urease gene (URE) from the pathogenic fungus *Coccidioides immitis*. *Gene* 198:387–391

Zampighi GA, Hall JE, Kreman M (1985) Purified lens junctional protein forms channels in planar lipid films. *Proc Natl Acad Sci USA* 82:8468–8472

Zardoya R (2005) Phylogeny and evolution of the major intrinsic protein family. *Biol Cell* 97:397–414

Zardoya R, Villalba S (2001) A phylogenetic framework for the aquaporin family in eukaryotes. *J Mol Evol* 52:391–404

Zardoya R, Ding X, Kitagawa Y, Chrispeels MJ (2002) Origin of plant glycerol transporters by horizontal gene transfer and functional recruitment. *Proc Natl Acad Sci USA* 99: 14893–14896

Zelazny E, Borst JW, Muylaert M, Batoko H, Hemminga MA, Chaumont F (2007) FRET imaging in living maize cells reveals that plasma membrane aquaporins interact to regulate their subcellular localization. *Proc Natl Acad Sci USA* 104: 12359–12364

Zeuthen T, Meinild AK, Klaerke DA, Loo DD, Wright EM, Belhage B, Litman T (1997) Water transport by the $\text{Na}^+/\text{glucose}$ cotransporter under isotonic conditions. *Biol Cell* 89:307–312