

Allicin inhibits cell growth and induces apoptosis through the mitochondrial pathway in HL60 and U937 cells

Talia Miron^{a,*}, Meir Wilchek^a, Ayala Sharp^b, Yoshihito Nakagawa^c, Makoto Naoi^c,
Yoshinori Nozawa^c, Yukihiro Akao^c

^aDepartment of Biological Chemistry, Weizmann Institute of Science, Rehovot 76100, Israel

^bDepartment of Biological Services, Weizmann Institute of Science, Rehovot 76100, Israel

^cGifu International Institute of Biotechnology, Kakamigahara, Gifu 504-0838, Japan

Received 22 April 2007; received in revised form 12 June 2007; accepted 20 June 2007

Abstract

In this article, the effects of allicin, a biological active compound of garlic, on HL60 and U937 cell lines were examined. Allicin induced growth inhibition and elicited apoptotic events such as blebbing, mitochondrial membrane depolarization, cytochrome *c* release into the cytosol, activation of caspase 9 and caspase 3 and DNA fragmentation. Pretreatment of HL60 cells with cyclosporine A, an inhibitor of the mitochondrial permeability transition pore (mPTP), inhibited allicin-treated cell death. HL60 cell survival after 1 h pretreatment with cyclosporine A, followed by 16 h in presence of allicin (5 μ M) was ~80% compared to allicin treatment alone (~50%). Also *N*-acetyl cysteine, a reduced glutathione (GSH) precursor, prevented cell death. The effects of cyclosporine A and *N*-acetyl cysteine suggest the involvement of mPTP and intracellular GSH level in the cytotoxicity. Indeed, allicin depleted GSH in the cytosol and mitochondria, and buthionine sulfoximine, a specific inhibitor of GSH synthesis, significantly augmented allicin-induced apoptosis. In HL60 cells treated with allicin (5 μ M, 30 min) the redox state for 2GSH/oxidized glutathione shifted from $E_{\text{GSH}} -240$ to -170 mV. The same shift was observed in U937 cells treated with allicin at a higher concentration for a longer period of incubation (20 μ M, 2 h). The apoptotic events induced by various concentrations of allicin correlate to intracellular GSH levels in the two cell types tested (HL60: 3.7 nmol/10⁶ cells; U937: 7.7 nmol/10⁶ cells). The emerging mechanistic basis for the antiproliferative function of allicin, therefore, involves the activation of the mitochondrial apoptotic pathway by GSH depletion and by changes in the intracellular redox status.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Allicin; Apoptosis; Garlic; GSH; HL60; U937

1. Introduction

Allicin, diallyl thiosulfinate, is the main biologically active compound derived from garlic. It is produced by the interaction of the enzyme alliinase (alliin lyase; EC 4.4.1.4) with its substrate alliin (*S*-allyl-L-cysteine sulfoxide) [1].


Abbreviations: BSO, D,L-buthionine *S,R*-sulfoximine; CsA, Cyclosporin A; DTNB, 5,5'-dithiobis-(2-nitrobenzoic acid); GGCS, γ -Glutamyl-L-cysteine synthetase; GSH, Reduced glutathione; GSSA, *S*-allylmercaptogluthathione; GSSG, Oxidized glutathione; mPTP, Mitochondrial permeability transition pore; NAC, *N*-acetyl-L-cysteine; XTT, 2,3-Bis (2-methoxy-4-nitro-5-sulphophenyl)-2*H*-tetrazolium-5-carboxanilide sodium salt.

* Corresponding author. Tel.: +972 8 9343627; fax: +972 8 9468256.

E-mail address: talia.miron@weizmann.ac.il (T. Miron).

Since alliinase and alliin are enclosed in different compartments within the garlic clove cells, intact garlic cloves do not contain allicin. When the garlic clove is crushed, alliin and alliinase interact, to form allicin, pyruvic acid and ammonia (Scheme 1). Allicin became an object of interest due to its potential to confer a vast spectrum of health beneficial effects including: anti microbial, antifungal and antiparasitic [2], antihypertensive [3], cardioprotective [4–6], anti-inflammatory [7] and anticancer activities [2,8,9].

In vitro studies of allicin's effect on human mammary MCF-7 cancer cells revealed that its antiproliferative activity is accompanied by accumulation of cells in the regulatory checkpoints, G₀/G₁ and G₂/M, of the cell cycle [9]. Interestingly, also other ally sulfur compound cause similar effects [10]. Allicin was shown to induce apoptosis in gastric



cancer SGC-7901 cells. The cause of apoptosis was related to decreased telomerase activity, an enzyme which alllicin inhibits in a time- and dose- dependent manner [11]. Alllicin induces apoptosis in human cervical cancer SiHa cells and mouse fibroblast-like L-929 cells, manifested through the appearance of characteristic apoptotic morphological changes in apoptotic bodies, through DNA fragmentation, and activation of caspases 8, 9 and 3 [12]. Alllicin was also shown to induce apoptosis in human epithelial carcinoma through a caspase-independent pathway, mediated by the release of apoptotic-inducing factor (AIF) from mitochondria and protein kinase A (PKA) activation [13]. More results describing the effects of alllicin or other garlic-derived products on various proteins participate in the apoptotic process have already been reported [14]. However, the mechanism underlying the induction of cell death by alllicin has not been fully elucidated.

Apoptosis is an ordered cascade of events that culminates in cell death. Two main pathways of apoptosis have been characterized. The extrinsic pathway is initiated through ligand stimulation of the cell surface death receptors such as TNFR or CD95R. In this pathway, cell death is executed via a cascade of proteolytic events with the sequential activation of caspase 8 and caspase 3. The intrinsic pathway is triggered by mitochondrial stress caused by various factors such as DNA damage, oxidative stress and heat shock (reviewed [15]). This pathway is initiated through the release of signal factors from mitochondria as a consequence of mitochondrial membrane permeability transition. Such changes lead to translocation of pro- and antiapoptotic proteins across the mitochondrial membranes [16,17]. Among these proteins is cytochrome *c*, which is released from the mitochondria to the cytosol and participates with other molecules in the formation of a complex that activates caspase 9, which in turn activates caspase 3. The activation of these caspases leads to the final fragmentation of nuclear DNA, with the typical apoptotic morphological manifestations.

Allicin easily diffuses through cell membranes (diffusion coefficient $5 \times 10^{-8} \text{ cm}^2 \text{ s}^{-1}$) [18] and exerts its biological effects by reacting with free thiols within the cell. In living

cells, reduced glutathione (GSH) is the major free thiol participating in cellular redox reactions and mixed disulfide formation. GSH is therefore the main cellular target of allicin reaction (Scheme 2). Both allicin and its glutathione derivative, *S*-allylmercaptogluthathione (GSSA), can prevent the formation of free radicals. GSSA is similar in this preventive capacity to GSH, both being less effective in this antioxidant activity than allicin [19–21]. Allicin can scavenge the chain-carrying peroxy radicals of the substrates by transferring its allylic hydrogen to the oxidized substrate. This renders it a stronger antioxidant than its derivatives [22].

Alliin is a short-lived compound, which rapidly reacts with free intracellular thiol groups [18,19,22]. It was found to disintegrate in the blood a few minutes after its administration, both in vitro in human blood [23] and in vivo in rats [24]. Therefore, the therapeutic effect of alliin administered orally may be restricted to targets that are close to the gastrointestinal tract. However, its main oxidation products, *S*-allylmercapto-glutathione and *S*-allylmercapto-cysteine, could exert their action in more remote sites within the body because they are more stable. The rapid disappearance of alliin can be exploited in tumor cells targeting. We previously showed that alliin kills tumor cells in vitro [25,26] and also in vivo if generated on their surface by conjugating alliinase to monoclonal antibodies directed to specific cell-surface receptors, such as ErbB2, overexpressed in breast and ovarian cancer [25]; CD20, a receptor expressed at high levels in human B chronic lymphocytic leukemia and other B-cell lymphomas [26].

The redox environment of a cell reflects the sum of the products of the reduction potential and the reducing capacity of the linked redox couples operating within the cell [27,28]. Glutathione is considered to be the major thiol-disulfide redox buffer of cells [29]. The redox state of the 2GSH/oxidized glutathione (GSSG) redox couple depends on their molar ratio. Upon reacting with GSH, allicin causes a decrease in free GSH concentration and an increase in mixed-disulfide glutathione products, which leads to an increase in the reduction potential values. Allicin also reacts with other



free sulfhydryl (SH)-bearing molecules in the cell, such as cysteine and SH residues in proteins, yet their relatively low concentrations contribute only little towards changing the reduction potential of the cell upon oxidation. The reduction potential of cells was proposed to reflect their growth cycle. Accordingly, the reduction potential of proliferating cells (pH 7.0) is $E_{\text{GSH}} -240$ mV. Increased values ($E_{\text{GSH}} -200$ mV) represent differentiation, and a further oxidative shift to a higher value ($E_{\text{GSH}} -170$ mV) elicits apoptosis [27]. In this study, we aimed to examine the initial events leading to apoptosis upon allicin treatment and the dependence of the apoptotic pathway on the reduction potential of cells.

2. Materials and methods

Alliin was synthesized from L-cysteine and allyl bromide followed by H_2O_2 oxidation [30]. Allicin (purity ~98%) was produced by applying synthetic alliin onto an immobilized alliinase column and its concentration was determined by high-performance liquid chromatography, as previously described [31].

2,3-Bis (2-methoxy-4-nitro-5-sulphophenyl)-2 *H*-tetrazolium-5-carboxanilide (XTT) sodium salt; D,L-buthionine S, R-sulfoximine (BSO); cyclosporin A (CsA); *N*-acetyl-L-cysteine (NAC); 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB); metaphosphoric acid; phenazine methosulfate (PMS) and 2-vinylpyridine were purchased from Sigma Chemical (St. Louis, MO, USA).

2.1. Cell culture, cell viability and morphological studies.

HL60 human promyelocytic leukemia-derived cells and U937 human myelomonocytic cells were maintained in RPMI-1640 supplemented with 2 mM L-glutamine, 10% (v/v) fetal bovine serum and antibiotics. Cell viability was measured using the XTT assay, based on the reduction of tetrazolium salt to soluble formazan compounds by mitochondrial enzymes. Cells (15,000–20,000 cells/well) were seeded in a 96-well plate. After 16-h incubation with allicin, in the presence or absence of NAC (0.1–1.0 mM) or after 16 h incubation with BSO (0.1 mM) alone, 50 μl of XTT/PMS mixture (50 μM PMS, 0.1% XTT in medium) were added onto the cells. After an incubation period of 3–4 h at 37°C, the absorbance of the samples was measured in an enzyme-linked immunosorbent assay (ELISA) reader at 450 nm. SDS (1%, 10 μl /well) was added to reference wells before adding the XTT/PMS solution.

HL60 cells (300,000/ml) were pretreated with CsA for 1 h. Cells were seeded in a 96-well plate (20,000 cells/well) in the presence of allicin. The viability was tested by the XTT assay. The analysis of apoptotic morphological changes was done by staining cells with Hoechst 33342 at 37°C for 30 min in medium, washing with phosphate-buffered saline (PBS) and examining by fluorescence microscopy. For evaluating DNA ladder formation, cellular DNA was extracted from cells by ethanol precipitation of the phenol/chloroform extract [32].

After electrophoresis (5–10 μg DNA/lane) on an agarose (2.5%) gel, DNA was visualized by ethidium bromide staining.

Mitochondrial membrane potential was measured by using the fluorescent dye, Mito Tracker Red CMXRos (Molecular Probes, Eugene, OR, USA) that accumulates selectively in active mitochondria. Cells were washed with medium, incubated with Mito-Tracker Red CMXRos for 30 min at 37°C, washed with PBS and examined under fluorescence microscopy (Olympus, Tokyo).

Mitochondria were prepared from cultured cells ($5\text{--}10\times 10^7$) as described elsewhere [33]. Cells were harvested and washed with PBS (600 g, 7 min). Cell pellets were suspended in 0.5 ml cold hepes isotonic mitochondrial buffer (HIM) buffer (200 mM mannitol, 70 mM sucrose, 1 mM EGTA, 10 mM HEPES-KOH buffer, pH 7.5) containing protease inhibitors, incubated 30 min on ice and homogenized by multiple passages through a 25-gauge needle (5/8 in.). Nuclei and unbroken cells were removed (1300 g, 8 min at 4°C). Protein and free SH were determined in the cytosol, supernatant (10,000 g, 30 min at 4°C) and in the mitochondria-enriched fraction (pellet, dissolved in HIM buffer containing 1% Triton X-100). Protein assay was done with the Biuret Reagent [34] or the BCA protein assay kit (Pierce, Rockford, IL, USA).

2.2. Cell cycle analysis

HL60 cells were pretreated with CsA (5 μM , 1 h) and then cultured further for 20 h in the presence or absence of allicin. After harvesting, washing and resuspending in 0.25 ml PBS, an equal volume of 0.005% propidium iodide solution containing 0.01% heated-RNase A and 0.3% Triton X-100 was added. Cells were analyzed by flow cytometry using fluorescence-activated cell sorting (Becton Dickinson FACScan Instrument using CellQuest software (BD Bioscience, San Jose, CA, USA).

2.3. Electrophoresis and Western blot analysis

Cell pellet was resuspended in lysis buffers A or B [32]. Lysis buffer A (2 \times PBS, 0.1% sodium dodecyl sulfate (SDS), 1% Nonidet P-40, 0.5% sodium deoxycholate) containing protease inhibitors was used to analyze of caspases 3, 8 and 9. Lysis buffer B (250 mM sucrose, 20 mM HEPES-KOH buffer, pH 7.5, 10 mM KCl, 1.5 mM MgCl_2 , 1 mM EDTA, 1 mM EGTA, 1 mM dithiothreitol [DTT]) containing protease inhibitors was used to analyze the presence of cytochrome *c* in the cytosol as described above (Section 2.2). After centrifugation (10,000g, 30 min, 4°C) the supernatant was separated by SDS-polyacrylamide gel electrophoresis using 15% gel and transferred electrophoretically onto a polyvinylidene difluoride (PVDF) membrane (Du Pont, Boston, MA, USA). The membrane was incubated overnight at 4°C with the following antihuman antibodies: anti β -actin (Sigma–Aldrich), anti-caspase 3 (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-caspase 8 (MBL, Nagoya, Japan), anti

caspase-9 (Novus Biological, Littleton, CO, USA) and anti cytochrome *c* (Upstate Biotechnology, Lake Placid, NY, USA). The membranes were washed with tris-buffered saline Tween-20 (TBST), incubated with HRP-conjugated secondary antibody for 1 h at room temperature and washed with TBST. Proteins were detected with an enhanced chemiluminescence (ECL) detection kit (New England Biolabs) and a chemiluminescence detector (LAS-1000, Fuji, Japan).

2.4. Determination of cell volume

The volume of nontreated cells was determined by calculating sphere volumes based on diameter measurement of 100 cells. Since the morphology of treated cells was not spherical, cell volume was estimated by weighing the pellet of $2\text{--}4 \times 10^6$ cells and evaluating the volume on the assumption of 1/1 w/v.

2.5. Assay for glutathione and free SH

Cells were collected (600 g, 7 min, 4°C), washed with PBS and the pellets were stored at -80°C . After protein precipitation with 5% metaphosphoric acid ($0.2\text{--}0.3\text{ ml}/3 \times 10^6$ cells) by centrifugation (10,000g, for 30 min), the supernatant was used for GSH quantitation, and the pellet, dissolved in $0.2\text{--}0.3\text{ ml}$ of 0.5 M NaOH, was used for protein determination. To determine GSH in the supernatant, a glutathione assay kit (Calbiochem) was used. GSH and GSSG content were also determined by the glutathione reductase recycling assay before and after modification with 2-vinyl pyridine [35]. Samples ($40\text{ }\mu\text{l}$) were neutralized with 2 M triethanolamine ($10\text{ }\mu\text{l}$) in a 96-well plate. The reaction was started by adding $200\text{ }\mu\text{l}$ per well of 0.4 U/ml enzyme in 143 mM phosphate buffer pH 7.5 containing 0.3 mM reduced nicotinamide adenine dinucleotide phosphate, 0.6 mM DTNB and 6.25 mM EDTA. The initial rate of 5-thio-2-nitrobenzoic acid formation was monitored. Determination of GSSG was done 1 h after modification of the free SH with 9 M 2-vinyl pyridine ($1\text{ }\mu\text{l}/\text{well}$) at room temperature. Oxidized glutathione ($0\text{--}6\text{ nmol}/\text{well}$) served as a reference.

Total free SH content in cell extracts or in cytosolic and mitochondrial fractions ($50\text{ }\mu\text{l}$) was determined with DTNB in a 96-well plate. Samples were acidified with 5% metaphosphoric acid ($50\text{ }\mu\text{l}/\text{well}$) to stabilize the reduced state and neutralized with 2 M triethanolamine ($50\text{ }\mu\text{l}/\text{well}$). DTNB ($50\text{ }\mu\text{l}$ of 1 mM solution in 50 mM phosphate buffer pH 7.2 containing 2 mM EDTA) was added. After 10 min of incubation, the absorbance was measured at 412 nm using an ELISA reader. GSH ($0\text{--}20\text{ nmol}/\text{well}$) precalibrated with DTNB served as a reference, using $E_M 14150\text{ M}^{-1}$ at 412 nm [36]. Reduction potential was calculated by using the Nernst equation for $\text{GSSG}/2\text{GSH}$: $E_{\text{GSH}} = E_0 - 30 \times \log ([\text{GSH}]^2 / [\text{GSSG}])$ [27].

2.6. Statistical analysis

Each experiment was performed at least three times, and the results were expressed as mean values \pm S.D. ($n=2\text{--}6$).

The Student *t* test was used to determine the significance of differences between the mean values obtained for the two cell lines. Otherwise, the various treatments, were analyzed using one-way analysis of variance (ANOVA) followed by either Dunnett or Bonferroni's multiple comparison test, considering $P < 0.05$ as significant.

3. Results

3.1. Allicin inhibited proliferation of HL60 and U937 cells

The proliferation rate of exponentially growing HL60 and U937 cells in the presence of allicin was determined at a density of $100,000\text{ cells ml}^{-1}$. Cells were incubated with increasing concentrations of allicin (HL60: $0\text{--}10\text{ }\mu\text{M}$; U937: $0\text{--}30\text{ }\mu\text{M}$) for growing time periods up to 72 h. Cell viability was determined by the trypan blue dye exclusion assay. Allicin inhibited cell growth in a concentration-dependent manner (Figs. 1A and 2A for HL60 and U937, respectively). In the presence of $5\text{ }\mu\text{M}$ allicin, HL60 cells exhibited about 50% inhibition of proliferation after 22 h, and at $10\text{ }\mu\text{M}$ allicin after 22 h, the rate of inhibition reached 80% (Fig. 1A). HL60 cells were more sensitive to allicin than U937 cells. Inhibition of U937 cell proliferation reached 50% and 60% at 15 and $20\text{ }\mu\text{M}$ allicin, respectively, after 22 h (Fig. 2A). We therefore chose to continue the study of HL60 and U937 cells ($100,000\text{ cells/ml}$) at 5 and $20\text{ }\mu\text{M}$ allicin, respectively.

3.2. Allicin induced apoptosis in the cells

3.2.1. Morphological changes

Morphological changes indicating apoptotic processes were observed by staining the cells with Hoechst 33342. Distinct nuclear condensation was observed in HL60 cells treated with allicin ($5\text{ }\mu\text{M}$) for 16 h (Fig. 1B) and U937 cells treated with allicin ($20\text{ }\mu\text{M}$) for 40 h (Fig. 2B). In both allicin-treated cell lines, more than 90% of the cells showed blebbing after 2 h (Fig. 3A, lower panel).

3.2.2. DNA fragmentation

A DNA ladder appeared in allicin ($5\text{ }\mu\text{M}$)-treated HL60 cells 6 h after treatment, whereas in allicin ($20\text{ }\mu\text{M}$)-treated U937 cells, it appeared only after 40 h (Figs. 1C and 2C).

3.2.3. Cytochrome *c* release into cytoplasm and activation of caspases

To better understand the mechanism of allicin-induced apoptosis, Western blot analysis was used to detect apoptosis-related proteins. Increased amounts of cytochrome *c* appeared in the cytosol of allicin ($5\text{ }\mu\text{M}$)-treated HL60 cells after 2.5 h of incubation. An increase in cleaved caspase 9 appeared 4 and 6 h after adding allicin, and it gradually decreased to the basal level after 30 h. Cleaved caspase 3 appeared after 16 h and was still observed 30 h after the treatment (Fig. 3B). U937 cells reacted at a slower rate to allicin ($20\text{ }\mu\text{M}$) treatment. Cytochrome *c* release into the cytosol began 6 h after adding allicin and reached a maximal




Fig. 1. Effect of allicin on cell growth of human leukemia HL60 cells. (A) Growth dependent of HL60 cells on allicin concentration. Viable cells were counted at various time intervals between 0–72 h after staining with trypan blue. Data represent the mean value of % viable cells (nontreated=100%) \pm S.D. (B) Apoptotic changes observed after Hoechst 33342 (5 μ g/ml) staining for 30 min. Cells were treated with 5 μ M allicin for 16 h. (C) Nucleosomal DNA fragmentation of nontreated cells and allicin-treated HL60 cells (5 μ M, 6 h, 5 μ g/lane). Lane M is a DNA size marker.




Fig. 2. Effect of allicin on cell growth of human leukemia U937 cells. (A) Growth dependence of U937 cells on allicin concentration. Viable cells were counted at various time intervals between 0–72 h after staining with trypan blue. Data represent the mean value of % viable cells (nontreated=100%) \pm S.D. (B) Hoechst 33342 (5 μ g/ml) staining for 30 min of cells treated with 20 μ M allicin for 40 h. (C) Nucleosomal DNA fragmentation of nontreated cells and allicin-treated U937 cells (20 μ M, 40 h, 10 μ g/lane). Lane M is a DNA size marker.




Fig. 3. (A) Morphological changes in HL60 and U937 cells treated with allixin. HL60 and U937 cells treated with allixin (5 or 20 μ M, respectively) for 2 h were collected and washed with PBS. Cells were visualized by confocal microscopy (original magnification $\times 40$). (Upper panel) Nontreated cells (the diameter of 100 cells was measured). (Lower panel) Allixin-treated cells. (B) Immunoblot analysis of extracts (20 μ g protein/lane) from allixin treated HL60 cells at various time of allixin (5 μ M) treatment. The degrees of cytochrome *c* release into the cytosol and of caspase 9 and caspase 3 activation were assessed. β -Actin served as a control to the amount of loaded protein. The analysis of caspase 3, 9 and β -actin was done on the same blot after stripping. The data were obtained from one representative experiment (total of three independent experiments). (C) Staining of HL60 cells with MitoTracker Red CMXRos. HL60 cells treated with allixin (5 μ M, 6 h) were collected, resuspended in fresh medium and were stained for 30 min with MitoTracker Red CMXRos (0.05 μ M, 30 min). After washing twice with PBS, cells were examined by fluorescence microscopy (original magnification $\times 40$).

level after 18 h. The band of cleaved caspase 9 appeared at 24 h, reached a maximal intensity at 48 h and was still seen at 72 h. Cleaved caspase 3 appeared at 30 h, became intense at 48 h and did not fade after 72 h (data not shown). Cleaved caspase 8 was not observed in either cell line after allixin treatment (data not shown).

3.2.4. Impaired mitochondrial activity

Mitochondrial activity was monitored with the XTT assay in cells treated with allixin (5 and 20 μ M for HL60 and U937, respectively) for 16 h (Fig. 4). Both cell lines behaved similarly, showing a residual mitochondrial activity $65 \pm 5\%$ in HL60 and $62 \pm 5\%$ in U937 cells. At this stage, the drop in




Fig. 4. (A) Allicin effect on cell viability in the absence or presence of *N*-acetyl cysteine. HL60 and U937 cells (100,000 cells/ml) were treated with allicin (5 or 20 μ M, respectively) for 16 h, in the absence or presence of *N*-acetyl cysteine at 0.1, 0.5 and 1.0 mM. Cell viability was measured by the XTT assay after 3 h of incubation at 37°C with 5% CO₂. Means \pm S.D. of one experiment ($n=6$) are shown (total of three independent experiments). ** $P<.01$ represents a significant difference from nontreated cells, based on one-way ANOVA followed by Dunnett's test. (B) Allicin effect on HL60 cell viability following a 1-h CsA (5 μ M) pretreatment. HL60 cells (100,000 cells/ml) were treated with or without allicin (5 μ M) for 16 h. Cell viability was measured by the XTT assay. Means \pm S.D. of one experiment ($n=6$) are shown (total of three independent experiments). *** $P<.001$, represents significant difference compared with nontreated cells by one-way ANOVA followed by Bonferroni's multiple comparison test.

mitochondrial activity was not accompanied by a significant change in cell number. Such a change was observed only after 24 h. Nontreated HL60 cells proliferated at a higher rate than nontreated U937 cells. Although after 24 h nontreated HL60 cells exceeded nontreated U937 cells in number, mitochondrial activity (measured by XTT assay) in U937 cells (0.869 ± 0.025 optical density [OD] units at 450 nm, $n=6$) was significantly higher than that in HL60 cells (0.498 ± 0.073 OD units, $n=6$) ($P<.001$). These results indicate a higher battery of mitochondrial enzymes in U937 cells and

suggest a higher resistance to allicin. When cells were treated with allicin in the presence of NAC (0.1–1.0 mM), mitochondrial metabolic activity was not impaired, and the antiproliferative effect of allicin was precluded (Fig. 4A).

3.3. Allicin induces changes in mitochondrial membrane potential

Nontreated cells showed strong staining with MitoTracker Red CMXRos, indicating the negative membrane potential of the mitochondria and the accumulation of the lipophilic dye. In HL60 cells treated with allicin (5 μ M, 6 h the staining intensity decreased significantly, indicating changes in the mitochondrial membrane potential (Fig. 3C). Reduced mitochondrial staining in allicin-treated (20 μ M) U937 cells occurred much slower; changes were observed 24 h after treatment, and the decline continued up to 40 h (data not shown).

A deeper insight into the involvement of mitochondrial permeability transition pore (mPTP) in the cytotoxicity of allicin was gained by using cyclosporine A (CsA), an inhibitor of adenine nucleotide translocator. HL60 cells were preincubated with 5 μ M CsA for 1 h and then treated with allicin for 16 or 20 h. CsA inhibited the detrimental effects of allicin on cell viability as determined by the XTT assay ($\sim 80\%$ residual viability for CsA pretreatment; $\sim 50\%$ for allicin alone). There was no significant difference between nontreated and CsA-treated cells ($P>.05$). No significant difference was found between CsA alone and CsA followed by allicin treated cells ($P>.05$) (Fig. 4B). CsA-pretreated cells were stained with Hoechst 33342 and MitoTracker Red CMXRos. CsA pretreatment prevented the nuclear condensation induced by treatment with allicin alone (Fig. 5). Mitochondrial staining also showed the preventive effect of CsA, whereby allicin-induced mitochondrial membrane depolarization was inhibited.

Fluorescence-activated cell sorting (FACS) analysis of HL60 cells treated with allicin (5 μ M, 20 h) showed that 35.0% of the cells were apoptotic, whereas CsA pretreatment reduced the apoptotic fraction to 17.1% ($P<.05$). Cell arrest at G2 was increased in allicin-treated cells with or without CsA pretreatment 34.9% and 59.0%, respectively (Fig. 6). The fraction of HL60 cells at the G2 stage was 17.3% for nontreated cells and 18.9% for cells treated with CsA only (the difference is not significant).

3.4. Differences in GSH levels between HL60 and U937

GSH concentrations (nmol/total cell volume) in nontreated HL60 and U937 cells were almost the same: 3.4 ± 0.48 mM and 3.7 ± 0.55 mM, respectively. However, due to the differences in cell size between the two cell lines, analysis of GSH concentration expressed as a function of protein levels or cell numbers was also used. This analysis revealed significant differences between the GSH concentrations in the two cell lines. In nontreated HL60 cells, GSH concentration was 20.94 ± 3.15 nmol/mg protein, and in nontreated U937 cells, 30.80 ± 6.04 nmol/mg protein ($P<.02$). The

HL60




Fig. 5. Morphological effects of allicin on CsA pretreated HL60 cells. Cells (200,000 cells/ml) with or without CsA pretreatment (5 μ M, 1 h) were diluted and incubated in the presence or absence of allicin (5 μ M) for 16 h. Cells were stained with Hoechst 33342 (5 μ g/ml) or with Mito-Tracker Red CMXRos (50 nM) for 30 min. Cells were washed twice with PBS and examined by fluorescence microscopy. (A) HL60 nontreated. (B) CsA-treated. (C) Allicin-treated. (D) CsA- and allicin-treated.

difference was found to be even higher when cell numbers were considered, GSH concentration in HL60 cells was 3.7 ± 0.56 nmol/ 10^6 cells whereas, in U937 cells, approximately twice as much [7.7 ± 1.08 nmol/ 10^6 cells ($P < .001$)].

3.5. Allicin-induced GSH depletion

Allicin treatment of HL60 cells caused a rapid decrease in GSH concentration, reaching its lowest point at about 30 min. A spontaneous recovery occurred, reaching the basal level of GSH (Fig. 7A). The highest GSH levels were observed in HL60 cells between 2–24 h after adding allicin. In allicin-treated U937 cells, a decrease in GSH levels to a minimal value occurred 2–4 h after treatment, and it was followed by a gradual increase to the basal GSH level (Fig. 7B).

3.6. Allicin-induced GSH depletion in the cells pretreated with BSO

In HL60 cells treated with BSO, an inhibitor of γ -glutamyl-cysteine synthetase (GGCS) (0.4 mM, 20 h) GSH levels decreased to 1.45 ± 0.25 nmol/ 10^6 cells. No cell death was observed, but there was a slight decrease in the proliferation rate ($\times 1.7$ after 20 h as compared to $\times 2.1$ in




Fig. 6. Cell cycle analysis of allicin-treated HL60 cells. Cells (200,000 cells/ml, 1.5×10^6) with or without CsA pretreatment (5 μ M, 1 h) were diluted and incubated in absence or presence of allicin (5 μ M) for 20 h. The cells were stained with propidium iodide in the presence of RNase A and Triton X-100, as described in Materials and methods. The data represent one experiment (total of three independent experiments).




Fig. 7. Time-dependent decrease of GSH levels in alliin-treated HL60 and U937 cells after BSO pretreatment. Cells were pretreated with or without 0.4 mM BSO for 20 h. HL60 and U937 cell cultures were diluted to 100,000 cells/ml and were grown in the presence of alliin (5 and 20 μ M, respectively). At each time point, 30-ml samples were harvested and washed twice with PBS. Cell pellets were extracted with 5% metaphosphoric acid and analyzed for GSH content and protein concentration. Means \pm S.D. of one representative experiment (total of three independent experiments).

nontreated cells). Alliin (5 μ M) applied to HL60 cells pretreated with BSO (0.4 mM, 20 h) caused an additional GSH decrease to 0.65 ± 0.13 nmol/10⁶ cells after 1 h (Fig. 7A, black column). There was no recovery of GSH as in the case of HL60 cells treated with alliin alone. In U937 cells pretreated with BSO (0.4 mM, 20 h), GSH levels decreased to 1.06 ± 0.14 nmol/10⁶ cells. Also, in this case, no cell death was observed and cells were still proliferating. When BSO-pretreated U937 cells were incubated with alliin (20 μ M), GSH levels further decreased to 0.33 ± 0.08 nmol/10⁶ cells within 2 h (Fig. 7B, black column). A high rate of cell death (60–80%) occurred in both cell lines 24 h after alliin was applied to the culture.

3.7. Reduction potential in alliin-treated cells

Since alliin-treated cells lost their spherical structure and their volume could not be calculated from diameter measurement, we had to use another parameter to calculate

changes in the intracellular reducing state resulting from fluctuations in 2GSH/GSSG ratios. Based on the assumption that 1-mg cell pellet is equivalent to 1 μ l of volume, cell pellets were weighed, and GSH molar concentration was calculated accordingly (GSH amount/cell weight). Therefore, calculated reduction potential represents only approximate values. The intracellular pH of HL60 cells is 7.0 [37], and the reported E_0 for this pH is –240 mV [27]. Using the above E_0 values, reducing potential (E_{GSH}) calculated for nontreated HL60 was –207 mV (GSH 3.42 ± 0.18 mM, GSSG 0.15 ± 0.023 mM) and for alliin-treated (0.5 h) –169 mV (GSH 1.4 ± 0.17 mM, GSSG 0.46 ± 0.082 mM). The intracellular pH of U937 cells is 7.04 [38] or 7.2 [39], and the




Fig. 8. Free SH contents in cytosolic and crude mitochondrial fractions from alliin-treated HL60 cells. Alliin (5 μ M) was added to cells growing at density of 100,000 cells/ml. At various intervals, 100-ml samples were harvested at 4°C, washed twice with PBS and stored at –80°C overnight. Cell fractionation was done as described in Materials and methods. Total free SH and protein concentrations were determined as described in Materials and methods. The SH content at various time intervals in the cytosol (upper panel) and in the crude mitochondrial fraction (lower panel) are shown. Means \pm S.D. of one representative experiment (total of three independent experiments) [40].

reported E_0 values are -240 mV or -252 mV, respectively. The calculated reducing potential values of nontreated U937 cells were -206 mV or -218 mV (GSH 3.7 ± 0.21 mM, GSSG 0.18 ± 0.040 mM), and for allicin-treated cells (2 h), -173 mV or -185 mV (GSH 1.54 ± 0.042 mM, GSSG 0.41 ± 0.021 mM). The values obtained for nontreated cells are characteristic of cells in proliferation/differentiation state, whereas those obtained after allicin treatment are typically obtained in the apoptotic state.

3.8. Free SH in the cytosolic and mitochondrial fractions of HL60 cells

Total free SH concentrations were determined in the cytosol and in the crude mitochondrial fractions prepared from allicin-treated HL60 cells. The decrease in free SH in the cytosol (Fig. 8, upper panel) was accompanied by a decrease of free SH in the mitochondria (Fig. 8, lower panel). Since mitochondrial GSH is supplied from the cytoplasm and upon allicin treatment it was depleted in both cellular compartments, it was followed by a substantial decrease in the cell-reducing potential. Consequently, depolarization of the mitochondrial membrane occurred and triggered a chain of apoptotic events leading to cell death.

4. Discussion

GSH, cysteine and accessible free SH groups of various proteins and peptides are rapidly oxidized by allicin to their respective allyl-SS derivatives [19,22]. The mixed disulfide formed is less active than allicin, yet it can easily react with free SH in thiol-disulfide exchange reactions, and produce, at the end of a series of reactions, the oxidized form of glutathione (GSSG), the oxidized form of cysteine (cystine) and glutathiolated S proteins. The leaving group (allyl-mercaptan) is volatile and rapidly disappears.

Glutathione is the major regulator of the thiol-disulfide redox state of living cells (reviewed in Refs. [29,41–43]). Alterations in cellular free thiols have already been described as a key event in the generation of apoptosis (reviewed in [44,45]).

In this study, allicin induced mitochondrial release of cytochrome *c*, activation of caspase 9 and 3 and DNA fragmentation in HL60 and U937 cell lines in a concentration- and time-dependent manner. GSH depletion had been shown to be associated with cell surface blebbing in hepatocytes [46]. Here, cell blebbing occurred in both HL60 and U937 cells treated with allicin. The rate of free SH depletion from the cytosol and the mitochondria was similar, indicating that allicin penetrated the mitochondria and the cell membrane equally well and reacted with free SH in both compartments. Allicin induced an initial drop in GSH levels that was followed by a regeneration process, occurring several hours after treatment. This may be due to decreased feedback inhibition of GSH biosynthetic enzymes such as GGCS by GSH [45]. Conversely, in BSO-pretreated cells,

there was no recovery of GSH after its allicin-induced depletion, as BSO is a direct inhibitor of GGCS. BSO pretreatment in allicin-treated U937 cells shortened the life span from 72 to 20 h, indicating that preliminary GSH depletion renders cells more sensitive to apoptosis. BSO-depleted GSH levels in U937 cells were, at this stage, similar to the inherent GSH level in HL60 cells and, hence, a similar degree of sensitivity to allicin.

Here, we have shown that allicin causes GSH depletion concomitantly in the cytosol and the mitochondria in HL60 cells. It is known that active GSH uptake by intact mitochondria occurs even at very low cytosolic GSH levels and that the rate of mitochondrial GSH depletion is much slower than that of the cytosol [47]. These are indications that mitochondria have more than one control mechanism to maintain GSH levels. Since BSO-induced depletion of GSH levels caused neither mitochondrial permeability transition nor apoptosis, it is likely that BSO treatment depletes GSH mainly from the cytosol, whereas allicin induced depletion of mitochondrial GSH trigger the apoptotic process. Moreover, CsA (mPTP inhibitor) pretreatment of HL60 cells inhibits allicin-induced mitochondrial permeability transition and cell death, corroborating the central role of mitochondria in the apoptotic process induced by allicin. Simultaneous addition of NAC and allicin to cells provided protection against the apoptotic damage. NAC sequestered the harmful activity by competing with GSH for the reaction with allicin.

The following allicin-induced apoptotic scenario can be deduced from this study: Allicin penetrates the cells and readily reacts with any exposed SH present in its vicinity. As GSH is the most abundant thiol molecule in the cytosol and the mitochondria, it is the main target for allicin reaction. The reaction with glutathione leads to an immediate change in the ratio GSH/GSSG. The increased concentration of GSSG and the depletion of GSH brings about a decrease in the cellular reduction potential state. Concomitantly, mitochondrial damage occurs, as is evident from changes in membrane potential, its decreased enzymatic activity and the release of cytochrome *c* to the cytosol. Upon its release, cytochrome *c* participates in the activation of the caspase 9, followed by activation of caspase 3.

Allicin induced cell cycle arrest at G2/M in HL60 cells ($\sim 35\%$). Upon CsA pretreatment, the percentage of cells at G2/M was even higher ($\sim 59\%$). After pretreatment of HL60 cells with CsA and 40-h incubation with allicin, there was no change in the number of viable cells, neither were there signs of cell breakage. By using CsA, it could be shown that the effect of allicin on cell cycle arrest was independent of its proapoptotic activity. The mechanism by which allicin induces G2/M cell arrest is not clear, but a clue may reside in its ability to inhibit microtubule assembly. It had been shown that microtubule assembly is disrupted by direct modification of tubulin SH groups [48–50] and that adding GSH to an *in vitro* microtubule assembly system disrupted by SH modification restores tubulin polymerization [51].

Here we have shown that, in allicin-treated cells, there is an increase in GSH synthesis. Therefore, cell cycle arrest would be expected to expire after tubulin recovery. However, the protracted G2/M phase induced by cyclosporine A and allicin suggests an additional modification of a critical pathway that needs an extended period to regenerate.

Allicin disappears from the medium very rapidly due to its penetration into the cells and its reaction with SH group. When applied to HL60 cells ($5 \mu\text{M}/10^5$ cells/ml) or to U937 cells ($20 \mu\text{M}/10^5$ cells/ml), it could not be detected in the medium after 2 h or 6–8 h in HL60 or U937 cells, respectively. The consumption of allicin within such a short time enables full activity of SH-dependent apoptotic enzymes such as the cysteine proteases and caspases 9 and 3 [52].

Cell sensitivity to allicin was shown to be dependent on the intracellular content of GSH. Since there is a significant difference in cell size between the two cell lines examined, comparing GSH amount/cell number showed that HL60 cells contained $3.7 \text{ nmol GSH}/10^6$ cells, whereas U937 cells contained $7.7 \text{ nmol GSH}/10^6$ cells. The differences in GSH intracellular content reflect differences in sensitivity to allicin. The higher the GSH, the less sensitive to allicin cells are, and the slower the apoptotic signs appear.

The allicin-induced decrease in GSH levels modulates the redox state of cells. In allicin-treated HL60 cells, the ratio of [GSH]/[GSSG or GSSA] changed within the first 30 min and the reducing potential reached -170 mV , whereas in U937 cells, it took 2 h to reach the same level. Such a high value is typical of the apoptotic state [27].

Since GSSA (the first main product of allicin reaction with GSH) is able to prevent the formation of free radicals [19], it is likely that the initial event inducing apoptosis does not occur via the generation of reactive oxygen species but rather through mitochondrial membrane depolarization or through a more finely tuned event mediated by protein modification upon creating a mixed disulfide [19,53]. Oxidative stress incurred by the accumulation of peroxides is a secondary event probably happening after the short-lived GSSA is eliminated from the cells.

It was reported that, in BSO-treated U937 cells, the depletion of GSH and the release of cytochrome *c* into the cytosol were counteracted by cell-repair systems involving nuclear factor κB and heat shock proteins [54]. Our results show that CsA and NAC prevented apoptosis induced by allicin. Taken together, the role allicin plays may be through direct modification of specific free SH in proteins, such as reside in apoptotic repair system, in the mPTP [55] or others.

Acknowledgment

We would like to thank Dr. A. Gross, Department of Biological Regulation, Weizmann Institute of Science for critical reading of the manuscript, and Mrs. Anna Gakamsky for her help in the statistical analysis.

References

- [1] Stoll A, Seebeck E. Chemical investigation on alliin, the specific principle of garlic. *Adv Enzymol* 1951;11:377–400.
- [2] Koch HP, Lawson LD. Garlic: the Science and therapeutic application of *Allium sativum* L. and related species. 2nd ed. Baltimore: Williams & Wilkins; 1996.
- [3] Elkayam A, Mirelman D, Peleg E, Wilchek M, Miron T, Rabinkov A, et al. The effects of allicin and enalapril in fructose-induced hyperinsulinemic hyperlipidemic hypertensive rats. *Am J Hypertens* 2001;14:377–81.
- [4] Eilat S, Oestraicher Y, Rabinkov A, Ohad D, Mirelman D, Battler A, et al. Alteration of lipid profile in hyperlipidemic rabbits by allicin, an active constituent of garlic. *Coron Artery Dis* 1995;6:985–90.
- [5] Abramovitz D, Gavri S, Harats D, Levkovitz H, Mirelman D, Miron T, et al. Allicin-induced decrease in formation of fatty streaks (atherosclerosis) in mice fed a cholesterol-rich diet. *Coron Artery Dis* 1999;10:515–9.
- [6] Gonen A, Harats D, Rabinkov A, Miron T, Mirelman D, Wilchek M, et al. The antiatherogenic effect of allicin: possible mode of action. *Pathobiology* 2005;72:325–34.
- [7] Lang A, Lahav M, Sakhnini E, Barshack I, Fidler HH, Avidan B, et al. Allicin inhibits spontaneous and TNF-alpha induced secretion of proinflammatory cytokines and chemokines from intestinal epithelial cells. *Clin Nutr* 2004;23:1199–208.
- [8] Agarwal KC. Therapeutic actions of garlic constituents. *Med Res Rev* 1996;16:111–24.
- [9] Hirsch K, Danilenko M, Giat J, Miron T, Rabinkov A, Wilchek M, et al. Effect of purified allicin, the major ingredient of freshly crushed garlic, on cancer cell proliferation. *Nutr Cancer* 2000;38:245–54.
- [10] Knowles LM, Milner JA. Possible mechanism by which allyl sulfides suppress neoplastic cell proliferation. *J Nutr* 2001;131:1061S–6S.
- [11] Sun L, Wang X. Effects of allicin on both telomerase activity and apoptosis in gastric cancer SGC-7901 cells. *World J Gastroenterol* 2003;9:1930–4.
- [12] Oommen S, Anto RJ, Srinivas G, Karunakaran D. Allicin (from garlic) induces caspase-mediated apoptosis in cancer cells. *Eur J Pharmacol* 2004;485:97–103.
- [13] Park SY, Cho SJ, Kwon HC, Lee KR, Rhee DK, Pyo S. Caspase-independent cell death by allicin in human epithelial carcinoma cells: involvement of PKA. *Cancer Lett* 2005;224:123–32.
- [14] Antlisperger DSM, Dirsch VM, Ferreira D, Su JL, Kuo ML, Vollmar AM. Ajoene-induced cell death in human promyeloleukemic cells does not require JNK but is amplified by the inhibition of ERK. *Oncogene* 2003;22:582–9.
- [15] Jin Z, El-Deiry WS. Overview of cell death signaling pathways. *Cancer Biol Ther* 2005;4:139–63.
- [16] Hsu YT, Wolter KG, Youle RJ. Cytosol-to-membrane redistribution of Bax and Bcl-X1 during apoptosis. *Proc Natl Acad Sci U S A* 1997;94:3668–72.
- [17] Wolter KG, Hsu YT, Smith A, Nechushtan A, Xi X, Youle RJ. Movement of Bax from the cytosol to mitochondria during apoptosis. *J Cell Biol* 1997;139:1281–92.
- [18] Miron T, Rabinkov A, Mirelman D, Wilchek M, Weiner L. The mode of action of allicin: its ready permeability through phospholipid membranes may contribute to its biological activity. *Biochim Biophys Acta* 2000;1463:20–30.
- [19] Rabinkov A, Miron T, Mirelman D, Wilchek M, Glozman S, Yavin E, et al. S-allylmercaptogluthathione: the reaction product of allicin with glutathione possesses SH-modifying and antioxidant properties. *Biochim Biophys Acta* 2000;1499:144–53.
- [20] Xiao H, Parkin KL. Antioxidant functions of selected allium thiosulfinates and S-alk(en)yl-L-cysteine sulfoxides. *J Agric Food Chem* 2002;50:2488–93.
- [21] Okada Y, Tanaka K, Sato E, Okajima H. Kinetic and mechanistic studies of allicin as an antioxidant. *Org Biomol Chem* 2006;4:4113–7.
- [22] Rabinkov A, Miron T, Konstantinovski L, Wilchek M, Mirelman D, Weiner L. The mode of action of allicin: trapping of radicals and

- interaction with thiol containing proteins. *Biochim Biophys Acta* 1998;1379:233–44.
- [23] Freeman F, Koder Y. Garlic chemistry-stability of *S*-(2-propenyl) 2-propene-1-sulfinothioate (allicin) in blood, solvents, and simulated physiological fluids. *J Agric Food Chem* 1995;43:2332–8.
- [24] Lachmann G, Lorenz D, Radeck W, Steiper M. The pharmacokinetics of the S35 labeled garlic constituents alliin, allicin and vinylthiine. *Arzneimittelforschung* 1994;44:734–43.
- [25] Miron T, Mironchik M, Mirelman D, Wilchek M, Rabinkov A. Inhibition of tumor growth by a novel approach: in situ allicin generation using targeted alliinase delivery. *Mol Cancer Ther* 2003;2:1295–301.
- [26] Arditti F, Rabinkov A, Miron T, Reisner Y, Berrebi A, Wilchek M, et al. Apoptotic killing of B-chronic lymphocytic leukemia tumor cells by allicin generated in situ using a rituximab–alliinase conjugate. *Mol Cancer Ther* 2005;4:325–31.
- [27] Schafer FQ, Buettner GR. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med* 2001;30:1191–212.
- [28] Jonas CR, Ziegler TR, Gu LH, Jones DP. Extracellular thiol/disulfide redox state affects proliferation rate in a human colon carcinoma (Caco2) cell line. *Free Radic Biol Med* 2002;33:1499–506.
- [29] Gilbert HF. Molecular and cellular aspects of thiol-disulfide exchange. *Adv Enzymol Relat Areas Mol Biol* 1990;63:69–172.
- [30] Stoll A, Seebeck E. Allium compounds V. The synthesis of natural alliin and its three optical active isomers. *Helv Chim Acta* 1951;34:481–7.
- [31] Miron T, SivaRaman H, Rabinkov A, Mirelman D, Wilchek M. A method for continuous production of allicin using immobilized alliinase. *Anal Biochem* 2006;351:152–4.
- [32] Nakagawa Y, Iinuma M, Matsuura N, Yi K, Naoi M, Nakayama T, et al. A potent apoptosis-inducing activity of a sesquiterpene lactone, arucanolide, in HL60 cells: a crucial role of apoptosis-inducing factor. *J Pharmacol Sci* 2005;97:242–52.
- [33] Grinberg M, Schwarz M, Zaltsman Y, Eini T, Niv H, Pietrokovski S, et al. Mitochondrial carrier homolog 2 is a target of tBID in cells signaled to die by tumor necrosis factor alpha. *Mol Cell Biol* 2005;25:4579–4590.
- [34] Gornall A, Bardawill C, David M. Determination of serum proteins by means of the biuret reaction. *J Biol Chem* 1949;177:751–66.
- [35] Anderson ME. Determination of glutathione and glutathione disulfide in biological samples. *Methods Enzymol* 1985;113:548–55.
- [36] Riddles PW, Blakeley RL, Zerner B. Ellman's reagent: 5,5'-dithiobis (2-nitrobenzoic acid) — a reexamination. *Anal Biochem* 1979;94:75–81.
- [37] Restrepo D, Kozody DJ, Spinelli LJ, Knauf PA. pH Homeostasis in promyelocytic leukemic HL60 cells. *J Gen Physiol* 1988;92:489–507.
- [38] Oehler R, Zellner M, Hefel B, Weingartmann G, Spittler A, Struse HM, et al. Influence of heat shock on cell volume regulation: protection from hypertonic challenge in a human monocyte cell line. *FASEB J* 1998;12:553–60.
- [39] Nilsson C, Kagedal K, Johansson U, Ollinger K. Analysis of cytosolic and lysosomal pH in apoptotic cells by flow cytometry. *Methods Cell Sci* 2003;25:185–94.
- [40] Amonkar SV, Reeves EL. Mosquito control with active principle of garlic, *Allium sativum*. *J Econ Entomol* 1970;63:1172–5.
- [41] Powis G. Anticancer drugs acting against signaling pathways. *Curr Opin Oncol* 1995;7:554–9.
- [42] Sies H. Glutathione and its role in cellular functions. *Free Radic Biol Med* 1999;27:16–21.
- [43] Moran LK, Gutteridge JM, Quinlan GJ. Thiols in cellular redox signalling and control. *Curr Med Chem* 2001;8:763–72.
- [44] Meister A. Glutathione metabolism. *Methods Enzymol* 1995;251:3–7.
- [45] Lash LH. Mitochondrial glutathione transport: physiological, pathological and toxicological implications. *Chem Biol Interact* 2006;163:54–67.
- [46] Jewell SA, Bellomo G, Thor H, Orrenius S, Smith M. Bleb formation in hepatocytes during drug metabolism is caused by disturbances in thiol and calcium ion homeostasis. *Science* 1982;217:1257–9.
- [47] Meister A. Mitochondrial changes associated with glutathione deficiency. *Biochim Biophys Acta* 1995;1271:35–42.
- [48] Kuriyama R, Sakai H. Role of tubulin-SH groups in polymerization to microtubules. Functional-SH groups in tubulin for polymerization. *J Biochem (Tokyo)* 1974;76:651–4.
- [49] Li M, Ciu JR, Ye Y, Min JM, Zhang LH, Wang K, et al. Antitumor activity of Z-ajoene, a natural compound purified from garlic: antimitotic and microtubule-interaction properties. *Carcinogenesis* 2002;23:573–9.
- [50] Xiao D, Pinto JT, Soh JW, Deguchi A, Gundersen GG, Palazzo AF, et al. Induction of apoptosis by the garlic-derived compound *S*-allylmercaptocysteine (SAMC) is associated with microtubule depolymerization and c-Jun NH(2)-terminal kinase 1 activation. *Cancer Res* 2003;63:6825–37.
- [51] Nishida E, Kobayashi T. Relationship between tubulin SH groups and bound guanine nucleotides. *J Biochem (Tokyo)* 1977;81:343–7.
- [52] Thornberry NA. Caspases: key mediators of apoptosis. *Chem Biol* 1998;5:R97–R103.
- [53] Patya M, Zahalka MA, Vanichkin A, Rabinkov A, Miron T, Mirelman D, et al. Allicin stimulates lymphocytes and elicits an antitumor effect: a possible role of p21(ras). *Int Immunol* 2004;16:275–81.
- [54] Filomeni G, Aquilano K, Rotilio G, Ciriolo MR. Antiapoptotic response to induced GSH depletion: involvement of heat shock proteins and NF-kappaB activation. *Antioxid Redox Signal* 2005;7:446–55.
- [55] McStay GP, Clarke SJ, Halestrap AP. Role of critical thiol groups on the matrix surface of the adenine nucleotide translocase in the mechanism of the mitochondrial permeability transition pore. *Biochem J* 2002;367:548.