

# Zebrafish (*Danio rerio*) fed vitamin E-deficient diets produce embryos with increased morphologic abnormalities and mortality<sup>☆</sup>

Galen W. Miller<sup>a,b</sup>, Edwin M. Labut<sup>a</sup>, Katie M. Lebold<sup>a</sup>, Abby Floeter<sup>a</sup>,  
Robert L. Tanguay<sup>b,c,e</sup>, Maret G. Traber<sup>a,b,d,e,\*</sup>

<sup>a</sup>The Linus Pauling Institute, Oregon State University, Corvallis, OR 97331, USA

<sup>b</sup>Molecular and Cellular Biology Program, Oregon State University, Corvallis, OR 97331, USA

<sup>c</sup>Department of Environmental and Molecular Toxicology, Oregon State University, Corvallis, OR 97331, USA

<sup>d</sup>Department of Nutrition and Exercise Sciences, Oregon State University, Corvallis, OR 97331, USA

<sup>e</sup>Environmental Health Sciences Center, Oregon State University, Corvallis, OR 97331, USA

Received 15 May 2010; received in revised form 4 January 2011; accepted 3 February 2011

## Abstract

Vitamin E ( $\alpha$ -tocopherol) is required to prevent fetal resorption in rodents. To study  $\alpha$ -tocopherol's role in fetal development, a nonplacental model is required. Therefore, the zebrafish, an established developmental model organism, was studied by feeding the fish a defined diet with or without added  $\alpha$ -tocopherol. Zebrafish (age, 4–6 weeks) were fed the deficient (E<sup>−</sup>), sufficient (E<sup>+</sup>) or lab diet up to 1 years. All groups showed similar growth rates. The exponential rate of  $\alpha$ -tocopherol depletion up to ~80 day in E<sup>−</sup> zebrafish was  $0.029 \pm 0.006$  nmol/g, equivalent to a depletion half-life of  $25 \pm 5$  days. From age ~80 days, the E<sup>−</sup> fish ( $5 \pm 3$  nmol/g) contained ~50 times less  $\alpha$ -tocopherol than the E<sup>+</sup> or lab diet fish ( $369 \pm 131$  or  $362 \pm 107$ , respectively;  $P < .05$ ). E-depleted adults demonstrated decreased startle response suggesting neurologic deficits. Expression of selected oxidative stress and apoptosis genes from livers isolated from the zebrafish fed the three diets were evaluated by quantitative polymerase chain reaction and were not found to vary with vitamin E status. When E-depleted adults were spawned, they produced viable embryos with depleted  $\alpha$ -tocopherol concentrations. The E<sup>−</sup> embryos exhibited a higher mortality ( $P < .05$ ) at 24 h post-fertilization and a higher combination of malformations and mortality ( $P < .05$ ) at 120 h post-fertilization than embryos from parents fed E<sup>+</sup> or lab diets. This study documents for the first time that vitamin E is essential for normal zebrafish embryonic development.

© 2012 Elsevier Inc. All rights reserved.

**Keywords:**  $\alpha$ -Tocopherol;  $\gamma$ -Tocopherol; Embryonic development; Liver; Ascorbic acid; Morphology

**Abbreviations:** AIF (*pdc8*), apoptosis-inducing factor (*programmed cell death 8*);  $\beta$ 2M, beta-2-microglobulin; BA, bent anterior–posterior axis; BAD, BCL2-antagonist of cell death;  $\beta$ -actin, beta-actin; CF, cranial facial malformations; E<sup>−</sup> diet,  $\alpha$ -tocopherol-deficient diet; E<sup>+</sup> diet,  $\alpha$ -tocopherol-sufficient diet; FW, fish water; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; GCLC, glutamate-cysteine ligase, catalytic subunit; GPx4a, glutathione peroxidase 4a; GPx4b, glutathione peroxidase 4b; GSTp1, glutathione S-transferase pi; hpf, hours postfertilization; HPLC-ECD, high-pressure liquid chromatography coupled with electrochemical detection; lab diet, standard zebrafish diet (see “Methods”); LOX, arachidonate 12-lipoxygenase; NRF2, nuclear factor (erythroid-derived 2)-like 2; PE, pericardial edema; PLA2gVI, phospholipase A2, calcium dependent; PLA2gIV, phospholipase A2, calcium independent; SB, swim bladder malformation; TTP,  $\alpha$ -tocopherol transfer protein; YSE, yolk-sac edema

<sup>☆</sup> Grants: This publication was made possible, in part, by the Aquatic Biomedical Models Facility Core of the Environmental Health Sciences Center, Oregon State University, and National Institute of Environmental Health Sciences (P30 ES000210) and by a grant from The Eunice Kennedy Shriver National Institute Of Child Health and Human Development (HD062109).

\* Corresponding author. Tel.: +1 541 737 7977; fax: +1 541 737 5077.

E-mail address: [maret.traber@oregonstate.edu](mailto:maret.traber@oregonstate.edu) (M.G. Traber).

## 1. Introduction

Vitamin E ( $\alpha$ -tocopherol) was discovered in 1922 by Evans and Bishop [1], who demonstrated that rats fed diets containing rancid fat were unable to carry offspring to term. Both low fetal and maternal  $\alpha$ -tocopherol concentrations may be important factors during fetal resorption. Retention of uterine  $\alpha$ -tocopherol by the mother appears essential to maintain pregnancy because the  $\alpha$ -tocopherol transfer protein (TTP) is increased at the site of implantation [2,3] and is also expressed in the placenta [3–6]. Both TTP and biomarkers of lipid peroxidation (malondialdehyde and 4-hydroxy-2-nonenal) have been detected in the first-trimester placenta [6]. Moreover, early failure of pregnancy is associated with lipid peroxidation with resultant damage to the placental syncytiotrophoblast [7]. Thus, it is likely that  $\alpha$ -tocopherol is needed by the mother to protect her from the oxidative stress of the rapidly growing fetus.

TTP is a liver protein in adult rodents and humans. However, Jauniaux et al. [8] showed that not only does the human placenta and uterus express TTP, the human yolk sac expresses TTP as well. They further suggest that during very early human fetal development, the human embryo obtains  $\alpha$ -tocopherol from the yolk sac. Our

hypotheses are dependent on the observation that TTP is expressed during the first 48 h of zebrafish embryonic development [9].

The zebrafish is an established vertebrate model organism frequently used in developmental studies [10–13]. Embryos undergo maturation from a single cell to an autonomous juvenile fish in 120 h (5 days) [14]. During development, embryos are optically clear, allowing for noninvasive observation of the entire embryo. *TTP* messenger RNA (mRNA) is expressed in embryonic zebrafish as early as 24 h postfertilization (hpf) [9]. Because the zebrafish reproduces by spawning and does not have a placenta, the vitamin E requirements of the embryo can be separated from those of the mother. However, available commercial diets contain many ingredients making the study of a specific nutrient difficult [15]. Many “fish foods” also contain ambiguous ingredients, such as “fish oil” and/or “fish meal,” the sources of which are not immediately apparent. The feeding of live food sources (artemia, paramecium, etc.) to zebrafish present similar difficulties [16]. This problem has been addressed using zebrafish and other teleost fish in the past with varying levels of success by partially defining laboratory diets [17–20].

To address these challenges, we hypothesized that development of a defined diet, which is sufficient to sustain zebrafish health, growth and reproduction, will allow us to study the action of vitamin E during embryogenesis. We further hypothesize based on studies in rodent models of vitamin E deficiency that (1) we can deplete adult zebrafish of  $\alpha$ -tocopherol, (2) these adults will remain reproductively active, (3) the embryos that are produced will also be deficient in  $\alpha$ -tocopherol, and (4)  $\alpha$ -tocopherol-deficient embryos will show phenotypic abnormalities as a result of their deficiency.

## 2. Methods

### 2.1. Fish husbandry

Tropical 5D strain zebrafish (*Danio rerio*) were housed in the Sinnhuber Aquatic Research Laboratory at Oregon State University. The zebrafish were studied in accordance with protocols approved by the Institutional Animal Care and Use Committee.

Adult zebrafish were kept at standard laboratory conditions of 28°C on a 14-h light/10-h dark photoperiod in fish water (FW) consisting of reverse osmosis water supplemented with a commercially available salt solution (0.6% Instant Ocean; Spectrum Brands, Inc., Atlanta, GA, USA). Zebrafish were fed experimental diets (described in section 2.2) or a conventional lab diet comprising artemia (Inve Aquaculture Inc., Ogden, UT, USA) and a combination of commercial flake foods including: Aquatox Flake (77% by weight; Zeigler Brothers Inc., Gardners, PA, USA), Cyclopeez (6% by weight; Argent Laboratories, Redmond, WA, USA), Golden Pearls (8.5% by weight; Artemia International LLC, Fairview, TX, USA) and Hikari Micropellets (8.5% by weight; Hikari, Hayward, CA, USA), hereafter called “lab” diet. Zebrafish were fed twice daily (AM and PM) with an amount of food sufficient for the fish to consume in ~5 min. Embryos used for our studies were not fed, but instead received all their nutrition from their yolk sacs. At 120 hpf, embryos were euthanized by an overdose of tricaine. Note that zebrafish younger than 1 month cannot consume standard food because their mouths are too small and instead consume paramecium, artemia and specially created diets (EZ Larval Diet; Zeigler Brothers Inc., Gardners, PA, USA). Thus, we are currently unable to generate adult zebrafish that were spawned from deficient parents and have only consumed the vitamin E-deficient diet.

### 2.2. Diet preparation

Ingredients for the defined diets (Table 1) were obtained from Dyets Inc. (Bethlehem, PA, USA) with exceptions, as indicated: wheat gluten (#402100), casein (#400627), egg whites (#401600), cellulose (#401850), vitamin mix (#310064), mineral mix (#210087), tocopherol-stripped soybean oil (#404365), Stay C (Vitamin C-3; Argent Chemical Laboratories Inc.), modified food starch (National Starch Food Innovation, Bridgewater, NJ, USA), lecithin (Ultralec without added tocopherol) and vitamin E (RRR- $\alpha$ -tocopherol). Both the lecithin and the vitamin E were generous gifts from Archer Daniels Midland (Decatur, IL, USA).

Diets were prepared in 100-g batches without added vitamin E (E–) or with (E+; ~500 mg RRR- $\alpha$ -tocopherol/kg diet). Each diet was prepared by mixing ingredients until homogenous, spreading the batter on a large baking sheet and then oven drying at 212°F for 1 h. The diet was then cooled, crushed into a powder and stored frozen at –20°C until fed to the zebrafish. Diets were used within 4 months of preparation.

Table 1  
Diet ingredients

Ingredient	g/100 g
Wheat gluten	15
Casein	30.5
Egg whites	4
Cellulose	3
Vitamin mix <sup>a</sup>	4
Mineral mix <sup>b</sup>	4
Starch	26.5
Tocopherol-stripped soybean oil	7
Tocopherol-stripped soy lecithin	5
Stay C (Argent Chemical Laboratories)	1
$\alpha$ -Tocopherol (E+ diets only)	0.05

<sup>a</sup> Containing the following (g/kg vitamin mix): vitamin A (500,000 IU/g), 0.15; vitamin D<sub>3</sub> (400,000 United States Pharmacopeia (USP)/mg), 6.2445; vitamin K, 0.025; thiamine, 0.15; riboflavin, 0.25; vitamin B<sub>6</sub>, 0.125; pantothenic acid, 0.75; niacin, 1.25; biotin, 0.005; folate, 0.05; vitamin B<sub>12</sub>, 0.0005; myoinositol, 6.25; para-amino benzoic acid (PABA), 1; celufil ( $\alpha$ -cellulose), 983.75.

<sup>b</sup> Containing the following (g/kg mineral mix): calcium carbonate, 19.23; calcium phosphate dibasic (2H<sub>2</sub>O), 766.29; citric acid, 5.28; cupric carbonate, 0.36; ferric citrate, 2.99; magnesium oxide, 22.89; manganese carbonate, 5.65; sodium chloride, 28.02; disodium hydrogen phosphate, 11.89; zinc carbonate, 0.97; potassium phosphate dibasic, 74.16; potassium sulfate, 62.26; potassium iodide, 0.01.

### 2.3. $\alpha$ - and $\gamma$ -tocopherol Measurements

Tocopherol concentrations of the diets, fish, and embryos were determined by high-pressure liquid chromatography using electrochemical detection (HPLC-ECD; Shimadzu, Columbia, MD, USA, and LC-4C, Bioanalytical Systems, Inc. West Lafayette, IN, USA, respectively), as described previously [21]. In brief, the samples were weighed (diet samples ~50 mg, whole fish 5–900 mg, and 5–30 embryos [embryo estimated wet weight was 1 mg]), saponified in alcoholic KOH with 1% ascorbic acid at 70°C for 30 min (young fish, embryos and diets) or up to 1 h (adult fish). The samples were cooled, extracted with hexane, dried under nitrogen gas and resuspended in 50:50 ethanol/methanol, and an appropriate aliquot was injected into the HPLC-ECD. Tocopherols were quantified by comparison to calibration curves generated from authentic  $\alpha$ - and  $\gamma$ -tocopherol standards.

### 2.4. Ascorbic acid measurements

Sample collection methods were based on Moreau and Dabrowski [22] and Matamoros et al. [23]. Briefly, fish were euthanized by an overdose of tricaine (MS 222, ethyl 3-aminobenzoate methanesulfonate salt; Sigma-Aldrich, St. Louis, MO, USA) and weighed, and the entire fish were homogenized with buffer (5% trichloroacetic acid [Sigma-Aldrich], 0.08% diethylenetriaminepentaacetic acid [Acros Organics, Morris Plains, NJ, USA], 250 mM perchloric acid [Fisher Scientific, Fair Lawn, NJ, USA] and 0.4 mM dithioerythritol [Sigma-Aldrich]). Samples were then centrifuged; the supernatants were frozen and stored at –80°C until analysis. Samples were analyzed by HPLC-ECD, as described previously [24].

### 2.5. Zebrafish $\alpha$ -tocopherol depletion kinetics

At 4–6 weeks of age, zebrafish were randomly separated into three dietary groups: E–, E+ or lab diets. Zebrafish ( $n=3$ /dietary group) were euthanized by overdose of tricaine at each of the indicated time points. Zebrafish were weighed and kept frozen until analysis for vitamin concentrations, as described in sections 2.3 and 2.4. This experiment was repeated using three separate generations of fish.  $\alpha$ -Tocopherol depletion kinetics were calculated using Microsoft Excel (Microsoft Corporation, Santa Rosa, CA, USA) by fitting a linear regression analysis to the logarithmic-transformed  $\alpha$ -tocopherol concentrations measured during the first ~80 days after initiation of the E– diet.

### 2.6. Zebrafish sensory testing

To evaluate sensory responses, computer-assisted video monitoring of swimming behavior was assessed using modification of the method of Eddins et al. [25]. Adult zebrafish ( $n=6$ ) from each diet (E+, E– and lab; 221 days on diet) were placed in individual 1.75-L tanks containing ~1.5 L of FW. Tanks were set inline on shelves with the broadside facing the camera, separated by dividers to isolate individual fish. Tanks were backed with blank white paper, evenly backlit. Room temperature was controlled at 28°C. Fish were fasted for the duration of the behavior trials and were given 24 h to acclimatize prior to beginning the trials.

Trials were recorded using a Sony HD camcorder (Sony Handycam HDR-SR11) coupled with the Noldus EthoVision XT V 7.0 analysis software (Leesburg, VA, USA). The swimming velocity was recorded for 16-min sessions (separated into 2-min intervals); the first 6 min was averaged to obtain the average velocity for the trial, and the next 10 min was not used for assessment (rest time). Two trials with no stimulus

were used to generate “background” swimming information, and results from each fish were averaged to generate its own baseline information. Two trials were done using a “single tap” generated by an electromagnetic solenoid to tap the tank at 2 min; results from each fish were averaged to generate its own single tap information. A final “multiple tap” trial was generated using the solenoids to strike the tanks once every 5 s starting at 2 min for 90 s (18 taps in total). Rest time was given between each stimulus trial to allow a return to baseline behavior.

2.7. Zebrafish embryo handling and scoring

Fertilized eggs were obtained from natural spawning of adult zebrafish according to methods in *The Zebrafish Book* [15]. Embryos were collected and staged, as described by Kimmel et al. [14]. Embryos were treated with a dilute bleach solution (0.0033% = 0.55 ml household bleach diluted to 1 L of FW) to clean their chorions and were then rinsed twice with FW before being placed into 10-cm petri dishes containing methylene blue (0.0002%) to inhibit fungal growth.

To assess morphology of embryos over time, embryos were placed individually in wells of 96-well plates in 150 µl methylene blue (0.0002%) and observed daily using stereomicroscopy, up to 120 hpf. Scoring was assessed on visible phenotypes (mortality at 24 hpf; mortality at 120 hpf, delayed development, lack of motility, abnormal touch response, spastic movement and malformations of heart, brain, yolk sac, notochord, body axis, trunk, circulatory system, eye, jaw, somites, snout, otic, fin, pigmentation or swim bladder).

2.8. Quantitative real-time polymerase chain reaction

Fish were euthanized by tricaine overdose; the livers were removed and placed in RNALater (Qiagen, Valencia, CA, USA) and stored at –20° C until RNA extraction. Total RNA was extracted using an RNeasy kit with DNase I treatment per manufacture's directions (Qiagen). RNA concentrations and purity were determined by UV absorption (NanoDrop ND-1000 UV-Vis Spectrophotometer; Thermo Scientific, Wilmington, DE, USA). Complement DNA was synthesized following the manufacture's directions using Superscript III First-Strand Synthesis SuperMix for qRT-PCR (Invitrogen, Carlsbad, CA, USA). Primers were designed for each target gene using the Primer-BLAST program (Primer3 combined with BLAST, National Center for Biotechnology Information (NCBI, www.ncbi.nlm.nih.gov) Web site) (Table 2). Plasmids were cloned from each primer product (TOPO TA cloning kit; Invitrogen), sequenced to verify correct product (ABI Prism 3730 Genetic Analyzer, ABI Prism 3730 Data Collection Software v. 3.0, ABI Prism DNA Sequencing Analysis Software v. 5.2, with BigDye Terminator v. 3.1 Cycle Sequencing Kit; Center for Genome Research and Biocomputing core facility, Oregon State University, Corvallis, OR, USA), and concentrations were measured by spectrophotometer. These plasmids were used to generate an absolute copy-number standard curve for real-time polymerase chain reaction (PCR) quantification. Samples were analyzed using Platinum SYBR Green qPCR SuperMix-UDG (Invitrogen) using a DNA Engine Opticon 2 System (BioRad, Hercules, CA, USA) with the Opticon Monitor Version 3.0 software for real-time PCR detection. Results were normalized using glyceraldehyde-3-phosphate dehydrogenase (GAPDH), β-actin or β-2-microglobulin (β2M) expression. Data are reported as fold changes relative to values from lab control fish livers. There were no statistically significant differences in the housekeeping genes between the diet groups. The β2M gene was chosen because its abundance was similar to our genes of interest.

2.9. Statistics

GraphPad Prism (GraphPad Software, Inc., La Jolla, CA, USA) and JMP (SAS Institute, Cary NC, USA) were used for statistical analyses. The data were logarithmically




Fig. 1. Diet vitamin E concentrations. Measured α- and γ-tocopherol concentrations (means ± S.E.M., n = 3 separate diet batches) of experimental diets.

transformed, when unequal variances were observed between groups. Sensory evaluation was estimated using average velocity of the fish generated by the Noldus EthoVision software. Comparisons between lab, E+ and E– diets were calculated using one-way analysis of variance, comparisons between E+ and E– by Tukey's honestly significant differences (HSD). Relationships between the three diet groups over time were analyzed using two-way analysis of variance. Post hoc tests were carried out using paired comparisons (Tukey's HSD). Data are reported as means ± S.E.M.; differences were considered significant at P < .05.

3. Results

3.1. Diet vitamin E concentrations

The E– and E+–defined diets were formulated identically with the exception that the E– diet had no added α-tocopherol. The E– diet contained 400 times less α-tocopherol than the E+ diet (P < .0001; Fig. 1), which contained ~500 mg/kg. γ-Tocopherol concentrations were also less in the E– than the E+ diets (6.2 ± 1.1 and 10.4 ± 1.0 mg/kg, respectively; P < .0001). The lab diet contained both α- and γ-tocopherols. The flake food components contained the following α- and γ-tocopherol concentrations, respectively: Aquatox Flake, 440 ± 18 and 16 ± 1 mg/kg; Golden Pearls, 305 ± 8 and 10 ± 1 and Hikari Micropellets, 283 ± 4 and 5.9 ± 0.1 (n = 3 measurements).

3.2. Zebrafish growth on defined diets and tocopherol concentrations

Zebrafish at ~4–6 weeks of age were divided into three groups: E–, E+ and lab diets; then they were fed exclusively the indicated diets and the whole-body vitamin E concentrations measured at the indicated time on diet (up to 310 days; Fig. 2). Although the zebrafish increased weight during the first 3–4 months on diet (time

Table 2  
Primers for genes of interest

NCBI gene ID no.	Abbreviation <sup>a</sup>	Forward primer	Reverse primer
373125	AIF	5'-AAAGTCCGGAAGAGGGTGT-3'	5'-GCCTGGAGCTCAGCATTAAC-3'
30400	β2M	5'-CCACTCCGAAAGTTCATGTGT-3'	5'-ATCTCTTTCTCTGGGGTGAA-3'
58100	BAD	5'-CTGAAAGGAGAGCAACTGGG-3'	5'-CCGCATCAGACTCTTTGTGA-3'
57934	β-actin	5'-AAGCAGGAGTACCATGAGTC-3'	5'-TGGAGTCTCAGATGACATG-3'
317743	GAPDH	5'-GAATTCCTGGATACACGGAG-3'	5'-AAAGGGGTACATCTACTCC-3'
326857	GCLC	5'-CTATCTGGAGAATCATGGAGG-3'	5'-CATTTTCTCTGTGACCGG-3'
352928	Gpx4a	5'-CCTTGGAAATGGCATCAAAT-3'	5'-CCACCACACTGGATCTCG-3'
352929	Gpx4b	5'-GAAGTGGATGAAGGAGCAGC-3'	5'-GCCGCAGCATACACACTCTA-3'
79381	GSTp1	5'-TTCAGTCCAACGCCATGC-3'	5'-ATGAGATCTGATCGCCAACC-3'
322732	LOX	5'-GGCAGACCAAAAAGCTGAGCG-3'	5'-ATCGAGCGTCTGGGGCAGA-3'
360149	NRF2	5'-CAGACGAGGAGGAGCGGGA-3'	5'-CCACTGCTGCAACTCTGGGA-3'
559087	PLA2gIV	5'-ATTTATCTCGACCCCAACC-3'	5'-ACACCAGCAATTGTTGTCCA-3'
406370	PLA2gVI	5'-AAACTGCAGAGATGAGCCGT-3'	5'-GCCTGGTGTTCCTCCCAACAT-3'
325906	TTP	5'-ATGGGAGACGCAAGAAATG-3'	5'-TGCATGTGAATCCGTGTTT-3'

<sup>a</sup> AIF (pdc8), apoptosis-inducing factor (programmed cell death 8); BAD, BCL2-antagonist of cell death; GCLC, glutamate-cysteine ligase, catalytic subunit; Gpx4a, glutathione peroxidase 4a; Gpx4b, glutathione peroxidase 4b; GSTp1, glutathione S-transferase pi; LOX, arachidonate 12-lipoxygenase; NRF2, nuclear factor (erythroid-derived 2)-like 2; PLA2gIV, phospholipase A2, calcium dependent; PLA2gVI, phospholipase A2, calcium independent.




Fig. 2. Vitamin E depletion kinetics. Time course of body weights (A–C) and whole-body concentrations of  $\alpha$ - (D–F) and  $\gamma$ -tocopherol (G–I) from individual zebrafish fed E– (squares,  $n=172$ ; A, D, G), E+ (triangles,  $n=177$ ; B, E, H) or lab (circles,  $n=168$ ; C, F, I) diets. Results are from three separate generations of fish, samples taken at noted days after initiation of dietary treatments. Body weights (A–C) were not significantly different between the diet groups, but increased over time (time effect,  $P<.0001$ , months  $1 < 2 < 3-5 < 6-10$ ; Tukey paired comparisons, each  $P<.05$ ). (D–F) By month 2 ( $>30$  days), and thereafter, the  $\alpha$ -tocopherol concentrations of the E– zebrafish were significantly less than those of the other groups [diet  $\times$  time interaction,  $P=.0007$ ; E– for months 2, 3–5 and 6–10, different than E+ or lab diet groups (which were not different from each other at each time interval); Tukey paired comparisons,  $P<.05$ ]. The line indicates the exponential rate of depletion. (G–I) Within the first month ( $<30$  days) and thereafter, the  $\gamma$ -tocopherol concentrations of the E– zebrafish were significantly less than those of the other groups [diet  $\times$  time interaction,  $P<.0001$ ; E– for months 1, 2, 3–5 and 6–10, different than E+ or lab (which were not different from each other at each time interval); Tukey HSD,  $P<.05$ ].



Table 3  
Zebrafish vitamin E concentrations and body weights taken at various times between ~80 and ~300 days consuming experimental diets

Diet	Observations (n)	α-Tocopherol (nmol/g)	γ-Tocopherol (nmol/g)	Body weight (g)
E–	80	5±3 <sup>a</sup>	1.8±1.0 <sup>a</sup>	379±154
E+	84	369±131 <sup>b</sup>	3.6±2.4 <sup>b</sup>	334±145
Lab	80	362±107 <sup>b</sup>	4.2±2.2 <sup>c</sup>	334±108

A significant diet effect ( $P<.0001$ ) was observed for both zebrafish α- and γ-tocopherol concentrations. Data in columns not bearing the same letter are significantly different ( $P<.05$ ). Time points are indicated in Fig. 2.

effect,  $P<.0001$ ), the various dietary treatments (E–, E+ and lab) had no significant effects on body weights over the course of the study (Fig. 2A–C). The rate of weight increase was  $2.3\pm0.1$ ,  $2.1\pm0.1$  and  $2.0\pm0.1$  g/d for the E–, E+ and lab fish.

Prior to initiating the experimental diets, zebrafish α-tocopherol concentrations were  $33.5\pm2.5$  nmol/g. When fed the E+ or lab diets, the α-tocopherol concentrations of zebrafish initially increased over time (Fig. 2E or F). In contrast, when zebrafish consumed the E– diet, the α-tocopherol concentrations decreased exponentially for the first ~80 days at a rate of  $-0.029\pm0.006$  nmol/g ( $n=3$  experiments), which yields a half-life of  $25\pm5$  days (Fig. 2D). After three half-lives, hypothetically only 12.5% of the starting vitamin E concentration should remain. Consistent with this prediction, between 80 and 300 days of diet consumption, the zebrafish fed the E– diets had α-tocopherol concentrations that were approximately 10% of their starting concentrations and more than ~50 times lower than zebrafish consuming either the E+ or lab diets (Fig. 2E or F, Table 3; diet effect,  $P=.0007$ ; paired comparisons,  $P<.05$ ). After this observed initial decrease, the fish maintained consistently low α-tocopherol concentrations (Fig. 2D; Table 3).

After 80 days, γ-tocopherol concentrations in zebrafish consuming the E– and E+ diets were lower than those in the fish consuming the lab diet; those of the zebrafish consuming the E– diet were less than half of those consuming the other diets (Fig. 2G–I; Table 3; diet effect,  $P<.0001$ ; paired comparisons,  $P<.05$ ).

3.3. Impaired sensory perception in zebrafish consuming E– diets

Sensory neuropathy is one of the first symptoms of vitamin E deficiency in humans [26]. To assess sensory neuropathy in




Fig. 3. Vitamin E deficiency alters startle responses. Adult zebrafish ( $n=6$ ) from each diet (E+, E– and lab; 221 days on diet) were placed in individual 1.75-L tanks containing ~1.5 L of FW. In the baseline trials and multitap trials, no significant differences were observed in the average swimming velocity between the diet groups. When startled by a single tap, the E+ and Lab zebrafish swam faster, while E– fish had an attenuated response (diet effect,  $P<.003$ ; bars not sharing the same letter are significantly different, Tukey HSD,  $P<.05$ ). When exposed to multiple taps, the fish did not show significant differences in swimming velocity.

Table 4  
mRNA expression of in adult zebrafish liver from fish fed E–, E+ or lab diets

Gene symbol	Diet		
	E–	E+	Lab
BAD	56±22	57±23	100±29
GCLC	76±42	67±29	100±6
GSTp1	100±48	70±40	100±20
GPX4b	64±19	56±31	100±14
LOX	73±26	77±28	100±42
NRF2	98±42	85±32	100±20
TTP	64±31	42±27	100±38

mRNA expression (mean±S.E.M.) in adult zebrafish livers from fish from each diet group ( $n=4$  per group) was analyzed by qPCR. Genes are defined, and primers are shown in Table 2. Expression levels were normalized to β2M and are shown as fold change over the average of the lab diet control liver mRNA, set to 100. No significant differences were found for a diet effect for the indicated genes.

zebrafish, we devised a “startle response” measurement using a percussive “tap” with responses monitored by video recording. In the baseline trial, no significant differences were observed in the average swimming velocity between the diet groups. Application of a single synchronized solenoid tap showed a characteristic startle




Fig. 4. α- and γ-Tocopherol concentrations of embryos and adults. Adult zebrafish ( $n=12$  per diet) were collected for vitamin E analysis between 250 and 300 days of consuming the diets. Zebrafish from this generation were spawned at 270, 278 and 284 days. Embryos were collected for vitamin E analysis at 48 hpf in groups of 15 embryos (E–,  $n=16$ ; E+,  $n=11$ ; lab,  $n=12$  replicates). (A) All embryo α-tocopherol concentrations (mean±S.E.M.; logarithmic scale) were less than those of the adult zebrafish; E– embryo α-tocopherol concentrations were less than those of E– adults; E– embryos compared with E+ or lab embryos had the lowest α-tocopherol concentrations (diet × life stage interaction,  $P=.016$ ; bars not sharing the same letter are significantly different, Tukey HSD,  $P<.05$ ). (B) Adult E+ and lab zebrafish γ-tocopherol concentrations (mean±S.E.M.) were greater than the adult E– zebrafish and γ-tocopherol concentrations in all of the embryos; the lab embryo γ-tocopherol concentrations were greater than those of the E+ embryos; E– embryo γ-tocopherol concentrations were not significantly different from the E– adults γ-tocopherol concentrations, while E+ and lab embryos were less than those of the adults (diet × life stage interaction,  $P<.0001$ ; bars not sharing the same letter are significantly different, Tukey HSD,  $P<.05$ ).




Fig. 5. Malformations and mortality of zebrafish embryos. (A) Increased mortality (mean  $\pm$  S.E.M.) was observed in the E– embryos at 24 and 120 hpf compared with the other diet groups (diet  $\times$  time interaction,  $P < .0001$ ), but at 24 hpf, the differences between diet groups did not reach statistical significance. Mortality increased from 24 to 120 hpf in the E– (squares;  $P < .01$ ) and the E+ embryos (triangles;  $P < .01$ ), but not in the lab diet embryos (circles). At 120 hpf, the E– embryos displayed significantly higher levels of mortality compared with the E+ and lab diet embryos (diet effect,  $P = .005$ ; E– (a)  $>$  E+ (b) or lab (b);  $P < .05$  for paired comparisons). (B) Higher levels of both malformations and mortality were observed at 120 hpf in the E– embryos compared with E+ ( $P < .05$ ; a) or lab diet embryos ( $P < .001$ ; b); E+ had greater malformations than did lab diet embryos ( $P < .05$ ; c). Embryos were analyzed in 96-well plates, one embryo per well with 48 to 120 embryos per group per spawn. Results are expressed as percentages affected per total number of embryos ( $n = 6$  spawns per group).

response in the E+ and lab diet zebrafish [25]. When startled by a single tap, the E+ and lab zebrafish both swam faster, while E– fish had an attenuated response (Fig. 3; swim effect,  $P < .003$ ; paired comparisons,  $P < .05$ ). These differences disappeared when the fish were exposed to a higher level of stimulation in the “multiple tap” trial, suggesting that a greater stimulus was needed to elicit a response from the E– fish.

#### 3.4. Embryo tocopherol concentrations

Fish from each of the diet groups were group spawned starting at 3–4 months of age (~80 days on diets). Despite differences in  $\alpha$ -tocopherol concentrations, all dietary groups of zebrafish routinely produced viable gametes. The embryos were observed up to 120 hpf without being fed. Normally, zebrafish embryos do not consume food during this period but receive their nutrition from the yolk sac [13,14].

Embryo  $\alpha$ -tocopherol concentrations reflected those of their parents. For example, adult zebrafish that had consumed the diets for  $>250$  days were spawned, and the  $\alpha$ - and  $\gamma$ -tocopherol concentrations were measured in representative adult fish and in 48-hpf embryos (Fig. 4). Adult zebrafish consuming the E+ and lab diets contained nearly 50 times higher  $\alpha$ -tocopherol concentrations

than did those consuming the E– diet; the embryos from the fish consuming the E+ and lab diets contained nearly 30 times higher  $\alpha$ -tocopherol concentrations than embryos in the E– group (diet  $\times$  life stage interaction,  $P = .016$ ;  $P < .05$  for comparisons; Fig. 4A).

The adult zebrafish consuming the E+ and lab diets contained nearly three times higher  $\gamma$ -tocopherol concentrations than did the adults consuming the E– diet. However, in the E– and lab embryos, the  $\gamma$ -tocopherol concentrations were almost double those of the E+ embryos (diet  $\times$  life stage interaction,  $P < .0001$ ; E+ compared with lab,  $P < .05$ ; Fig. 4B). Thus, E– embryo  $\gamma$ -tocopherol concentrations were not significantly different from the E– adults'  $\gamma$ -tocopherol concentrations, while the  $\gamma$ -tocopherol concentrations of the embryos from the E+ and lab diet groups were lower than those of their parents (E+ or lab adults,  $P < .05$ ). These findings indicate that the adult female zebrafish transfer very little  $\gamma$ -tocopherol to the eggs, even during vitamin E deficiency.

Embryo vitamin E status did not vary significantly from 48 to 120 hpf (data not shown).

#### 3.5. Embryo morphology and mortality

Embryos from spawns of each diet group were observed up to 120 hpf for morphologic changes. Low levels of abnormalities are inherent in zebrafish embryos and were observed in embryos from all the diet groups. In the E– embryos, there was increased mortality at 24 and 120 hpf compared with the other diet groups (Fig. 5A; diet  $\times$  time interaction,  $P < .0001$ ), but at 24 hpf, the differences between diet groups did not reach statistical significance for paired comparisons. At




Fig. 6. Typical zebrafish morphology at 120 hpf. Representative pictures from the three diet groups are shown after 5 days (120 h). The eye and otic vesicle are indicated on all fry; malformations are illustrated on the image of the deficient fish. CF, cranial–facial malformation; BA, bent anterior–posterior axis; PE, pericardial edema; SB, swim bladder malformation; YSE, yolk-sac edema.




Fig. 7. Ascorbic acid concentrations in adult fish. Ascorbic acid concentrations (mean  $\pm$  S.E.M.) were analyzed from two generations of zebrafish ( $n=7$  per diet; Diet effect,  $P<.0035$ ; bars not sharing the same letter are significantly different,  $P<.05$ ).

120 hpf, the E- embryos displayed significantly higher levels of mortality compared with the E+ or lab diet embryos (diet effect,  $P=.005$ ;  $P<.05$  for paired comparisons).

At 120 hpf, nearly 70% of the E- embryos experienced either malformations or death. These levels were higher than those observed in the E+ or lab diet embryos; E+ had greater malformations than did lab diet embryos (Fig. 5B;  $P<.05$ ). Malformations commonly observed included cranial-facial malformations, bent anterior-posterior axis, pericardial edema, swim bladder malformations and yolk-sac edema (Fig. 6).

### 3.6. Whole-fish ascorbic acid concentrations

Ascorbic acid concentrations were measured in adult zebrafish after consuming E-, E+ or lab diets for various amounts of time ( $>181$  days). Ascorbic acid concentrations of the fish on the defined diets were nearly double those of the fish consuming the lab diet (diet effect,  $P<.0035$ ;  $P<.05$  for paired comparisons; Fig. 7).

### 3.7. Liver gene expression

To assess long-term effects of vitamin E deficiency on mRNA expression levels of liver genes of interest, we used quantitative PCR (qPCR) to measure transcripts of genes involved in oxidative stress response, lipid metabolism, vitamin E trafficking and cell death [27–31] (genes are listed in Table 2). Expression levels were normalized to  $\beta 2M$ ,  $\beta$ -actin or *GAPDH*; all three housekeeping genes yielded similar results; there were no statistically significant differences between the diets for each of the housekeeping genes. The data shown in Table 4 are normalized to  $\beta 2M$  and are shown as fold changes over the average of the lab diet control liver mRNA, set to 100. We observed no significant changes in mRNA levels due to chronic vitamin E deficiency in the genes assayed. However, the expression of four genes, which have previously been identified to be important in nerve protection [32], were found to be significantly reduced in the zebrafish fed E- or E+ diets relative to those in the control fish (Fig. 8).

## 4. Discussion

Vitamin E deficiency caused developmental abnormalities and death in zebrafish embryos. This finding elucidates for the first time that the embryo, as well as the mother, requires vitamin E. The expression of TTP by the human embryo-yolk sac [8], as well as by the zebrafish embryo-yolk sac [9], suggested that the embryo itself might require vitamin E. Initially, in 1922, vitamin E was discovered because rats fed rancid fat failed to carry their offspring to term [1]. This




Fig. 8. mRNA expression in adult zebrafish liver. mRNA expression (mean  $\pm$  S.E.M.) in adult zebrafish livers from fish from each diet group ( $n=4$  per group, on diets  $>200$  days) was analyzed by qPCR. Genes are defined, and primers are shown in Table 2. Expression levels were normalized to  $\beta 2M$  and are shown as fold change over the average of the lab diet control liver mRNA, set to 100 (diet effect for *GPX4a*,  $P=.0011$ ; *PLA2*,  $P=.0027$ ; *PLA2GIV*,  $P=.0058$ ; *AIF*,  $P=.0222$ ; bars not sharing the same letter are significantly different,  $P<.05$ ).

finding was the basis for the “fetal resorption test” that remains in use today as an assay of biologic activities of various vitamin E forms [33]. In order to prevent fetal resorption, vitamin E must be administered to the vitamin E-deficient rat mother on postfertilization days 5 to 9 [34,35]. Interestingly, this is the same critical period where the 12/15-lipoxygenase-dependent pathway appears to mediate implantation [36]. Similarly, *glutathione peroxidase 4* (*GPx4*) expression increases at day 7.5 and is the same time at which *GPx4* knockout mice embryos are resorbed [37]. Taken together, these findings suggest that during days 5 to 9 of rodent embryogenesis, lipid peroxidation can be especially damaging. Importantly, vitamin E is a potent, lipid-soluble antioxidant and thus may be critical for the embryo to have sufficient amounts to protect against lipid peroxidation [38].

Both low fetal and maternal  $\alpha$ -tocopherol concentrations may be important factors during fetal resorption. Retention of uterine  $\alpha$ -tocopherol by the mother appears essential to maintain pregnancy because TTP is increased at the site of implantation [2,3] and is also expressed in the syncytiotrophoblast of the human placenta [3–6]. Early failure of pregnancy is associated with lipid peroxidation with resultant damage to the syncytiotrophoblast in the placenta [7]. Thus, it is likely that  $\alpha$ -tocopherol is needed by the mother to protect her from the oxidative stress of the rapidly growing fetus. Herein, we demonstrate that the embryo independently requires vitamin E because the E- zebrafish embryos suffered increased developmental abnormalities, as well as increased mortality (Figs. 5 and 6). These E- embryos contained significantly lower  $\alpha$ -tocopherol concentrations than the embryos in the other diet groups (Fig. 4).

Importantly, the E- diet depleted the adult zebrafish of  $\alpha$ -tocopherol (Fig. 2, Table 3). The exponential rate of depletion predicted that the zebrafish fed the E- diet would experience nearly 90% depletion of their initial  $\alpha$ -tocopherol concentration by 80 days on diet. The  $\alpha$ -tocopherol measurements beyond 100 days showed that further depletion did not occur, or at least was slowed to statistically imperceptible changes. Given that the E- diet had some  $\alpha$ -tocopherol and that the zebrafish experienced deficiency symptoms, these data suggest that the dietary  $\alpha$ -tocopherol concentration (approximately 2 mg/kg diet) is below the requirement for the zebrafish. Using the estimate that a fish weighing 500 mg eats about 10 mg food [39], the zebrafish is probably consuming about 50 pmol



$\alpha$ -tocopherol, then after 80 days on the E– diet, its whole body contains about 2500 pmol  $\alpha$ -tocopherol.

Remarkably, like humans [40,41], zebrafish show a preference for  $\alpha$ - over  $\gamma$ -tocopherol. This finding is most easily appreciated in the comparison of the vitamin E distribution within the E– diets compared with the E– zebrafish. The E– diets had approximately 5 times more  $\gamma$ - than  $\alpha$ -tocopherol (Fig. 1), while the E– zebrafish contained approximately 2.5 times more  $\alpha$ - than  $\gamma$ -tocopherol (Table 3). Given that the zebrafish expresses *TTP* gene [9], it is not surprising that the zebrafish shows a preference for  $\alpha$ -tocopherol. More remarkable is the apparent lack of  $\gamma$ -tocopherol transfer to the embryos (Fig. 4). Although the E+ and lab adult zebrafish contained  $\gamma$ -tocopherol at three times the E– adult zebrafish's  $\gamma$ -tocopherol concentrations, the E– embryo  $\gamma$ -tocopherol concentrations were not significantly different from those of either the E+ and lab embryos. These data emphasize that the embryos are preferentially enriched in  $\alpha$ -tocopherol and that the E– embryos were limited in  $\gamma$ -tocopherol even when the zebrafish adults were severely limited in both forms of vitamin E. These findings are reminiscent of those in *TTP* null mice that were fed  $\gamma$ -tocopherol-enriched diets yet did not accumulate  $\gamma$ -tocopherol in tissues or plasma [42]. Thus, vitamin E deficiency does not drive the use of  $\gamma$ -tocopherol in place of  $\alpha$ -tocopherol.

This study also demonstrates that the adult zebrafish require vitamin E because we detected abnormal swimming responses to a “tap” test (Fig. 3). In general, testing of neurologic responses of adult zebrafish has been limited. Eddins et al. [25] studied chlorpyrifos neurotoxicity that was induced during early development and then persisted into adulthood. They showed that the neurotoxicity caused an increase in swimming velocity in the affected adult zebrafish [25]. In contrast, we found that vitamin E deficiency in zebrafish decreased their responsiveness, as shown by decreased swimming velocity, suggesting diminished sensory perception. It should be noted that muscle degeneration is also associated with vitamin E deficiency, but in the current study, the zebrafish were apparently able to swim faster in response to multiple taps; they just did not respond quickly to a single stimulus. This finding suggests that it took greater stimulus to elicit a response. Given that the vitamin E deficiency symptom in humans is a dying back of the sensory neurons [43], it seems likely that the E– zebrafish also had a sensory deficit, but further studies are needed to evaluate the extent and mechanism of the deficit.

To assess long-term effects of vitamin E deficiency on mRNA expression levels of liver genes of interest, we used quantitative PCR (qPCR) to measure transcripts of genes involved in oxidative stress response, lipid metabolism, vitamin E trafficking and cell death [27–31]. Liver expression of these genes was not dependent on vitamin E status since the fish from the E– and E+ groups displayed similar responses. However, the lab diet contains various ingredients, such as fish oil, that are not present in the defined diets. These “control” livers had higher expression of *GPX4a*, *phospholipase A2* (*PLA2*), *phospholipase A2* (calcium dependent) and *apoptosis-inducing factor* (*AIF*), suggesting that the more oxidizable lipids had induced expression of these protective enzymes. However, we have not measured lipid oxidation products in these fish, so the mechanisms for the differential expression remain to be investigated. It is also apparent that the embryo is more sensitive than is the adult fish to insufficient vitamin E. We plan further studies to investigate vitamin E-dependent gene regulation in the embryo.

We formulated a diet containing sufficient nutrients for the zebrafish based on previous research in other fresh water teleost fish [17–20]. Our diet formulation (Table 1) was sufficient for zebrafish nutrition based on growth and body weights (Fig. 2, Table 3). Animal health, behavior and mortality were observed daily during feeding, and no overt visual changes were noted in fish consuming the defined diets. Previously, commercially available diets for zebrafish were

insufficient to maintain growth and had to be supplemented with live food [39]. Our diets are, therefore, a major step forward in zebrafish husbandry given that 45-day-old zebrafish fed these diets grow as well as those fed the commercial mixture of foods and live food.

In addition, we analyzed whole zebrafish ascorbic acid concentrations, as vitamin C status has been linked to vitamin E status [44], and teleost fish lack the  $\alpha$ -gulonolactone oxidase gene and thus cannot synthesize ascorbic acid [45]. Zebrafish ascorbic acid concentrations were higher in the fish fed the defined diets than in the lab controls, showing that the defined diets are more than sufficient in ascorbic acid (Fig. 7).

In conclusion, our findings document for the first time that vitamin E is required not only for the mother but also for the embryo during development. This zebrafish model has allowed the successful separation of maternal and embryonic nutrition, permitting the study of embryogenesis without the fetal resorption that occurs in mammalian models.

## Acknowledgments

The authors gratefully acknowledge Cari Buchner, Jane LaDu, Sumitra Sengupta and the Sinnhuber Aquatic Research Laboratory staff for excellent assistance with zebrafish husbandry; Brent Flickinger from Archer Daniels Midland for the gifts of vitamin E and lecithin and Andrew Ross and Yee-Ling “Caryn” Ong in the Crop and Food Science, Oregon State University, for the use of laboratory facilities for diet preparation.

## References

- [1] Evans HM, Bishop KS. On the existence of a hitherto unrecognized dietary factor essential for reproduction. *Science* 1922;56:650–1.
- [2] Jishage K, Arita M, Igarashi K, Iwata T, Watanabe M, Ogawa M, et al. Alpha-tocopherol transfer protein is important for the normal development of placental labyrinthine trophoblasts in mice. *J Biol Chem* 2001;276:1669–72.
- [3] Kaempf-Rotzoll DE, Igarashi K, Aoki J, Jishage K, Suzuki H, Tamai H, et al. Alpha-tocopherol transfer protein is specifically localized at the implantation site of pregnant mouse uterus. *Biol Reprod* 2002;67:599–604.
- [4] Kaempf-Rotzoll DE, Horiguchi M, Hashiguchi K, Aoki J, Tamai H, Linderkamp O, et al. Human placental trophoblast cells express alpha-tocopherol transfer protein. *Placenta* 2003;24:439–44.
- [5] Muller-Schmehl K, Beninde J, Finckh B, Florian S, Dudenhausen JW, Brigelius-Flohe R, et al. Localization of alpha-tocopherol transfer protein in trophoblast, fetal capillaries' endothelium and amnion epithelium of human term placenta. *Free Radic Res* 2004;38:413–20.
- [6] Rotzoll DE, Scherling R, Etzl R, Stepan H, Horn LC, Poschl JM. Immunohistochemical localization of alpha-tocopherol transfer protein and lipoperoxidation products in human first-trimester and term placenta. *Eur J Obstet Gynecol Reprod Biol* 2008;140:183–91.
- [7] Hempstock J, Jauniaux E, Greenwold N, Burton GJ. The contribution of placental oxidative stress to early pregnancy failure. *Hum Pathol* 2003;34:1265–75.
- [8] Jauniaux E, Cindrova-Davies T, Johns J, Dunster C, Hempstock J, Kelly FJ, et al. Distribution and transfer pathways of antioxidant molecules inside the first trimester human gestational sac. *J Clin Endocrinol Metab* 2004;89:1452–8.
- [9] Usenko CY, Harper SL, Tanguay RL. Fullerene C60 exposure elicits an oxidative stress response in embryonic zebrafish. *Toxicol Appl Pharmacol* 2008;229:44–55.
- [10] Dodd A, Curtis PM, Williams LC, Love DR. Zebrafish: bridging the gap between development and disease. *Hum Mol Genet* 2000;9:2443–9.
- [11] Udvardi AJ, Linney E. Windows into development: historic, current, and future perspectives on transgenic zebrafish. *Dev Biol* 2003;256:1–17.
- [12] Wixon J. Featured organism: *Danio rerio*, the zebrafish. *Yeast* 2000;17:225–31.
- [13] Kimmel CB. Genetics and early development of zebrafish. *Trends Genet* 1989;5:283–8.
- [14] Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic development of the zebrafish. *Dev Dyn* 1995;203:253–310.
- [15] Westerfield M. The zebrafish book: a guide for the laboratory use of zebrafish (*Danio rerio*). 4. ed. Eugene, OR: M. Westerfield; 2000.
- [16] Peterson C, Gustin M. Mercury in the air, water and biota at the Great Salt Lake (Utah, USA). *Sci Total Environ* 2008;405:255–68.
- [17] DeKoven DL, Nunez JM, Lester SM, Conklin DE, Marty GD, Parker LM, et al. A purified diet for medaka (*Oryzias latipes*): refining a fish model for toxicological research. *Lab Anim Sci* 1992;42:180–9.
- [18] Markovich ML, Rizzuto NV, Brown PB. Diet affects spawning in zebrafish. *Zebrafish* 2007;4:69–74.



- [19] National Research Council (U.S.). Committee on Animal Nutrition. Nutrient requirements of fish. Washington, D.C.: National Academy Press; 1993.
- [20] Siccardi AJ, Garriss HW, Jones WT, Moseley DB, D'Abramo LR, Watts SA. Growth and survival of zebrafish (*Danio rerio*) fed different commercial and laboratory diets. *Zebrafish* 2009;6:275–80.
- [21] Podda M, Weber C, Traber MG, Packer L. Simultaneous determination of tissue tocopherols, tocotrienols, ubiquinol, and ubiquinones. *J Lipid Res* 1996;37:893–901.
- [22] Moreau R, Dabrowski K. Body pool and synthesis of ascorbic acid in adult sea lamprey (*Petromyzon marinus*): an agnathan fish with gulonolactone oxidase activity. *Proc Natl Acad Sci U S A* 1998;95:10279–82.
- [23] Matamoros MA, Loscos J, Coronado MJ, Ramos J, Sato S, Testillano PS, et al. Biosynthesis of ascorbic acid in legume root nodules. *Plant Physiol* 2006;141:1068–77.
- [24] Frei B, England L, Ames BN. Ascorbate is an outstanding antioxidant in human blood plasma. *Proc Natl Acad Sci U S A* 1989;86:6377–81.
- [25] Eddins D, Cerutti D, Williams P, Linney E, Levin ED. Zebrafish provide a sensitive model of persisting neurobehavioral effects of developmental chlorpyrifos exposure: comparison with nicotine and pilocarpine effects and relationship to dopamine deficits. *Neurotoxicol Teratol* 2009;32:99–108.
- [26] Ouahchi K, Arita M, Kayden H, Hentati F, Ben Hamida M, Sokol R, et al. Ataxia with isolated vitamin E deficiency is caused by mutations in the alpha-tocopherol transfer protein. *Nat Genet* 1995;9:141–5.
- [27] Todorovic M, Skugor S, Ruyter B. Alterations in oxidative stress status modulate terminal differentiation in Atlantic salmon adipocytes cultivated in media rich in n-3 fatty acids. *Comp Biochem Physiol B Biochem Mol Biol* 2010;156:309–18.
- [28] Surh YJ, Kundu JK, Li MH, Na HK, Cha YN. Role of Nrf2-mediated heme oxygenase-1 upregulation in adaptive survival response to nitrosative stress. *Arch Pharm Res* 2009;32:1163–76.
- [29] Na YR, Seok SH, Baek MW, Lee HY, Kim DJ, Park SH, et al. Protective effects of vitamin E against 3,3',4,4',5-pentachlorobiphenyl (PCB126) induced toxicity in zebrafish embryos. *Ecotoxicol Environ Saf* 2009;72:714–9.
- [30] Johnson JA, Johnson DA, Kraft AD, Calkins MJ, Jakel RJ, Vargas MR, et al. The Nrf2-ARE pathway: an indicator and modulator of oxidative stress in neurodegeneration. *Ann N Y Acad Sci* 2008;1147:61–9.
- [31] Feng Z, Liu Z, Li X, Jia H, Sun L, Tian C, et al. Alpha-tocopherol is an effective phase II enzyme inducer: protective effects on acrolein-induced oxidative stress and mitochondrial dysfunction in human retinal pigment epithelial cells. *J Nutr Biochem* 2010;21:1222–31.
- [32] Seiler A, Schneider M, Forster H, Roth S, Wirth EK, Culmsee C, et al. Glutathione peroxidase 4 senses and translates oxidative stress into 12/15-lipoxygenase dependent- and AIF-mediated cell death. *Cell Metab* 2008;8:237–48.
- [33] Traber MG. Vitamin E. In: Shils ME, Olson JA, Shike M, Ross AC, editors. *Modern nutrition in health and disease*. Baltimore: Williams & Wilkins; 1999. p. 347–62.
- [34] Ames SR. Biopotencies in rats of several forms of alpha-tocopherol. *J Nutr* 1979;109:2198–204.
- [35] Leth T, Sondergaard H. Biological activity of vitamin E compounds and natural materials by the resorption-gestation test, and chemical determination of the vitamin E activity in foods and feeds. *J Nutr* 1977;107:2236–43.
- [36] Li Q, Cheon YP, Kannan A, Shanker S, Bagchi IC, Bagchi MK. A novel pathway involving progesterone receptor, 12/15-lipoxygenase-derived eicosanoids, and peroxisome proliferator-activated receptor gamma regulates implantation in mice. *J Biol Chem* 2004;279:11570–81.
- [37] Imai H, Hirao F, Sakamoto T, Sekine K, Mizukura Y, Saito M, et al. Early embryonic lethality caused by targeted disruption of the mouse PHGPx gene. *Biochem Biophys Res Commun* 2003;305:278–86.
- [38] Traber MG, Atkinson J. Vitamin E, antioxidant and nothing more. *Free Radic Biol Med* 2007;43:4–15.
- [39] Lawrence C. The husbandry of zebrafish (*Danio rerio*): a review. *Aquaculture* 2007;269:1–20.
- [40] Leonard SW, Paterson E, Atkinson JK, Ramakrishnan R, Cross CE, Traber MG. Studies in humans using deuterium-labeled alpha- and gamma-tocopherols demonstrate faster plasma gamma-tocopherol disappearance and greater gamma-metabolite production. *Free Radic Biol Med* 2005;38:857–66.
- [41] Traber MG, Kayden HJ. Preferential incorporation of alpha-tocopherol vs gamma-tocopherol in human lipoproteins. *Am J Clin Nutr* 1989;49:517–26.
- [42] Traber MG, Siddens LK, Leonard SW, Schock B, Gohil K, Krueger SK, et al.  $\alpha$ -Tocopherol modulates Cyp3a expression, increases  $\gamma$ -CEHC production and limits tissue  $\gamma$ -tocopherol accumulation in mice fed high  $\gamma$ -tocopherol diets. *Free Radic Biol Med* 2005;38:773–85.
- [43] Food and Nutrition Board. Institute of Medicine. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Washington: National Academy Press; 2000.
- [44] Bruno RS, Leonard SW, Atkinson J, Montine TJ, Ramakrishnan R, Bray TM, et al. Faster plasma vitamin E disappearance in smokers is normalized by vitamin C supplementation. *Free Radic Biol Med* 2006;40:689–97.
- [45] Dabrowski K. Gulonolactone oxidase is missing in teleost fish. The direct spectrophotometric assay. *Biol Chem Hoppe Seyler* 1990;371:207–14.