

STUDIES IN THE GENERAL THEORY OF RULED SURFACES*

BY

E. J. WILCZYNSKI†

The congruence Γ , which is made of all the generators of the first kind on the osculating hyperboloids of a ruled surface, has a great many interesting properties. Some of them have been considered in a previous paper.‡ We shall continue the consideration of this congruence and of configurations associated with it, completing in this way some of our previous investigations very essentially. We shall also study the osculating linear complex, and the point-to-plane correspondence to which it gives rise, enabling us to generalize some well-known theorems of CREMONA and LIE.

For the most part we shall, in this paper, confine our attention to the general case, when the flecnodal curve intersects every generator in two distinct points. The case of coincidence will be left for a future occasion, as it requires the use of a different normal form for the equations than that here adopted.

The notations are the same as in previous papers. To save space they are not again explained.

§ 1. *The derivative cubic curve.*

If α_1 and α_2 are arbitrary, $\alpha_1 y_k + \alpha_2 z_k$ will represent the coördinates of an arbitrary point on the generator g of the ruled surface, where (y_k, z_k) for $k = 1, 2, 3, 4$ are four simultaneous systems of solutions of our system of differential equations, whose determinant does not vanish. We shall usually write $\alpha_1 y + \alpha_2 z$, suppressing the index k , as in previous papers. Of course this is essentially a form of vector analysis, which enables us to make one equation do the work of four. The point $\alpha_1 \rho + \alpha_2 \sigma$ of the corresponding generator g' of S' , will then be such that the line joining it to $\alpha_1 y + \alpha_2 z$ is a generator of the sec-

* Presented to the Society (San Francisco) December 19, 1903. Received for publication October 20, 1904.

† Of the Carnegie Institution of Washington.

‡ On a certain congruence associated with a given ruled surface. Transactions of the American Mathematical Society, vol. 4 (1903), pp. 185–200, hereafter referred to as *Congruence*. My other papers will be quoted by the initial words of their title. See *Covariants*, vol. 3 (1902), p. 423, footnote.

and kind on the hyperboloid H osculating S along g . Therefore if β_1 and β_2 are arbitrary,

$$\beta_1(\alpha_1 y + \alpha_2 z) + \beta_2(\alpha_1 \rho + \alpha_2 \sigma)$$

will be an arbitrary point of H .

If we choose the tetrahedron $P_y P_z P_\rho P_\sigma$ as tetrahedron of reference for a system of homogeneous coördinates, we may take them so that the coördinates of any point represented by an expression of the form $\lambda y + \mu z + \nu \rho + \kappa \sigma$ will be $x_1 = \lambda$, $x_2 = \mu$, $x_3 = \nu$, $x_4 = \kappa$. We have then

$$x_1 = \beta_1 \alpha_1, \quad x_2 = \beta_1 \alpha_2, \quad x_3 = \beta_2 \alpha_1, \quad x_4 = \beta_2 \alpha_2$$

as the coördinates of an arbitrary point on H , and therefore

$$(1) \quad x_1 x_4 - x_2 x_3 = 0$$

as the equation of H in this system of coördinates.

Let us consider now the hyperboloid H' , which osculates S' along g' . The coördinates of P_ρ and P_σ were obtained from the system of differential equations defining S by forming

$$\rho = 2y' + p_{11}y + p_{12}z, \quad \sigma = 2z' + p_{21}y + p_{22}z.$$

We shall obtain the coördinates of two points on a generator g'' of the derivative of S' with respect to x , by applying the same process to the equations [Congruence, (15)] which define S' . The ruled surface S'' thus obtained shall be naturally called the second derivative of S with respect to x . Its generator g'' is then a generator of the hyperboloid H' which osculates S' along g' . The following quantities

$$(2) \quad \begin{aligned} 2\rho' + P_{11}\rho + P_{12}\sigma &= u_{11}y + u_{12}z + \lambda_{11}\rho + \lambda_{12}\sigma, \\ 2\sigma' + P_{21}\rho + P_{22}\sigma &= u_{21}y + u_{22}z + \lambda_{21}\rho + \lambda_{22}\sigma \end{aligned}$$

are the coördinates of two points on g'' .

These equations show that g'' intersects g , if and only if $\lambda_{11}\lambda_{22} - \lambda_{12}\lambda_{21} = 0$, i. e., if $K = v_{11}v_{22} - v_{12}v_{21} = 0$, provided we assume that S' is not developable. By changing the independent variable one can always change K into \bar{K} such that $\bar{K} = 0$. The equation for η , [Congruence, equ. (4)], which must be satisfied so as to make $\bar{K} = 0$, is of the second order. Therefore there exist ∞^2 non-developable ruled surfaces in the congruence Γ , each of which, when considered as the first derivative of S , gives rise to a second derivative whose generators intersect the corresponding generators of S .

Let us consider any point on g' , whose coördinates are $\epsilon_1 \rho + \epsilon_2 \sigma$. The corresponding point on g'' will be given by

$$(\epsilon_1 u_{11} + \epsilon_2 u_{21})y + (\epsilon_1 u_{12} + \epsilon_2 u_{22})z + (\epsilon_1 \lambda_{11} + \epsilon_2 \lambda_{21})\rho + (\epsilon_1 \lambda_{12} + \epsilon_2 \lambda_{22})\sigma.$$

Therefore, the expression

$$\delta_1(\epsilon_1 u_{11} + \epsilon_2 u_{21})y + \delta_1(\epsilon_1 u_{12} + \epsilon_2 u_{22})z \\ + [\delta_1(\epsilon_1 \lambda_{11} + \epsilon_2 \lambda_{21}) + \delta_2 \epsilon_1] \rho + [\delta_1(\epsilon_1 \lambda_{12} + \epsilon_2 \lambda_{22}) + \delta_2 \epsilon_2] \sigma$$

will, for arbitrary values of $\epsilon_1, \epsilon_2, \delta_1, \delta_2$, represent an arbitrary point of H' . If we introduce again our special system of coördinates we have

$$x_1 = u_{11} \delta_1 \epsilon_1 + u_{21} \delta_1 \epsilon_2, \quad x_3 = \lambda_{11} \delta_1 \epsilon_1 + \lambda_{21} \delta_1 \epsilon_2 + \delta_2 \epsilon_1, \\ x_2 = u_{12} \delta_1 \epsilon_1 + u_{22} \delta_1 \epsilon_2, \quad x_4 = \lambda_{21} \delta_1 \epsilon_1 + \lambda_{22} \delta_1 \epsilon_2 + \delta_2 \epsilon_2,$$

as the coördinates of an arbitrary point of H' . If we eliminate $\delta_1, \delta_2, \epsilon_1, \epsilon_2$, we find the equation of H' :

$$(3) \quad (u_{22}x_1 - u_{21}x_2)[Jx_4 - (\lambda_{12}u_{22} - \lambda_{22}u_{12})x_1 + (\lambda_{12}u_{21} - \lambda_{22}u_{11})x_2] \\ + (u_{12}x_1 - u_{11}x_2)[Jx_3 - (\lambda_{11}u_{22} - \lambda_{21}u_{12})x_1 + (\lambda_{11}u_{21} - \lambda_{21}u_{11})x_2] = 0.$$

We shall mostly assume that P_y and P_z are the flecnodes of g , so that $u_{12} = u_{21} = 0$, and (3) simplifies into

$$(3a) \quad -\lambda_{12}u_{22}^2x_1^2 + \lambda_{21}u_{11}^2x_2^2 + u_{11}u_{22}(\lambda_{11} - \lambda_{22})x_1x_2 + u_{11}u_{22}(u_{22}x_1x_4 - u_{11}x_2x_3) = 0.$$

It is easy to see from (3a) that H' cannot coincide with H , unless S is a quadric.

The hyperboloids H and H' have the straight line g' in common. The rest of their intersection is therefore, in general, a space cubic. We shall call it the *derivative cubic*, and discuss some of its properties. It is interesting to notice that we obtain in this way associated with every ruled surface, a surface containing a single infinity of twisted cubics. This surface we shall also consider.

Let us again assume that P_y and P_z are the flecnodes of g . Then, it follows from (1) and (3a) that we may take

$$(4) \quad x_1 = tx_2, \quad x_2 = J(u_{11} - u_{22})t, \quad x_3 = tx_4, \\ x_4 = -\lambda_{12}u_{22}^2t^2 + u_{11}u_{22}(\lambda_{11} - \lambda_{22})t + \lambda_{21}u_{11}^2$$

as the parametric equations of the cubic. From these, the following corollaries follow at once. If $\lambda_{11} - \lambda_{22} = 0$, the derivative cubic, and therefore the hyperboloid H' , intersects g in two points, which are harmonic conjugates with respect to the flecnodes. If

$$(\lambda_{11} - \lambda_{22})^2 + 4\lambda_{12}\lambda_{21} = 0,$$

the cubic is tangent to g . The congruence Γ contains ∞^2 surfaces S' corresponding to each of these properties of the derivative cubic. For, the corresponding equations for η are again of the second order.

The equation of a plane which is tangent to H' at a point (x'_1, x'_2, x'_3, x'_4) , P_y and P_z being flecnodes, is

$$\begin{aligned}
 (5) \quad & [-2\lambda_{12}u_{22}^2x_1' + u_{11}u_{22}(\lambda_{11} - \lambda_{22})x_2' + u_{11}u_{22}^2x_4']x_1 \\
 & + [u_{11}u_{22}(\lambda_{11} - \lambda_{22})x_1' + 2\lambda_{21}u_{11}^2x_2' - u_{11}^2u_{22}x_3']x_2 \\
 & - u_{11}^2u_{22}x_2'x_3 + u_{11}u_{22}^2x_1'x_4 = 0.
 \end{aligned}$$

Consider the two points of g'' which correspond to P_y and P_z , viz.,

$$P_1'' = (u_{11}, 0, \lambda_{11}, \lambda_{12}), \quad P_2'' = (0, u_{22}, \lambda_{21}, \lambda_{22}).$$

The equations of the two planes tangent to S'' or H' at these points respectively, are

$$-u_{22}\lambda_{12}x_1 - u_{11}\lambda_{22}x_2 + u_{11}u_{22}x_4 = 0, \quad u_{22}\lambda_{11}x_1 + u_{11}\lambda_{21}x_2 - u_{11}u_{22}x_3 = 0.$$

They intersect g , i. e., the line $x_3 = x_4 = 0$, in two points

$$P_1 = (u_{11}\lambda_{22}, -u_{22}\lambda_{12}, 0, 0), \quad P_2 = (u_{11}\lambda_{21}, -u_{22}\lambda_{11}, 0, 0),$$

which coincide, if and only if $\lambda_{11}\lambda_{22} - \lambda_{12}\lambda_{21} = 0$, i. e., if and only if g'' intersects g , as is moreover geometrically evident. P_1 and P_2 are harmonic conjugates with respect to the flecnodes if $\lambda_{11}\lambda_{22} + \lambda_{12}\lambda_{21} = 0$, i. e., *under this condition the planes, tangent to the second derivative at the points which correspond to the flecnodes of g , divide these flecnodes harmonically.*

The line P_zP_p has, besides P_p , another point in common with H' . Its coördinates are found to be $(0, u_{22}, \lambda_{21}, 0)$. Similarly P_yP_p intersects H' in a second point $(u_{11}, 0, 0, \lambda_{12})$. Join these two points. The coördinates of any point on this line joining them will be $(\mu u_{11}, \lambda u_{22}, \lambda \lambda_{21}, \mu \lambda_{12})$, where $\lambda:\mu$ determines the position of the particular point. It is easily seen, by substituting in (3a), that this line is entirely on H' if $\lambda_{11} - \lambda_{22} = 0$, and in no other case, provided that S' is not developable. We have therefore the following result. *Corresponding to every point of S we have a point of S' . If each of the two flecnodes on a generator g of S be joined to the point of S' which corresponds to the other, the two straight lines thus obtained intersect the hyperboloid H' osculating S' along g' in two new points. The line joining these latter points lies entirely on H' , if H' intersects g in two points which are harmonic conjugates with respect to the flecnodes. The converse of the theorem is also true.*

Let us introduce the following abbreviations:

$$\begin{aligned}
 (6) \quad & A = 2u_{11}u_{22}(u_{11} - u_{22}), \quad \psi = Bt_1^2 + 2Ct_1t_2 + Dt_2^2, \\
 & B = v_{12}u_{22}, \quad C = \frac{1}{2}(u_{11}v_{22} - u_{22}v_{11}), \quad D = -v_{21}u_{11},
 \end{aligned}$$

and let us write the parameter t of the cubic curve in homogeneous form. Then we may write instead of (4),

$$\begin{aligned}
 (7) \quad & x_1 = At_1^2t_2, \quad x_3 = (Bt_1^2 + 2Ct_1t_2 + Dt_2^2)t_1 = \psi t_1, \\
 & x_2 = At_1t_2^2, \quad x_4 = (Bt_1^2 + 2Ct_1t_2 + Dt_2^2)t_2 = \psi t_2.
 \end{aligned}$$

If the cubic degenerates, each irreducible part will be a plane curve (a conic or a straight line). If therefore the cubic degenerates it must be possible to satisfy the equation

$$a_1 x_1 + a_2 x_2 + a_3 x_3 + a_4 x_4 = 0,$$

for all values of $t_1 : t_2$, the coefficients being independent of $t_1 : t_2$. If we substitute in this equation the values (7), and equate to zero the coefficients of t_1^3 , $t_1^2 t_2$, etc., we find

$$(8) \quad a_3 B = 0, \quad a_4 D = 0, \quad a_1 A + 2a_3 C + a_4 B = 0, \quad a_2 A + a_3 D + 2a_4 C = 0.$$

Let us assume first that neither B nor D vanishes. Then $a_3 = a_4 = 0$, and $A = 0$, for if A were not zero, we would have also $a_1 = a_2 = 0$, i. e., there would be no plane containing the (supposedly) degenerate cubic. But from $A = 0$ follows either $u_{11} - u_{22} = 0$, which would make S a quadric, or else u_{11} or u_{22} would vanish, which however contradicts the assumption that B and D shall not be zero.

Let us now assume $B \neq 0$, $D = 0$. Then $a_3 = 0$, and either u_{11} or v_{21} must vanish, i. e., either S' is developable or S has a straight line directrix. Similarly if $B = 0$, $D \neq 0$. Finally if $B = 0$, $D = 0$, either S has two straight line directrices, or else it has one while S' is developable. We have therefore the following theorem. *If the surface has one or more straight line directrices the derivative cubic always degenerates. In all other cases, the only way to obtain a degenerate derivative cubic consists in taking as derivative ruled surface of S , one of the developables of the congruence Γ .*

Another question at once suggests itself. To every value of x , i. e., to every generator of S there belongs a derivative cubic. In general, the cubics belonging to values of x , differing from each other by an infinitesimal δx , will not intersect. Their shortest distance will be an infinitesimal of the same order as δx . It may happen however that, for an appropriately chosen variable, this distance becomes infinitesimal of a higher order, or as we may say briefly, that consecutive cubics intersect. We ask now: is it possible to choose the independent variable in such a way that every pair of consecutive derivative cubics may intersect?

By putting $y = y_k$, $z = z_k$ ($k = 1, 2, 3, 4$) in

$$(9) \quad \phi = A t_1 t_2 (t_1 y + t_2 z) + \psi(t_1 \rho + t_2 \sigma),$$

we obtain the coördinates of any point P_ϕ on the cubic. As x changes we go from one cubic to another; as $t_1 : t_2$ changes we go from one point on a certain cubic to another point of the same curve. Equation (9) gives therefore, if both x and $t_1 : t_2$ be taken as variables, the locus of all such points P_ϕ , i. e., the surface generated by all of the derivative cubics of S . If $t_1 : t_2$ be chosen as a function of x , a curve is picked out upon this surface. Let us differentiate ϕ

totally, i. e., assuming that t_1 and t_2 are functions of x , and consider the quantity

$$\phi + \frac{d\phi}{dx} \delta x,$$

where δx is an infinitesimal. This will clearly represent the coördinates of a point on the adjacent derivative cubic determined by the parameters $x + \delta x$ and $t_k + dt_k/dx \delta x$. If the original cubic and this second one, infinitesimally close to it, intersect, it must be possible to choose t_k as functions of x in such a way, that the corresponding points of the two curves coincide up to infinitesimals of higher than the first order. Therefore $d\phi/dx$ must differ from a multiple of ϕ only by an infinitesimal quantity. Proceeding to the limit we must therefore have

$$(10) \quad \frac{d\phi}{dx} = \omega \phi,$$

We find by differentiation

$$(10a) \quad \begin{aligned} \frac{d\phi}{dx} = & [(t_1 t_2' + t_1' t_2) A + A' t_1 t_2] (t_1 y + t_2 z) \\ & + y [A t_1 t_2 (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) + \tfrac{1}{2} \psi u_{11} t_1] \\ & + z [A t_1 t_2 (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) + \tfrac{1}{2} \psi u_{22} t_2] \\ & + \rho [\tfrac{1}{2} A t_1^2 t_2 + \psi (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) + \psi' t_1] \\ & + \sigma [\tfrac{1}{2} A t_1 t_2^2 + \psi (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) + \psi' t_2], \end{aligned}$$

denoting as usual differentiation by strokes.

If we substitute in (10) we find the following four equations:

$$(11) \quad \begin{aligned} (a) \quad & [(t_1 t_2' + t_1' t_2) A + A' t_1 t_2] t_1 + A t_1 t_2 (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) \\ & + \tfrac{1}{2} \psi u_{11} t_1 = \omega A t_1^2 t_2, \\ (b) \quad & [(t_1 t_2' + t_1' t_2) A + A' t_1 t_2] t_2 + A t_1 t_2 (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) \\ & + \tfrac{1}{2} \psi u_{22} t_2 = \omega A t_1 t_2^2, \\ (c) \quad & \tfrac{1}{2} A t_1^2 t_2 + \psi (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) + \psi' t_1 = \omega \psi t_1, \\ (d) \quad & \tfrac{1}{2} A t_1 t_2^2 + \psi (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) + \psi' t_2 = \omega \psi t_2. \end{aligned}$$

If we multiply both members of (a) by t_2 , of (b) by $-t_1$, and add, and if we treat (c) and (d) in the same way, we find

$$\begin{aligned} (e) \quad & A t_1 t_2^2 (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) - A t_1^2 t_2 (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) \\ & + \tfrac{1}{2} \psi t_1 t_2 (u_{11} - u_{22}) = 0, \\ (f) \quad & \psi t_2 (t_1' - \tfrac{1}{2} p_{11} t_1 - \tfrac{1}{2} p_{21} t_2) - \psi t_1 (t_2' - \tfrac{1}{2} p_{12} t_1 - \tfrac{1}{2} p_{22} t_2) = 0. \end{aligned}$$

Let us assume first $\psi \neq 0$. Divide both members of (f) by ψ , and compare with (e). We find

$$\psi t_1 t_2 (u_{11} - u_{22}) = 0,$$

i. e., either S is a quadric, or else either t_1 or t_2 must vanish identically. Assume $t_1 = 0$. Then (e) is satisfied. From (f) we would find $p_{21} = 0$, for t_2 cannot be zero, since the ratio $t_1 : t_2$ is the parameter which determines a point on the cubic. But from $u_{21} = p_{21} = 0$ follows further $v_{21} = 0$, i. e., $D = 0$, so that in this case $\psi = Bt_1^2 + 2Ct_1 t_2 = 0$ contrary to our assumption. It is therefore impossible to satisfy (11) except by putting $\psi = 0$. According to (c) and (d) this gives either $t_1 = 0$, or $t_2 = 0$, or $A = 0$. Assume $A \neq 0$ and say $t_1 = 0$. All equations (11) are now satisfied. But from $t_1 = 0$, $\psi = 0$ follows $D = 0$, i. e., either u_{11} or v_{21} must vanish, i. e., either S has a straight line directrix or else S' is developable. If however $A = 0$, either S is a quadric or else S' is developable.

In connection with our last theorem, we may therefore say: *It is impossible to choose the surface S' of the congruence Γ in such a way that consecutive derivative cubics may intersect, except in the trivial cases when the cubics are degenerate.*

If in (9) we put $\psi = 0$, we obtain the locus of the intersections of the cubic with the generator of S to which it belongs, i. e., a certain curve cutting every generator twice. This may be an asymptotic curve. It is, in fact, if the further conditions

$$(12) \quad 2t'_1 - p_{11}t_1 - p_{21}t_2 = \omega t_1, \quad 2t'_2 - p_{12}t_1 - p_{22}t_2 = \omega t_2$$

are satisfied, where ω is arbitrary. For, as (10a) shows, the line joining P_ϕ to $P_{\phi+\phi'\delta x}$, i. e., the tangent to this curve, is then a generator of the second kind on the hyperboloid osculating S along g . In other words it is a tangent to an asymptotic curve of S .

In general, of course the conditions (12) and $\psi = 0$ can not both be satisfied at once. The question is: when are these conditions consistent? We find from (12)

$$t'_1 = \frac{1}{2}[(p_{11} + \omega)t_1 + p_{21}t_2], \quad t'_2 = \frac{1}{2}[p_{12}t_1 + (p_{22} + \omega)t_2].$$

Let us substitute these values of t'_1 and t'_2 in the equation $d\psi/dx = 0$. We shall find

$$\begin{aligned} [B' + B(p_{11} + \omega) + Cp_{12}]t_1^2 + [D' + D(p_{22} + \omega) + Cp_{21}]t_2^2 \\ + [2C' + Bp_{12} + Dp_{21} + C(p_{11} + p_{22} + 2\omega)]t_1 t_2 = 0, \end{aligned}$$

which compared with $\psi = 0$ gives,

$$B' + B(p_{11} + \omega) + Cp_{12} = \tau B,$$

$$D' + D(p_{22} + \omega) + Cp_{21} = \tau D,$$

$$2C' + Bp_{12} + Dp_{21} + C(p_{11} + p_{22} + 2\omega) = 2\tau C,$$

where τ is a proportionality factor. Eliminating τ , we find

$$(a) \ 2(B'C - BC') + BC(p_{11} - p_{22}) + (2C^2 - BD)p_{12} - B^2p_{21} = 0,$$

$$(13) \ (b) \ 2(D'C - DC') + DC(p_{22} - p_{11}) + (2C^2 - BD)p_{21} - D^2p_{12} = 0,$$

$$(c) \ BD' - B'D + BD(p_{11} - p_{22}) + C(Dp_{12} - Bp_{21}) = 0.$$

We may always assume that our system of differential equations has been so written that $p_{11} - p_{22} = 0$. Equations (13) may then be regarded as three homogeneous linear equations for $1, p_{12}, p_{21}$. Their determinant must therefore be zero. This gives rise to three alternatives: either $C^2 - BD$ or C or $BD' - B'D$ must vanish.

Consider first the case $C^2 - BD = 0$. Equations (13) become

$$(14) \quad 2(B'C - BC') - B^2p_{21} = 0, \quad 2(D'C - DC') - D^2p_{12} = 0,$$

$$BD' - B'D + CDp_{12} - CBp_{21} = 0.$$

If we multiply both members of the first two equations by CD and $-CB$ respectively, add and make use of $C^2 = BD$, we find

$$2C^2(DB' - D'B) + C^2(CDp_{12} - CBp_{21}) = 0,$$

whence, if $C \neq 0$,

$$-2(BD' - B'D) + CDp_{12} - CBp_{21} = 0.$$

But if we consider the last equation of (14), this gives $BD' - B'D = 0$, $Dp_{12} - Bp_{21} = 0$, which last equation may be written

$$p_{12}p_{21}(u_{11} - u_{22})^2 = 0,$$

i. e., the surface S has at least one straight line directrix.

In the second place let $C = 0$. Then (13) becomes

$$B(Dp_{12} + Bp_{21}) = 0, \quad D(Dp_{12} + Bp_{21}) = 0, \quad BD' - B'D = 0.$$

According to the first two equations S must either have at least two straight line directrices, or else $u_{11} + u_{22}$ must be zero. The third condition gives, on substituting the values of B and D ,

$$\frac{u'_{22}}{u_{22}} - \frac{u'_{11}}{u_{11}} + \frac{v'_{12}}{v_{12}} - \frac{v'_{21}}{v_{21}} = 0,$$

if we assume that S has no straight line directrix, so that v_{12} and v_{21} are not

zero. By integration we find

$$\frac{v_{12}}{v_{21}} \frac{u_{22}}{u_{11}} = \text{const.},$$

whence, since $u_{11} + u_{22} = 0$ in this case,

$$\frac{p_{12}}{p_{21}} = \text{const.}$$

But if we substitute into the condition $\Delta = 0$ for a surface belonging to a linear complex the assumptions $u_{12} = u_{21} = p_{11} - p_{22} = 0$, which we have made, we find either $u_{11} - u_{22} = 0$, i. e., S is a quadric, or $p_{12}p'_{21} - p'_{12}p_{21} = 0$, i. e., $p_{12}/p_{21} = \text{const.}$ Therefore we see that if $C = 0$, S either has straight line directrices, or at least belongs to a linear complex. Moreover in that case the independent variable must be so chosen as to make $u_{11} + u_{22}$ vanish.

If finally $BD' - B'D = 0$, we find from (13) either $C = 0$, which leads us back to the case just considered, or else $Dp_{12} - Bp_{21}$ must vanish, which gives again a surface S with at least one straight line directrix.

In all of these cases the surface S belongs to a linear complex. In the special cases the asymptotic lines thus obtained are merely two of the straight line directrices, if there are two, or else one straight line directrix and another curved asymptotic line obtained by putting $u_{11} + u_{22} = 0$. If we leave aside the trivial cases, we can say that *if a ruled surface belongs to a linear complex, and if the independent variable is so chosen that $u_{11} + u_{22} = 0$, the surface of derivative cubics determines an asymptotic curve upon it which intersects every generator twice.*

We have seen (*Congruence*, p. 197), that there exists in the congruence Γ a single infinity of ruled surfaces S' for which $u_{11} + u_{22} = 0$. They are those ruled surfaces of Γ whose intersections with the flecnodal surface of S are asymptotic lines upon them. *But we can also characterize them by saying that such a surface is made up of the lines of Γ intersecting any asymptotic curve on the flecnodal surface of S .* In fact, if we assume $u_{12} = u_{21} = 0$, $p_{11} = p_{22} = 0$, we may write the equations of the sheet F' of the flecnodal surface as follows:*

$$(15) \quad \begin{aligned} y'' - 2 \frac{q_{12}}{p_{12}} y' - \rho' - q_{11} y + \frac{q_{12}}{p_{12}} \rho &= 0, \\ \rho'' + [2(q_{11} + q_{22}) - p_{12}p_{21}]y' - 2 \frac{q_{12}}{p_{12}} \rho' + [2q'_{11} - p_{12}q_{21} - 4 \frac{q_{12}}{p_{12}} q_{11}]y - q_{22}\rho &= 0. \end{aligned}$$

If $u_{11} + u_{22} = 0$, the coefficient of y' in the second equation vanishes, which proves that the curve C_p is an asymptotic curve on F' . *The asymptotic curves*

* The system of equations for the flecnodal surface, in *Congruence*, p. 190, equ. (12), does not agree exactly with (15) because there the further assumption $u_{11} = 0$ was made. This should have been stated there.

on F' and F'' must therefore correspond to each other. The congruence Γ is therefore a so-called W -congruence.*

We may now state our theorem as follows: *In order that the surface of derivative cubics may intersect the ruled surface S in an asymptotic curve, S must belong to a linear complex. Moreover, its derivative ruled surface must intersect the flecnodal surface of S along an asymptotic curve.*

The asymptotic curve on S , thus determined, is unique. For the ratio $B:D$ which determines it cannot be changed by any transformation of the variables subject to the conditions $p_{11} - p_{22} = u_{12} = u_{21} = u_{11} + u_{22} = 0$. We may then take any asymptotic curve of the flecnodal surface and consider the ruled surface of Γ made up of the lines which intersect it. We obtain thus as a consequence a single infinity of surfaces made up of derivative cubics. All of these intersect S along the same asymptotic curve.

But we notice further that $C = 0$. Therefore, *this asymptotic curve intersects every generator in two points which are harmonic conjugates with respect to the flecnodes.* We shall meet this special asymptotic curve again in a later paragraph.

Another question suggests itself. Is it possible to choose the independent variable of our system in such a way that the derivative cubics shall be asymptotic lines on the surface generated by them?

In order to answer this question, we must first find the coördinates of the osculating plane of the cubic at any one of its points. For the moment we prefer to take the equation of the cubic referred to a non-homogeneous parameter t , i. e., in the form

$$x_1 = At^2, \quad x_2 = At, \quad x_3 = Bt^3 + 2Ct^2 + Dt, \quad x_4 = Bt^2 + 2Ct + D.$$

Its intersections with the plane $\sum u_i x_i = 0$, whose coördinates are (u_1, \dots, u_4) , will be given by solving the cubic equation

$$Bu_3 t^3 + (Au_1 + 2Cu_3 + Bu_4)t^2 + (Au_2 + 2Cu_4 + Du_3)t + Du_4 = 0.$$

The three roots of this cubic must coincide if the plane is an osculating plane of the cubic curve. They must therefore also satisfy the equations obtained from the above by twofold differentiation with respect to t . This gives the following conditions:

$$\begin{aligned} 3Bu_3 t + Au_1 + 2Cu_3 + Bu_4 &= 0, \\ (Au_1 + 2Cu_3 + Bu_4)t + Au_2 + 2Cu_4 + Du_3 &= 0, \\ (Au_2 + 2Cu_4 + Du_3)t + 3Du_4 &= 0. \end{aligned}$$

Of course, only the ratios of $u_1 \dots u_4$ are of interest, so that we may multiply $u_1 \dots u_4$ by a common factor if we please. We find

* Cf. BIANCHI-LUKAT, *Vorlesungen über Differentialgeometrie*, p. 315.

$$(16) \quad \begin{aligned} u_1 &= B^2 t^3 - 3BDt - 2CD, & u_3 &= AD, \\ u_2 &= 2BCt^3 + 3BDt^2 - D^2, & u_4 &= -ABt^3, \end{aligned}$$

as the coördinates of the plane osculating the derivative cubic belonging to the generator g at the point whose parameter is t , or in homogeneous form,

$$(17) \quad \begin{aligned} u_1 &= B^2 t_1^3 - 3BDt_1 t_2^2 - 2CDt_2^3, & u_3 &= ADt_2^3, \\ u_2 &= 2BCt_1^3 + 3BDt_1^2 t_2 - D^2 t_2^3, & u_4 &= -ABt_1^3. \end{aligned}$$

We have already made use of the expression

$$\phi = At_1 t_2 (t_1 y + t_2 z) + \psi(t_1 \rho + t_2 \sigma)$$

as giving the coördinates of a point $(t_1 : t_2)$ of the cubic curve which belongs to the argument x , or as giving the coördinates of a point $(x, t_1 : t_2)$ of the surface formed by the aggregate of all of these curves. The plane which is tangent to this surface at the point $(x, t_1 : t_2)$ must contain also the point $t_1 : t_2$ of the adjacent cubic, i. e., the point whose coördinates are given by

$$\phi + \frac{\partial \phi}{\partial x} \delta x,$$

where in forming $\partial \phi / \partial x$, x , t_1 and t_2 are regarded as independent variables. The tangent plane must therefore contain the point whose coördinates are $\partial \phi / \partial x$. We have

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= y [A' t_1^2 t_2 - \tfrac{1}{2} A t_1 t_2 (p_{11} t_1 + p_{21} t_2) + \tfrac{1}{2} \psi u_{11} t_1] \\ &\quad + z [A' t_1 t_2^2 - \tfrac{1}{2} A t_1 t_2 (p_{12} t_1 + p_{22} t_2) + \tfrac{1}{2} \psi u_{22} t_2] \\ &\quad + \rho [\tfrac{1}{2} A t_1^2 t_2 - \tfrac{1}{2} \psi (p_{11} t_1 + p_{21} t_2) + \frac{\partial \psi}{\partial x} t_1] \\ &\quad + \sigma [\tfrac{1}{2} A t_1 t_2^2 - \tfrac{1}{2} \psi (p_{12} t_1 + p_{22} t_2) + \frac{\partial \psi}{\partial x} t_2]. \end{aligned}$$

The point, whose coördinates (ξ_1, \dots, ξ_4) are the coefficients of y, z, ρ, σ in this expression, must be in the tangent plane of the point $(x, t_1 : t_2)$. If therefore, the cubic curve is an asymptotic line upon the surface, its osculating plane must contain the point (ξ_1, \dots, ξ_4) , i. e., we must have

$$(18) \quad u_1 \xi_1 + u_2 \xi_2 + u_3 \xi_3 + u_4 \xi_4 = 0.$$

We find in the first place

$$(19) \quad \begin{aligned} \xi_1 &= u_{11} B t_1^3 + (2A' + 2u_{11} C - A p_{11}) t_1^2 t_2 + (u_{11} D - A p_{21}) t_1 t_2^2, \\ \xi_2 &= u_{22} D t_2^3 + (u_{22} B - A p_{12}) t_1^2 t_2 + (2A' + 2u_{22} C - A p_{22}) t_1 t_2^2, \end{aligned}$$

$$\begin{aligned}
 \xi_3 &= (2B' - Bp_{11})t_1^3 + (A - Bp_{21} + 4C' - 2Cp_{11})t_1^2 t_2 \\
 &\quad + (2D' - Dp_{11} - 2Cp_{21})t_1 t_2^2 - Dp_{21} t_2^3, \\
 (19) \quad \xi_4 &= -Bp_{12} t_1^3 + (2B' - Bp_{22} - 2Cp_{12})t_1^2 t_2 \\
 &\quad + (A - Dp_{12} + 4C' - 2Cp_{22})t_1 t_2^2 + (2D' - Dp_{22})t_2^3.
 \end{aligned}$$

If these values, and the values (17) for u_1, \dots, u_4 be substituted in (18), and the coefficients of $t_1^6, t_1^5 t_2$, etc., be successively equated to zero, the following seven equations make their appearance:

$$\begin{aligned}
 (20) \quad &u_{11} B^3 + AB^2 p_{12} = 0, \\
 &B^2(2A' + 2u_{11}C - Ap_{11}) + 2BC(u_{22}B - Ap_{12}) \\
 &\quad - AB(2B' - Bp_{22} - 2Cp_{12}) = 0, \\
 &B^2(u_{11}D - Ap_{21}) - 3B^2Du_{11} + 2BC(2A' + 2u_{22}C - Ap_{22}) \\
 &\quad + 3BD(u_{22}B - Ap_{12}) - AB(A - Dp_{12} + 4C' - 2Cp_{22}) = 0, \\
 &-3BD(2A' + 2u_{11}C - Ap_{11}) - 2BCD(u_{11} - u_{22}) \\
 &\quad + 3BD(2A' + 2u_{22}C - Ap_{22}) \\
 &\quad + AD(2B' - Bp_{11}) - AB(2D' - Dp_{22}) = 0, \\
 &-3BD(u_{11}D - Ap_{21}) - 2CD(2A' + 2u_{11}C - Ap_{11}) \\
 &\quad + 3BD^2u_{22} - D^2(u_{22}B - Ap_{12}) + AD(A - Bp_{21} + 4C' - 2Cp_{11}) = 0, \\
 &-2CD(u_{11}D - Ap_{21}) - D^2(2A' + 2u_{22}C - Ap_{22}) \\
 &\quad + AD(2D' - Dp_{11} - 2Cp_{21}) = 0, \\
 &-u_{22}D^3 - AD^2p_{21} = 0.
 \end{aligned}$$

Let us assume that both B and D are different from zero. Then the first and last equations give

$$u_{11}u_{22}v_{12} + 2u_{11}u_{22}(u_{11} - u_{22})p_{12} = 0, \quad u_{11}u_{22}v_{21} - 2u_{11}u_{22}(u_{11} - u_{22})p_{21} = 0,$$

or

$$u_{11}u_{22}(u_{11} - u_{22})p_{12} = u_{11}u_{22}(u_{11} - u_{22})p_{21} = 0.$$

But all of the possibilities here suggested give zero values to either B or D , or both, except $u_{11} - u_{22} = 0$, in which case S is a quadric. But this is only apparently an exception, arising from the fact that in this case the flecnodal curve is indeterminate. If two of the straight lines of the second set be taken for the curves C_y and C_z , we have in this case also $B = D = 0$.

There remain two possibilities. If $B = D = 0$ all of the equations (20) are

satisfied. If S' is not developable, S must then belong to a linear congruence. If S' is developable, it is sufficient for S to have one straight line directrix. In either case the derivative cubic degenerates into a straight line, and is therefore obviously an asymptotic curve upon the surface of cubics.

Finally, it might happen that one only of the two quantities B and D is zero. Say $B = 0$, $D \neq 0$. Then the first four equations of (20) are satisfied. The other three become

$$\begin{aligned} -4u_{11}C^2 + ADp_{12} + A^2 &= 0, \\ -2Cu_{11}D - D(2A' + 2u_{22}C - Ap_{22}) + A(2D' - Dp_{11}) &= 0, \\ -u_{22}D - Ap_{21} &= 0. \end{aligned}$$

From the last of these we find

$$u_{11}u_{22}(u_{11} - u_{22})p_{21} = 0,$$

whence, since $D \neq 0$, follows $u_{22} = 0$, and therefore $A = 0$. The second equation now gives us $C = 0$, which also satisfies the first. We have again S' a developable surface, so that the cubic degenerates. Therefore

The derivative cubics of a ruled surface are asymptotic curves, upon the surface formed by their totality, only in the trivial cases when they degenerate into straight lines.

§ 2. Null-system of the derivative cubic.

A twisted cubic always determines a null-system, i. e., a point-to-plane correspondence with incident elements. Geometrically this correspondence may be set up as follows. An arbitrary plane intersects the curve in three points. The three planes, which osculate the curve in these points, intersect again in a point which is situated in the original plane. This is the point which corresponds to the plane in the null-system of the cubic.

We shall now set up the equations for this null-system. For this purpose it is more convenient to use the equations of the cubic referred to a non-homogeneous parameter t .

Let t_1, t_2, t_3 be the three values of t which correspond to the three points in which the plane

$$v_1x_1 + v_2x_2 + v_3x_3 + v_4x_4 = 0$$

intersects the cubic. Then t_1, t_2, t_3 are the roots of the cubic equation

$$Bv_3t^3 + (Av_1 + 2Cv_3 + Bv_4)t^2 + (Av_2 + Dv_3 + 2Cv_4)t + Dv_4 = 0.$$

Therefore we shall have

$$\frac{Av_1 + 2Cv_3 + Bv_4}{Bv_3} = -(t_1 + t_2 + t_3),$$

$$\frac{Av_2 + Dv_3 + 2Cv_4}{Bv_3} = t_2 t_3 + t_3 t_1 + t_1 t_2,$$

$$\frac{Dv_4}{Bv_3} = -t_1 t_2 t_3.$$

If we solve these equations for v_1/v_3 , v_2/v_3 , v_4/v_3 , then make them homogeneous, and multiply $v_1 \cdots v_4$ by the common factor A , we shall find

$$(21) \quad \begin{aligned} v_1 &= -2CD + B^2 t_1 t_2 t_3 - BD(t_1 + t_2 + t_3), & v_3 &= AD, \\ v_2 &= -D^2 + 2BCt_1 t_2 t_3 + BD(t_2 t_3 + t_3 t_1 + t_1 t_2), & v_4 &= -ABt_1 t_2 t_3. \end{aligned}$$

The coördinates of the planes which osculate the curve in the three points t_1 , t_2 , t_3 , are, according to (16),

$$\begin{aligned} u_1^{(k)} &= B^2 t_k^3 - 3BDt_k - 2CD, & u_3^{(k)} &= AD, \\ u_2^{(k)} &= 2BCt_k^3 + 3BDt_k^2 - D^2, & u_4^{(k)} &= -ABt_k^3 \end{aligned} \quad (k=1, 2, 3).$$

Let x_1, \dots, x_4 be the coördinates of the point of intersection of these three planes. We must then have

$$\sum_{i=1}^4 u_i^{(k)} x_i = 0 \quad (k=1, 2, 3).$$

Solving these equations we find

$$(22) \quad \begin{aligned} x_1 &= A(t_2 t_3 + t_3 t_1 + t_1 t_2), \\ x_2 &= A(t_1 + t_2 + t_3), \\ x_3 &= 3Bt_1 t_2 t_3 + 2C(t_2 t_3 + t_3 t_1 + t_1 t_2) + D(t_1 + t_2 + t_3), \\ x_4 &= B(t_2 t_3 + t_3 t_1 + t_1 t_2) + 2C(t_1 + t_2 + t_3) + 3D, \end{aligned}$$

and a simple calculation will show that $\sum v_i x_i = 0$, i. e., as we have stated, the point of intersection of the three osculating planes lies in the plane of the three points of osculation.

In our null-system then, the plane (21) and the point (22) correspond to each other. To find the explicit equations for this correspondence, we need only eliminate t_1 , t_2 , t_3 between equations (21) and (22). Denoting by ω and ω' two proportionality factors, we find:

$$(23) \quad \begin{aligned} \omega v_1 &= * & + 4(C^2 - BD)x_2 & + ABx_3 - 2ACx_4, \\ \omega v_2 &= -4(C^2 - BD)x_1 & + * & + 2ACx_3 - ADx_4, \\ \omega v_3 &= -ABx_1 & - 2ACx_2 & + * + A^2 x_4, \\ \omega v_4 &= 2ACx_1 & + ADx_2 & - A^2 x_3 + *; \end{aligned}$$

$$\begin{aligned}
 \omega'x_1 &= * + A^2v_2 + ADv_3 + 2ACv_4, \\
 \omega'x_2 &= -A^2v_1 + * - 2ACv_3 - ABv_4, \\
 \omega'x_3 &= -ADv_1 + 2ACv_2 + * + 4(C^2 - BD)v_4, \\
 \omega'x_4 &= -2ACv_1 + ABv_2 - 4(C^2 - BD)v_3 + *.
 \end{aligned}
 \tag{24}$$

But associated with the null-system we have a linear complex, made up of all of the lines passing through a point, which lie at the same time in the plane corresponding to this point in the null-system. Introduce line coördinates by putting $\omega_{ik} = x_i y_k - x_k y_i$, $\omega'_{ik} = v_i w_k - v_k w_i$, where $x_1 \dots x_4$ and $y_1 \dots y_4$ are coördinates of two points on the line, and where $v_1 \dots v_4$, $w_1 \dots w_4$ are the coördinates of two planes containing the line. Then the equation of the complex may be written in either of the forms

$$(25) \quad -4(C^2 - BD)\omega_{12} - AB\omega_{13} + 2AC\omega_{14} - 2AC\omega_{23} - AD\omega_{42} - A^2\omega_{34} = 0,$$

$$\begin{aligned}
 (26) \quad & -4(C^2 - BD)\omega'_{34} - AB\omega'_{42} + 2AC\omega'_{23} \\
 & - 2AC\omega'_{14} - AD\omega'_{13} - A^2\omega'_{12} = 0.
 \end{aligned}$$

This complex becomes special if $A^2BD = 0$, i. e., only if the cubic degenerates.

To the flecnode $P_y(x_2 = x_3 = x_4 = 0)$ corresponds the plane

$$[0, -4(C^2 - BD), -AB, 2AC].$$

Therefore, if $C^2 - BD = 0$, i. e., if the derivative cubic is tangent to the generator g , the corresponding plane passes through g . If $C = 0$, i. e., $u_{11}/u_{22} = \text{const.}$ the plane passes through $P_\sigma P_y$. Therefore, if the intersections of the cubic with g and the flecnodes form a harmonic group on g , the plane corresponding to each flecnode passes through that point of the derived ruled surface which corresponds to the other.

§ 3. The osculating linear complex.

There is another linear complex associated with every generator of a ruled surface, even more important than the one just considered. A linear complex is determined by five of its lines, provided that these have no two straight line intersectors. Let us consider five generators of a ruled surface, g and four others. As the four other generators approach coincidence with g , a definite linear complex will in general be obtained as a limit. We shall speak of it as the linear complex osculating S along g .

Instead of determining the complex by five consecutive generators of S , it will be advisable to determine it by means of two pairs of lines which are reciprocal polars with respect to it. Two such pairs are obviously constituted by

the flecnode tangent of g and of another generator infinitesimally close to g . Let us denote by g_x, f'_x, f''_x, g'_x respectively the generator of S , the first and the second flecnode tangents and the generator of S' which belong to the argument x . Similarly we denote by $g_{x+\delta x}$, etc., the corresponding lines belonging to the argument $x + \delta x$, where δx is an infinitesimal.

Clearly f'_x and f''_x are the directrices of the osculating linear congruence, (determined by four consecutive generators). Therefore all lines intersecting f'_x and f''_x belong to the osculating linear complex, which must therefore be of the form

$$(27) \quad a\omega_{13} + b\omega_{42} = 0.$$

Change the parameter x by an infinitesimal quantity δx . The flecnode tangents $f'_{x+\delta x}$ and $f''_{x+\delta x}$ must again be the directrices of a linear congruence contained in the complex. We have

$$y_{x+\delta x} = y_x + y'_x \delta x, \quad \rho_{x+\delta x} = \rho_x + \rho'_x \delta x, \text{ etc.}$$

If we substitute the values of y', ρ' , etc., from *Covariants*, equations (21) and (33), we shall find

$$\begin{aligned} 2y_{x+\delta x} &= 2y + (\rho - p_{11}y - p_{12}z) \delta x, \\ 2z_{x+\delta x} &= 2z + (\sigma - p_{21}y - p_{22}z) \delta x, \\ 2\rho_{x+\delta x} &= 2\rho + [u_{11}y + u_{12}z - p_{11}\rho - p_{12}\sigma] \delta x, \\ 2\sigma_{x+\delta x} &= 2\sigma + [u_{21}y + u_{22}z - p_{21}\rho - p_{22}\sigma] \delta x, \end{aligned}$$

where of course u_{12} and u_{21} may be equated to zero, since C_y and C_z constitute the flecnode curve.

Now clearly, the coefficients of y, z, ρ, σ in the expressions

$$\rho_{x+\delta x} + \lambda y_{x+\delta x} \quad \text{and} \quad \sigma_{x+\delta x} + \mu z_{x+\delta x}$$

will be the coördinates of two arbitrary points P_1 and P_2 , situated on $f'_{x+\delta x}$ and $f''_{x+\delta x}$ respectively. If (27) is the equation of the osculating complex, the plane which corresponds in it to P_1 must contain P_2 for arbitrary values of λ and μ . This consideration will enable us to determine the ratio $a : b$.

We find first, remembering that $u_{12} = u_{21} = 0$, for P_1 and P_2 the coördinates :

	x_1	x_2	x_3	x_4
P_1	$2\lambda + (u_{11} - \lambda p_{11})\delta x$	$-\lambda p_{12}\delta x$	$2 + (-p_{11} + \lambda)\delta x$	$-p_{12}\delta x$
P_2	$-\mu p_{21}\delta x$	$2\mu + (u_{22} - \mu p_{22})\delta x$	$-p_{21}\delta x$	$2 + (-p_{22} + \mu)\delta x$

But, if we denote by u_1, \dots, u_4 the coördinates of the plane which corresponds to the point x_1, \dots, x_4 in the linear complex (27), we find

$$u_1 = -ax_3, \quad u_2 = +bx_4, \quad u_3 = ax_1, \quad u_4 = -bx_2.$$

Substituting for x_1, \dots, x_4 the coördinates of P_1 , and writing down the condition that P_2 shall lie in the plane corresponding to P_1 , we find that we must have

$$a(\mu - \lambda)p_{21} + b(\lambda - \mu)p_{12} = 0$$

for arbitrary values of λ and μ , i. e., $a:b = p_{12}:p_{21}$.

Therefore, *the equation of the osculating linear complex, in the system of coördinates here employed, is*

$$(28) \quad p_{12}\omega_{13} + p_{21}\omega_{42} = 0.$$

The point-plane correspondence, determined by this complex, is given by the equations

$$(29) \quad u_1 = p_{12}x_3, \quad u_2 = -p_{21}x_4, \quad u_3 = -p_{12}x_1, \quad u_4 = p_{21}x_2.$$

Let us consider a point on the generator g . There will correspond to it, in this complex, a plane, obviously containing the generator itself. But to every point of g there also corresponds another plane through g , viz., the plane tangent to the ruled surface at that point. Clearly, there will exist in general two points on g at which these two planes will coincide. We shall call them the *complex points* of g , and their locus on S , the *complex curve* of the surface. We proceed to determine the complex points of g .

The plane corresponding to any point of g , $(x_1, x_2, 0, 0)$, in the osculating linear complex, has the coördinates

$$(30) \quad u_1 = 0, \quad u_2 = 0, \quad u_3 = -p_{12}x_1, \quad u_4 = p_{21}x_2.$$

The coördinates of the plane, tangent to S at the same point, are found most easily by computing the equation of the plane tangent to the osculating hyperboloid H at that point. They are $(0, 0, x_2, -x_1)$. This plane and (30) coincide if and only if $-p_{12}x_1:p_{21}x_2 = x_2:-x_1$, i. e., if

$$(31) \quad p_{12}x_1^2 - p_{21}x_2^2 = 0.$$

This shows that *the complex points and the flecnodes form a harmonic group on every generator of the surface.*

If in (7) we put $u_{11} + u_{22} = 0$, we find that the derivative cubic intersects g precisely in the complex points. Therefore: *if the surface S' of the congruence Γ is so chosen that it intersects the flecnodal surface of S in an asymptotic curve, the surface of derivative cubics will intersect S along its complex curve.*

If S is contained in a linear complex, the complex curve is at the same time an asymptotic curve.

For, we notice in the first place that under our assumptions the factors of the expression $p_{12}z^2 - p_{21}y^2$ determine the complex points. Let us assume $p_{11} - p_{22} = 0$, which we may do without affecting the generality of our argument. Then the condition $\Delta = 0$ for a linear complex becomes $p_{12}/p_{21} = \text{const.}$ If we now make the transformation

$$\eta = \sqrt{p_{21}}y + \sqrt{p_{12}}z, \quad \zeta = \sqrt{p_{21}}y - \sqrt{p_{12}}z,$$

we find that in the transformed system of differential equations, $\pi_{12} = \pi_{21} = 0$, the coefficients of this transformed system being denoted by Greek letters. But this proves that C_η and C_ζ are asymptotic lines on S . It is geometrically evident that the tangents of this asymptotic curve will be lines of the linear complex.

LIE seems to have been the first to notice the existence of this special asymptotic curve on a ruled surface belonging to a linear complex. He noticed in 1871 that its determination requires no integration, and that all other asymptotic curves can be obtained by quadratures.* These latter remarks we can also verify at once from our theory. PICARD found the same theorems independently in 1877.† These theorems on the determination of all of the asymptotic curves by quadratures if one of them is known, follow at once from the fact first noted by BONNET that their equation is of the RICCATI form, and had already been explicitly formulated and applied to special surfaces by CLEBSCH.‡ It seems that VOSS§ was the first to notice that this asymptotic curve and the flecnodal curve divide the generators of the surface harmonically. CREMONA|| however had already observed this in the special case of a surface with two straight line directrices. The general notion of the complex curve, its relation to the derivative surface and to the surface of derivative cubics do not seem to occur in the literature of the subject.

There always exists a pair of points harmonically conjugate to each of two given pairs. We see easily that the pair

$$(32) \quad p_{21}y^2 + p_{12}z^2$$

is thus situated with respect to the flecnodes as well as the complex points. They are therefore the double points of an involution of which the flecnodes

* LIE, *Verhandl. d. Ges. d. Wiss. Christiania* (1871), *Mathematische Annalen*, vol. 5 (1872).

† PICARD, *Thèse*, Paris (1877). See also DARBOUX, *Bulletin* (1877), p. 335, and *Annales de l'Ecole Normale* (1877).

‡ CLEBSCH, *Crelle's Journal*, vol. 68.

§ VOSS, *Mathematische Annalen*, vol. 8 (1875).

|| CREMONA, *Annali di Matematiche* (1867-68).

and the complex points are two pairs. We will call them the *involute points*, their locus the *involute curve*. Of course these three pairs cannot be real at the same time.

Consider the expression

$$(33) \quad \gamma = \theta_4 E - \theta'_4 C,$$

where

$$C = u_{12}z^2 - u_{21}y^2 + (u_{11} - u_{22})yz, \quad E = v_{12}z^2 - v_{21}y^2 + (v_{11} - v_{22})yz.$$

It is easy to prove that γ is a covariant. Moreover it reduces to (32) under our special assumptions. Therefore, the factors of the covariant $\theta_4 E - \theta'_4 C$ give the expressions for the involute points in invariant form. If

$$(34) \quad \theta'_4 u_{12} - \theta_4 v_{12} = 0, \quad \theta'_4 u_{21} - \theta_4 v_{21} = 0,$$

the combined locus of C_y and C_z constitutes the involute curve. If however

$$(35) \quad u_{11} - u_{22} = 0, \quad v_{11} - v_{22} = 0,$$

then C_y and C_z together constitute the complex curve.

We can also write down a covariant whose factors give directly the complex points. It is found by writing down the conditions that a quadratic in y and z shall represent points which divide the pair of the flecnodes, and the pair of involute points harmonically. We find in this way, that the factors of

$$(36) \quad [(u_{11} - u_{22})v_{12} - (v_{11} - v_{22})u_{12}]z^2 + [(u_{11} - u_{22})v_{21} - (v_{11} - v_{22})u_{21}]y^2 + 2[u_{12}v_{21} - u_{21}v_{12}]yz$$

represent the complex points. One of the covariants (33) or (36) should be added to the list (*Covariants*, p. 433), which is incomplete as it stands.

We can easily show that the derivative cubic cannot intersect g in its involute points unless S is a quadric. Moreover if the derivative cubic intersects g in two points which are harmonic conjugates with respect to the complex point, S can only be a quadric. The cubic will however intersect g in two points harmonic conjugates with respect to the involute points, provided that $u_{11} + u_{22} = 0$, i. e., provided that S' intersects the flecnode surface of S in an asymptotic curve. It then passes through the complex points.

It is further clear, geometrically as well as analytically, that the two complex points, as well as the two involute points, can coincide only if S has a straight line directrix or if the flecnodes coincide. They become indeterminate if S has two straight line directrices.

To every point P' of g' there corresponds a plane in the osculating linear complex, as well as the plane tangent to S' at P' . When do these planes coincide?

Let the coördinates of P' be $(0, 0, \alpha_1, \alpha_2)$. The plane, corresponding to P' in the complex, has the coördinates $(\alpha_1 p_{12}, -\alpha_2 p_{21}, 0, 0)$, so that it contains g' . The coördinates of the plane tangent to S' at P' are of course the same as those of the plane tangent to H' at P' , which may be obtained from (5), viz: $(u_{11} u_{22}^2 \alpha_2, -u_{11}^2 u_{22} \alpha_1, 0, 0)$. These planes coincide if and only if

$$u_{11} p_{12} \alpha_1^2 = u_{22} p_{21} \alpha_2^2.$$

The corresponding points on g are again harmonic conjugates with respect to the flecnodes, i. e., those asymptotic tangents of S which join g to the points of g' , the planes corresponding to which in the linear complex are the planes tangent to S' , are harmonic conjugates with respect to the flecnodal tangents.

The planes which correspond to these two points of g' in the null-system of the cubic, do not contain g' .

§ 4. *Relation of the osculating linear complex to the linear complex of the derivative cubic.*

The equations of the two complexes are

$$\begin{aligned} \Omega_1 &= 4(C^2 - BD)\omega_{12} + AB\omega_{13} + AD\omega_{42} \\ (37) \quad &+ A^2\omega_{34} + 2AC\omega_{23} - 2AC\omega_{14} = 0, \\ \Omega_2 &= p_{12}\omega_{13} + p_{21}\omega_{42} = 0. \end{aligned}$$

Their simultaneous invariant is

$$(38) \quad -2u_{11}u_{22}(u_{11} - u_{22})^2(u_{11} + u_{22})p_{12}p_{21},$$

which, leaving aside the cases when S' is developable or when S has one or more straight line directrices, vanishes if and only if $u_{11} + u_{22} = 0$. Therefore, *the osculating linear complex and the complex of the derivative cubic are in involution if the first derivative ruled surface cuts out asymptotic curves on the flecnodal surface of S , and the cubic passes through the complex points of g .* Some of our previous theorems are consequences of this.

The two special complexes which are contained in the family

$$\lambda\Omega_1 + \mu\Omega_2 = 0,$$

where λ and μ are constants, are those for which

$$-3A^2BD\lambda^2 - 2u_{11}u_{22}(u_{11} - u_{22})^2(u_{11} + u_{22})p_{12}p_{21}\lambda\mu + p_{12}p_{21}\mu^2 = 0,$$

or, discarding again the case when S has a straight directrix,

$$-12u_{11}^3u_{22}^3(u_{11} - u_{22})^4\lambda^2 - 2u_{11}u_{22}(u_{11} - u_{22})^2(u_{11} + u_{22})\lambda\mu + \mu^2 = 0.$$

They coincide if

$$u_{11}^2u_{22}^2(u_{11} - u_{22})^4(u_{11} + u_{22})^2 + 12u_{11}^3u_{22}^3(u_{11} - u_{22})^4 = 0,$$

i. e., if S' is developable, if S is a quadric, or if

$$(39) \quad (u_{11} + u_{22})^2 + 12u_{11}u_{22} = 0.$$

We can always choose the independent variable so as to satisfy this condition. In fact, if we change the independent variable by putting $\xi = \xi(x)$, according to *Congruence*, equation (5), we shall have

$$(\bar{u}_{11} + \bar{u}_{22})^2 + 12\bar{u}_{11}\bar{u}_{22} = 0,$$

if ξ be taken as any solution of the equation

$$(40) \quad 64\mu^2 + 32(u_{11} + u_{22})\mu + (u_{11} + u_{22})^2 + 12u_{11}u_{22} = 0,$$

where

$$\mu = \{ \xi, x \} = \eta' - \frac{1}{2}\eta^2, \quad \eta = \frac{\xi''}{\xi'}.$$

Therefore, there exist two families of ∞^1 non-developable ruled surfaces in the congruence Γ such that the linear congruence, common to the osculating linear complex of S and the linear complex of the derivative cubic, shall have coincident directrices. Any four surfaces of one family intersect all of the asymptotic tangents of S in a point row of constant anharmonic ratio. The two families never coincide unless $\theta_4 = 0$, i. e., unless the flecnodal curve intersects every generator in two coincident points. But in this case the congruence is not defined. If S has a straight line directrix this congruence is degenerate.

The coördinates of the plane, which corresponds to a point $(x_1, x_2, 0, 0)$ of g in the null-system of the cubic, are

$$4(C^2 - BD)x_2, \quad -4(C^2 - BD)x_1, \quad -ABx_1 - 2ACx_2, \quad 2ACx_1 + ADx_2.$$

This plane contains g if and only if $C^2 - BD = 0$, i. e., if the derivative cubic is tangent to g . It will coincide with the plane tangent to S at this point, if further

$$-ABx_1 - 2ACx_2 = \omega x_2, \quad 2ACx_1 + ADx_2 = -\omega x_1,$$

where ω is a proportionality factor, or

$$ABx_1 + (2AC + \omega)x_2 = 0, \quad (2AC + \omega)x_1 + ADx_2 = 0,$$

whence follows $\omega = -AC$ or $-3AC$. We have therefore

$$x_1 : x_2 = -C : B = -D : C \quad \text{or} \quad x_1 : x_2 = C : B = D : C.$$

These points are harmonic conjugates with respect to the flecnodes.

Therefore, if the derivative cubic is tangent to g , there are two points of g whose tangent planes are the planes corresponding to them in the null-system of the cubic. These points and the flecnodes form a harmonic group on g . They never coincide with the complex points unless the ruled surface has a straight line directrix.

The planes, corresponding to a point of g in the null-system of the cubic and in the osculating complex, coincide if

$$\begin{aligned} C^2 - BD &= 0, \\ (AB - \omega p_{12})x_1 + 2ACx_2 &= 0, \\ 2ACx_1 + (AD - \omega p_{21})x_2 &= 0, \end{aligned}$$

where ω is a root of the quadratic

$$(41) \quad \omega^2 + 2u_{11}u_{22}(u_{11} - u_{22})^2(u_{11} + u_{22})\omega - 12u_{11}^3u_{22}^3(u_{11} - u_{22})^4 = 0,$$

neglecting again the case when S has a straight line directrix. These two points of g coincide if $(u_{11} + u_{22})^2 + 12u_{11}u_{22} = 0$.

More generally, if we write down the conditions that the same plane shall correspond to a point (x_1, x_2, x_3, x_4) in the osculating linear complex and in the complex of the cubic, we shall obtain as the locus of these points two straight lines, the directrices of the congruence common to the two complexes. These conditions are as follows, $x_1 \cdots x_4$ must satisfy the equations:

$$\begin{aligned} * + 4(C^2 - BD)x_2 + (AB - \omega p_{12})x_3 - 2ACx_4 &= 0, \\ (42) \quad -4(C^2 - BD)x_1 + * + 2ACx_3 - (AD - \omega p_{21})x_4 &= 0, \\ -(AB - \omega p_{12})x_1 - 2ACx_2 + * + A^2x_4 &= 0, \\ 2ACx_1 + (AD - \omega p_{21})x_2 - A^2x_3 + * &= 0, \end{aligned}$$

the vanishing of whose skew-symmetric determinant gives for ω the quadratic equation (41), which may also be written

$$(41a) \quad (AB - \omega p_{12})(AD - \omega p_{21}) + 4A^2(C^2 - BD) - 4A^2C^2 = 0.$$

Let ω_1 and ω_2 be the two roots of this equation. If we eliminate x_3 from the first two, x_4 from the last two equations of (42), if we make use of (41a) and assume that neither A nor $C^2 - BD$ is zero, we shall find

$$\begin{aligned} -(AB - \omega_k p_{12})x_1 - 2ACx_2 + A^2x_4 &= 0, \\ -4(C^2 - BD)x_2 - (AB - \omega_k p_{12})x_3 + 2ACx_4 &= 0 \end{aligned} \quad (k=1, 2),$$

whence

$$\begin{aligned} -2C(AB - \omega_k p_{12})x_1 - 4ABDx_2 + A(AB - \omega_k p_{12})x_3 &= 0, \\ (43) \quad -(AB - \omega_k p_{12})x_1 - 2ACx_2 + A^2x_4 &= 0 \end{aligned} \quad (k=1, 2),$$

the equations of the two directrices in simpler form than in (42).

A line joining the point $(x_1, x_2, 0, 0)$ of g to the point $(0, 0, x_1, x_2)$ of g' is a generator of the second kind on H . It is not difficult to see that it will intersect the directrix (43) if and only if

$$(44) \quad -(AB - \omega_k p_{12})^2 x_1^2 + 4A^2 BD x_2^2 = 0.$$

Hence, the two points, in which either of the directrices of the congruence common to the two complexes intersects the osculating hyperboloid, determine upon this hyperboloid two generators of the second set which are harmonic conjugates with respect to the flecnodal tangents.

It also follows easily that the two pairs thus obtained, one corresponding to each directrix, coincide only if $(u_{11} + u_{22})^2 + 12u_{11}u_{22} = 0$, i. e., if the directrices themselves coincide. Further if one of these pairs intersects g in the involute points, the same is true of the other pair, so that this can only happen if the directrices coincide. Finally, such a pair of generators of H can pass through the complex points only if S has a straight line directrix, or if S' is developable.

The line joining the points $(x_1, 0, x_3, 0)$ and $(0, x_1, 0, x_3)$ is a generator of the first set on H . The coördinates of an arbitrary point of this line are $(\lambda x_1, \mu x_1, \lambda x_3, \mu x_3)$. This line will therefore intersect one of the directrices of the congruence if x_1, x_3, λ, μ can be determined so as to satisfy the equations

$$\begin{aligned} \lambda [-2C(AB - \omega_k p_{12})x_1 + A(AB - \omega_k p_{12})x_3] - 4\mu ABDx_1 &= 0, \\ -\lambda(AB - \omega_k p_{12})x_1 + \mu(-2ACx_1 + A^2x_3) &= 0, \end{aligned}$$

which gives either $A = 0$, $AB - \omega_k p_{12} = 0$, or

$$(45) \quad 4(C^2 - BD)x_1^2 - 4ACx_1x_3 + A^2x_3^2 = 0.$$

The first two cases give either a surface S with a straight line directrix, or else a developable surface S' . Leaving these cases aside we notice that (45) does not contain ω_k so that if the line on H here considered intersects one of the directrices it intersects the other also. Combining this with our previous result, we see that the following theorem holds.

The four points in which the directrices of the congruence, common to the osculating linear complex and the linear complex of the derivative cubic, intersect the osculating hyperboloid can be grouped into two pairs, such that the line joining the members of each pair shall be a generator of the first set upon the hyperboloid. Upon this generator this pair of points, together with the intersections of the generator with the flecnodal tangents, form a harmonic group.

The plane, corresponding to a point $(x_1, x_2, 0, 0)$ of g in the null-system of the cubic, intersects the flecnodal tangents

$$\begin{aligned} f' &\text{ in the point } [ABx_1 + 2ACx_2, 0, 4(C^2 - BD)x_2, 0], \\ f'' &\text{ in the point } [0, 2ACx_1 + ADx_2, 0, 4(C^2 - BD)x_1]. \end{aligned}$$

The line joining these points is a generator of H , if either

$$A = 0, \quad \text{or} \quad C^2 - BD = 0, \quad \text{or} \quad Bx_1^2 - Dx_2^2 = 0.$$

Therefore, there exist in general two points on g , harmonic conjugates with

respect to the flecnodes, such that the planes, corresponding to them in the null-system of the derivative cubic, pass through a generator of the osculating hyperboloid. If the cubic is tangent to g the null-plane of any point of g contains a generator of H , viz, g itself. If $A = 0$ likewise, all points of g satisfy the condition of the theorem. Their null-planes all pass through g' .

§ 5. *Various theorems concerning the flecnode surface. The principal surface of the congruence Γ .*

Let us consider the planes which osculate the flecnode curve of S at P_y and P_z . We have of course $u_{12} = u_{21} = 0$. If x_1, x_2, x_3, x_4 are the coördinates of an arbitrary point of the plane osculating C_y at P_y , we have for the equation of this plane

$$\begin{vmatrix} x_1 & x_2 & x_3 & x_4 \\ y_1 & y_2 & y_3 & y_4 \\ y'_1 & y'_2 & y'_3 & y'_4 \\ y''_1 & y''_2 & y''_3 & y''_4 \end{vmatrix} = 0.$$

But

$$\begin{aligned} 2y'_k &= \rho_k - p_{11}y_k - p_{12}z_k, & 2z'_k &= \sigma_k - p_{21}y_k - p_{22}z_k, \\ -y''_k &= p_{11}y'_k + p_{12}z'_k + q_{11}y_k + q_{12}z_k, \text{ etc.} \end{aligned}$$

If we assume again $p_{11} = p_{22} = 0$, and substitute into the above equation, it becomes

$$\begin{vmatrix} x_1, & x_2, & \dots \\ y_1, & y_2, & \dots \\ \rho_1 - p_{12}z_1, & \rho_2 - p_{12}z_2, & \dots \\ p_{12}\sigma_1 + 2q_{12}z_1, & p_{12}\sigma_2 + 2q_{12}z_2, & \dots \end{vmatrix} = 0.$$

But if we introduce again our fundamental tetrahedron $P_y P_z P_\rho P_\sigma$, this becomes

$$(46a) \quad p_{12}x_2 + p_{12}^2x_3 - 2q_{12}x_4 = 0.$$

In the same way we find the equation of the plane osculating C_z at P_z to be

$$(46b) \quad p_{21}x_1 - 2q_{21}x_3 + p_{21}^2x_4 = 0.$$

To these equations must be added the conditions $u_{12} = u_{21} = 0$, if C_y and C_z are the two portions of the flecnode curve.

Let us assume that this is the case. The osculating planes at P_y and P_z intersect along a straight line, whose intersections with the osculating hyperboloid may now be found. If x_1, \dots, x_4 are the coördinates of one of these points of intersection, we find

$$\begin{aligned} (47) \quad p_{12}p_{21}^2x_2 &= -2q_{12}p_{21}x_1 - (p_{12}^2p_{21}^2 - 4q_{12}q_{21})x_3, \\ p_{12}p_{21}^2x_4 &= -p_{12}p_{21}x_1 + 2q_{21}p_{12}x_3, \end{aligned}$$

where the ratio of $x_1 : x_3$ is determined by the quadratic

$$(48) \quad (p_{12}^2 p_{21}^2 - 4q_{12} q_{21}) x_3^2 + 2(p_{12} q_{21} + p_{21} q_{12}) x_1 x_3 - p_{12} p_{21} x_1^2 = 0.$$

Therefore if $p_{12} q_{21} + p_{21} q_{12} = 0$, i. e., if $p_{12} p_{21} = \text{const.}$, the two generators of the first kind on H which pass through these points are harmonic conjugates with respect to g and g' . If

$$(49) \quad (p_{12} q_{21} - p_{21} q_{12})^2 + p_{12}^3 p_{21}^3 = 0,$$

the intersection of the two osculating planes is tangent to the hyperboloid. This latter property is obviously characteristic of a class of ruled surfaces, and can be expressed in invariant form.

If $\theta_4^3 \theta_9^2 + 16\theta_{10}^3 = 0$, the ruled surface S has the following characteristic property. The planes, which osculate the flecnode curve at the two points of its intersection with any generator, intersect in a line which is tangent to the osculating hyperboloid.

To prove this it is sufficient to notice that the invariant equation

$$(49a) \quad \theta_4^3 \theta_9^2 + 16\theta_{10}^3 = 0$$

reduces to (49) under our special assumptions. Further we notice:

If the independent variable is chosen so that $\theta_{10}/\theta_4 = \text{const.}$, the generator g' of the first derived surface is the harmonic conjugate of g with respect to the two generators of the same kind on H which are determined by the points in which the line of intersection of the two osculating planes intersects H .

The plane osculating the flecnode curve at P_y intersects the flecnode tangent f'' which passes through P_z in the point $(0, 2q_{12}, 0, p_{12})$. Similarly, the plane osculating C_z at P_z intersects f' in the point $(2q_{21}, 0, p_{21}, 0)$. Therefore, the line joining these points is a generator of H if and only if $p_{12} q_{21} - p_{21} q_{12} = 0$, i. e., if S belongs to a linear complex. In other words: the points in which the two planes, osculating the flecnode curve at its points of intersection with any generator, intersect the flecnode tangents are situated upon the same generator of the osculating hyperboloid if and only if the surface belongs to a linear complex.

To each of the two planes (46a) and (46b) corresponds a point in that plane by means of the osculating linear complex. These points have the coördinates

$$(-p_{12} p_{21}, -2q_{12}, 0, -p_{12}) \quad \text{and} \quad (2q_{21}, p_{12} p_{21}, p_{21}, 0).$$

We find that the line joining them intersects H in two points which form a harmonic group with the first two, if S belongs to a linear complex. It is tangent to H if (49) is satisfied. Therefore, if the two planes, osculating the flecnode curve at its two points of intersection with a generator, intersect in a line which is tangent to the osculating hyperboloid, the line joining the two points of these planes which correspond to them in the osculating linear complex is also tangent to the osculating hyperboloid, and conversely.

We have seen that, under the assumptions $u_{12} = u_{21} = p_{11} = p_{22} = 0$, the equations of the sheet F' of the flecnode surface assume the form (15). Let us denote by u_{ik} the quantities formed for this system according to the same law as are the quantities u_{ik} for the equations of S . Then we shall have

$$u_{12} = 0, \quad u_{21} = -4(q'_{11} - q'_{22}) + 8 \frac{(q_{11} - q_{22})q_{12}}{p_{12}}, \quad u_{11} - u_{22} = 4(q_{11} - q_{22}).$$

The curve C_y is a branch of the flecnode curve on F' as well as on S . The other branch is the locus of the point

$$u_{21}y - (u_{11} - u_{22})\rho.$$

Now if the transformation $\xi = \xi(x)$ is made, ρ is converted into

$$\bar{\rho} = \frac{1}{\bar{\xi}'}(\rho + \eta y), \quad \text{where} \quad \eta = \frac{\xi''}{\xi'}.$$

Therefore, if a transformation $\xi_1 = \xi_1(x)$ is made such that the derivative surface of S with respect to ξ_1 shall cut out upon F' the second branch of its flecnode curve, ξ_1 must be so chosen that

$$(50a) \quad \eta_1 = \frac{q'_{11} - q'_{22}}{q_{11} - q_{22}} - 2 \frac{q_{12}}{p_{12}}.$$

Similarly the second branch of the flecnode curve on F'' will be obtained by putting

$$(50b) \quad \eta_2 = \frac{q'_{11} - q'_{22}}{q_{11} - q_{22}} - 2 \frac{q_{21}}{p_{21}}.$$

The two surfaces of Γ thus obtained coincide only if S belongs to a linear complex, i. e., *the second branches of the flecnode curves on the two sheets of the flecnode surface of S correspond to each other only if S belongs to a linear complex.*

We have seen that the plane osculating C_y at P_y intersects f'' in the point $2q_{12}z + p_{12}\sigma$. The corresponding point on f' , i. e., the point obtained by finding the intersection of f' with the corresponding generator of H , is given by $2q_{12}y + p_{12}\rho$. We find a surface of Γ which intersects f' and f'' in these points by making a transformation of the independent variables for which $\eta = 2q_{12}/p_{12}$, or if we denote this special value of η by η_1 ,

$$\bar{\eta}_1 = 2 \frac{q_{12}}{p_{12}}.$$

If we now denote by η the expression

$$\eta = \frac{1}{2} \frac{q'_{11} - q'_{22}}{q_{11} - q_{22}},$$

we find

$$\eta = \frac{1}{2}(\eta_1 + \bar{\eta}_1),$$

and similarly

$$\eta = \frac{1}{2}(\eta_2 + \bar{\eta}_2).$$

This gives us a very important result. For, we have frequently made use of a normal form for our system of differential equations, in which $\theta_4 = \text{const.}$ But in order to have $\theta_4 = \text{const.}$, we must make precisely the transformation determined by η . On account of its importance, we will call the surface of Γ which is thus obtained, the *principal surface* of the congruence, and the curves in which it intersects the two sheets of the flecnodal surface of S their *principal curves*. We see then that the principal surface may be constructed as follows.

We consider the flecnodes P_y and P_z of g , the planes p_y and p_z osculating the flecnodal curve at these points, and the points P' and P'' upon the flecnodal tangents f' and f'' whose loci are the second branches of the flecnodal curves on the two sheets F' and F'' of the flecnodal surface. The plane p_y intersects f'' in a certain point to which corresponds a point on f' such that the line joining them is a generator of the osculating hyperboloid. This latter point together with P' constitute a pair, such that the harmonic conjugate of P_y with respect to it is the point in which the principal surface intersects f' . The intersection with f'' is found in the same way.

We might also say, that in this way there is determined, upon the generators of H , an involution whose double elements are g and the generator of the principal surface.

This construction is especially important for the complete geometrical interpretation of the covariant C_3 . For, the interpretation given in a former paper, (*Covariants*, p. 450), rests upon the use of the principal surface, since θ_4 is assumed to be constant, a fact which appears clearly from the detailed proof of the theorem there given, but which is not mentioned explicitly in its enunciations as it should have been.

By combining a number of our previous results with the notion of the principal surface, we obtain a number of theorems, which may be easily verified. They provide interpretations for the vanishing of certain invariants, and therefore furnish characteristic properties of certain families of ruled surfaces. I write them down without proof.

If $\theta_{15} = 0$, $\theta_4 \neq 0$, the principal surface is the harmonic conjugate of S with respect to the two ruled surfaces of Γ which cut out the second branches of the flecnodal curves on F' and F'' .

If $\theta_6\theta_4 - 9\theta_{10} = 0$, the principal surface intersects F' and F'' along asymptotic curves.

If $4\theta_4^5 - (\theta_4\theta_6 - 9\theta_{10})^2 = 0$, the principal surface is developable.

If $\theta_{15}^2 - \theta_9^2\theta_4^3 = 0$, the principal surface intersects one of the sheets of the flecnodal surface along the second branch of its flecnodal curve. It thus intersects both sheets if $\theta_{15} = \theta_9 = 0$.

PARIS, October 6, 1903.